

Trigonometry Through  
Wayfinding and Navigation Across  
the Pacific

# Trigonometry Through Wayfinding and Navigation Across the Pacific

Kamuela E Yong  
University of Hawai'i - West O'ahu

April 10, 2024

**Website:** [kamuelayong.com](https://www.kamuelayong.com)<sup>1</sup>

©2022 – 2024 Kamuela E Yong

This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. To view a copy of this license, visit [CreativeCommons.org](https://creativecommons.org/licenses/by-nc-sa/4.0)<sup>2</sup>

---

<sup>1</sup><https://www.kamuelayong.com/trigonometry/>

<sup>2</sup>[creativecommons.org/licenses/by-nc-sa/4.0](https://creativecommons.org/licenses/by-nc-sa/4.0)

For my family

# Acknowledgements

This project is partially supported by NSF grant #1853533 and the University of Hawai'i System. Additionally, I would like to thank BarbaraJean Kahawaii and Dr. Walter Kahumoku, III for their patience, encouragement, and sharing their knowledge. Bonnie Kahapea and the crew of Kānehūnāmoku Voyaging Academy; Kala Baybayan Tanaka and the crew from Hui o Wa'a Kaulua; Peia Patai and the crew from Te Puna Marama; the crew of Nā Kālai Wa'a; the Friends of Hōkūle'a and Hawai'iloa. Carina Chernisky and UHWO OER committee. I truly valued the discussions I have had with Vicente Diaz, Linda Furuto, Kekai Lee, and so many more.

# Preface

About this book: This book integrates navigation into trigonometry curriculum.

# Download PDF

Chapter 1 is available to down as a PDF at <https://www.kamuelayong.com/Trigonometry-Yong-Chapter-1.pdf>

# Core Standards

The curriculum in this book aligns with the Common Core State Standards: [thecorestandards.org](http://thecorestandards.org)<sup>3</sup>

- High School: Algebra
  - [Creating Equations](#)<sup>4</sup>
    - Create equations that describe numbers or relationships.
      - HSA.CED.A.2: Create equations in two or more variables to represent relationships between quantities; graph equations on coordinate axes with labels and scales. ([Section 2.3](#))
- High School: Functions
  - [Interpreting Functions](#)<sup>5</sup>
    - Analyze functions using different representations.
      - HSF.IF.C.7: Graph functions expressed symbolically and show key features of the graph, by hand in simple cases and using technology for more complicated cases. ([Section 2.1](#), [Section 2.2](#))
      - HSF.IF.C.7.e: Graph exponential and logarithmic functions, showing intercepts and end behavior, and trigonometric functions, showing period, midline, and amplitude. ([Section 2.1](#), [Section 2.2](#))
  - [Building Functions](#)<sup>6</sup>
    - Build a function that models a relationship between two quantities.
      - HSF.BF.A.1: Write a function that describes a relationship between two quantities. ([Section 2.3](#))
    - Build new functions from existing functions.
      - HSF.BF.B.3: Identify the effect on the graph of replacing  $f(x)$  by  $f(x) + k$ ,  $kf(x)$ ,  $f(kx)$ , and  $f(x + k)$  for specific values of  $k$  (both positive and negative); find the value of  $k$  given the graphs. Experiment with cases and illustrate an explanation of the effects on the graph using technology. Include recognizing even and odd functions from their graphs and algebraic expressions for them. ([Subsection 2.1.4](#), [Subsection 2.2.9](#))

---

<sup>3</sup><http://www.thecorestandards.org/>

<sup>4</sup><https://www.thecorestandards.org/Math/Content/HSA/CED/>

<sup>5</sup><https://www.thecorestandards.org/Math/Content/HSF/IF/>

<sup>6</sup><https://www.thecorestandards.org/Math/Content/HSF/BF/>



○ [Trigonometric Functions](#)<sup>7</sup>

- Extend the domain of trigonometric functions using the unit circle.
  - HSF.TF.A.1: Understand radian measure of an angle as the length of the arc on the unit circle subtended by the angle. ([Definition 1.2.18](#))
  - HSF.TF.A.2: Explain how the unit circle in the coordinate plane enables the extension of trigonometric functions to all real numbers, interpreted as radian measures of angles traversed counterclockwise around the unit circle. ([Subsection 1.3.2](#))
  - HSF.TF.A.3: Use special triangles to determine geometrically the values of sine, cosine, tangent for  $\pi/3$ ,  $\pi/4$  and  $\pi/6$ , and use the unit circle to express the values of sine, cosine, and tangent for  $x$ ,  $\pi + x$ , and  $2\pi - x$  in terms of their values for  $x$ , where  $x$  is any real number. ([Subsection 1.4.2](#), [Subsection 1.5.3](#))
  - HSF.TF.A.4: Use the unit circle to explain symmetry (odd and even) and periodicity of trigonometric functions. Model periodic phenomena with trigonometric functions. ([Subsection 1.5.4](#), [Subsection 1.5.7](#))
- Model periodic phenomena with trigonometric functions.
  - HSF.TF.B.5: Choose trigonometric functions to model periodic phenomena with specified amplitude, frequency, and midline. ([Subsection 2.1.4](#), [Section 2.3](#))
  - HSF.TF.B.6: Understand that restricting a trigonometric function to a domain on which it is always increasing or always decreasing allows its inverse to be constructed. ([Subsection 2.4.1](#))
  - HSF.TF.B.7: Use inverse functions to solve trigonometric equations that arise in modeling contexts; evaluate the solutions using technology, and interpret them in terms of the context. ([Exercise 2.1.5.46](#), [Exercise Group 2.1.5.47–50](#), [Exercise Group 2.2.10.37–42](#), [Exercise Group 2.2.10.43–52](#), [Example 2.4.18](#), [Example 2.4.19](#), [Example 2.4.17](#))
- Prove and apply trigonometric identities.
  - HSF.TF.C.8: Prove the Pythagorean identity  $\sin^2 \theta + \cos^2 \theta = 1$  and use it to find  $\sin \theta$ ,  $\cos \theta$ , or  $\tan \theta$  given  $\sin \theta$ ,  $\cos \theta$ , or  $\tan \theta$  and the quadrant of the angle. ([Definition 1.5.20](#))
  - HSF.TF.C.9: Prove the addition and subtraction formulas for sine, cosine, and tangent and use them to solve problems. ([Proof 3.2.1.1](#), [Example 3.2.10](#), [Proof 3.2.3.1](#))

• High School: Geometry

○ [Similarity, Right Triangles, and Trigonometry Define trigonometric ratios and solve problems involving right triangles.](#)<sup>8</sup>

- Define trigonometric ratios and solve problems involving right triangles

<sup>7</sup><https://www.thecorestandards.org/Math/Content/HSF/TF/>

<sup>8</sup><https://www.thecorestandards.org/Math/Content/HSG/SRT/>

- HSG.SRT.C.6: Understand that by similarity, side ratios in right triangles are properties of the angles in the triangle, leading to definitions of trigonometric ratios for acute angles. ([Definition 1.4.1](#))
- HSG.SRT.C.7: Explain and use the relationship between the sine and cosine of complementary angles. ([Definition 1.4.7](#), [Remark 1.4.9](#))
- HSG.SRT.C.8: Use trigonometric ratios and the Pythagorean Theorem to solve right triangles in applied problems. ([Subsection 1.4.6](#))
- Apply trigonometry to general triangles
  - HSG.SRT.D.9: Derive the formula  $\text{Area} = \frac{1}{2}ab \sin(C)$  for the area of a triangle by drawing an auxiliary line from a vertex perpendicular to the opposite side.
  - HSG.SRT.D.9: Prove the Laws of Sines and Cosines and use them to solve problems.
  - HSG.SRT.D.9: Understand and apply the Law of Sines and the Law of Cosines to find unknown measurements in right and non-right triangles (e.g., surveying problems, resultant forces).
- [Circles](#)<sup>9</sup>
  - Find arc lengths and areas of sectors of circles
    - HSG.C.B.5: Derive using similarity the fact that the length of the arc intercepted by an angle is proportional to the radius, and define the radian measure of the angle as the constant of proportionality; derive the formula for the area of a sector. ([Definition 1.2.18](#), [Theorem 1.2.25](#), [Definition 1.2.28](#))

---

<sup>9</sup><https://www.thecorestandards.org/Math/Content/HSG/C/>

# Contents

<b>Acknowledgements</b>	<b>v</b>
<b>Preface</b>	<b>vi</b>
<b>Download PDF</b>	<b>vii</b>
<b>Core Standards</b>	<b>viii</b>
<b>1 Trigonometric Functions</b>	<b>1</b>
1.1 Pacific Island Navigation . . . . .	1
1.2 Angles and Their Measure . . . . .	12
1.3 Unit Circle . . . . .	33
1.4 Right Triangle Trigonometry. . . . .	47
1.5 Trigonometric Functions of Any Angles . . . . .	65
<b>2 Graphs of the Trigonometric Functions</b>	<b>88</b>
2.1 Graphs of the Sine and Cosine Functions . . . . .	88
2.2 Graphs of Other Trigonometric Functions . . . . .	113
2.3 Sinusoidal Curve Fitting and Graphical Analysis . . . . .	135
2.4 Inverse Trigonometric Functions . . . . .	155
<b>3 Analytic Trigonometry</b>	<b>178</b>
3.1 Trigonometric Identities . . . . .	178
3.2 Addition and Subtraction Formulas . . . . .	183
3.3 Double-Angle and Half-Angle Formulas . . . . .	197
3.4 Product-to-Sum and Sum-to-Product Formulas . . . . .	209
3.5 Basic Trigonometric Equations . . . . .	214
3.6 Trigonometric Equations - Advanced Techniques . . . . .	219
<b>Back Matter</b>	
<b>Glossary (Math)</b>	<b>224</b>

*CONTENTS*

xii

**Glossary (Canoe and Wayfinding)**

**225**

**Index**

**226**

# Chapter 1

## Trigonometric Functions

### 1.1 Pacific Island Navigation

Pacific Islanders have been navigating across long distances in the Pacific for centuries, even before the use of magnetic compasses or modern instruments. They relied on observations of celestial bodies, such as stars and the sun, as well as natural elements like ocean patterns, winds, bird behaviors, and other environmental cues to determine their position relative to known landmarks such as islands, reefs, and continents. Over time, much of this traditional navigational knowledge was lost in many parts of the Pacific. However, some islands, particularly in Micronesia and on Taumako Island in the Solomon Islands, managed to preserve the art and science of traditional navigation. These places continued to uphold the practice, teaching new generations to build ocean-going canoes and develop navigational skills based on profound knowledge of the natural world.

#### 1.1.1 Micronesia


Navigators in Micronesia utilize the *paafu* mat or map (shown in [Figure 1.1.1](#)). It is often misunderstood and misinterpreted as a Star “Compass” due to its use of stars and constellations for direction finding. However, *paafu* serves a different purpose and is not equivalent to the cardinal directionality marked by compasses (North, South, East, and West). Instead, it is a learning and teaching tool designed to teach the locational positions of islands, locales, or canoes relative to other places. This is achieved by observing the rising and setting points of stars and constellations, which act as markers for different locations.



**Figure 1.1.1** Paafu mat or map - Photo courtesy of Kānehūnāmoku Voyaging Academy

A constellation is a cluster of stars whose shapes and meanings reflect and carry cultural significance. In modern society, a well-known cluster of stars in the southern hemisphere, shaped like a cruciform, is commonly referred to as the “Southern Cross.” However, for the Polowatese and other islanders from the Central Carolines region in the western Pacific islands, this same constellation resembles the triggerfish, and so it is named accordingly.

In the Central Carolines, the general location in the celestial sky where stars appear to rise after sundown is referred to as “*tan*.” This term is often mistakenly translated as “east” due to the modern association of stars (like the sun) with “rising” in the east. However, it’s important to note that “*tan*” means “rising” and not “eastward.”



**Figure 1.1.2** The paafu, or Micronesian Star Compass. Stars are identified using the Polowat dialect of the Chuukese language as it is used by members of the Werieng School of navigation.

Figure 1.1.2 orients the cardinal direction known as “east” at the top of the page, and so the top half of this diagram is also identified as “tan” – where stars appear to rise. The diagram illustrates the apparent path of stars across the sky each night and day (though most stars are not visible during the day) and throughout a year. In this system, the rising and presence of specific stars mark months, and these same stars will eventually set at a point horizontally opposite to where they rose. This point corresponds to the cardinal direction known as “west,” which is referred to as “tolon” – the area where the stars “set” or go down.

In the paafu “map” shown in Figure 1.1.2, a canoe is placed at the center, and the star called Mailap (Altair) marks due east. The time and location when Mailap rises are referred to as “Tan Mailap” (rising Mailap), while the time and place it sets are referred to as “Tolon Mailap” (setting Mailap). The map’s orientation places east, or the rising points of the navigation or paafu stars, at the top of the circle, and west, or the setting points of these stars, at the bottom of the circle. As a map, paafu uses the rising and setting points of stars to mark places around a given locale, which is placed at the center of the circle. Table 1.1.3 displays the star and constellation names, provided in both Polowat and according to the International Astronomical Union, listed in the order of their rising during the third week of March in Polowat.

**Table 1.1.3 Star and constellation names in Polowat**

Polowat	International Astronomical Union
Wenenwenenfuwmaket	Polaris (always above the horizon)
Tan Mwarikar [Mahrah-ker]	Pleiades aka Seven Sisters
Tan Un [Oon]	Aldebaran
Tan Uliul [Ooh-lee-ool]	Orion's Belt
Tan Harapwel [Ah-rah-pwol]	Gamma Corvus
Tan Mailapenefang [My Lap in a Fang]	Beta Ursa Minor in Big Dipper
Tan Up [Oop]	Crux or Southern Cross at Rising
Machemeas [Matche-may-ess]	Crux or S. Cross at 45° <sup>1</sup>
Tan Welo [Well-Ah]	Alpha Ursa Major in Big Dipper
Wenenwenenup [Wehneh wehnen Oop]	Crux or S. Cross at Meridian or upright
Tan Tumor [Two More]	Antares or Scorpio's tail
Tan Maharuw [Maa-Haa-Roo]	Shaula or Scorpio's stinger
Tan Mol [Mohl]	Vega
Tan Mailap [My Lap]	Altair
Tan Paiefung [Pie Efung]	Gamma Aquila
Tan Paior [Pie Or]	Beta Aquila
Tan Ukinik [Icky Nick]	Cassiopeia

Paafu can also be used to identify the direction in which moving objects, such as canoes, or creatures like birds, fish, and humans, are heading or coming from. This version of paafu utilizes the Polowat dialect of the Chuukese language, as used by members of the Weriyeeng School of navigation.


### 1.1.2 Hawai'i

With the aim of reviving wayfinding in Hawai'i, Nainoa Thompson journeyed to the island of Satawal in the Federated States of Micronesia to learn from master navigator Mau Piailug, affectionately known as Papa Mau. Using this knowledge, Thompson adopted the paafu method, leading to the creation of the Hawaiian Star Compass, also referred to as the *Kūkuluokalani* (Figure 1.1.4).

In the star compass, featuring the figure of an 'iwa or great frigatebird at its center, Thompson divides the visual horizon into 32 equidistant points around a circle, referred to as houses. Each house in the Hawaiian Star Compass represents a specific space on the horizon (11.25°) where celestial bodies such as the sun, stars, moon, and planets rise and set. In the same way that we use addresses to locate homes, each celestial body has its own address represented by these houses.

<sup>1</sup>The "Tan" prefix is not used for this position, because it is no longer rising





**Figure 1.1.4** Hawaiian Star Compass, also known as the Kūkuluokalani.

The four cardinal points align with particular houses. Stars rise from the horizon called *Hikina* (“To Arrive”) or East and set on the horizon called *Komohana* (“To Enter”) or West. If you face *Komohana* (West) with your back towards *Hikina* (East), your right will point towards *‘Ākau* (“Right”) or North, and your left will point towards *Hema* (“Left”) or South. The Hawaiian Star Compass is oriented with North at the top.


The star compass is divided into four quadrants, each named after winds in Hawai‘i. *Ko‘olau* is the Northeast quadrant named for the trade winds; *Ho‘olua* is the Northwest quadrant, *Kona* is the Southwest quadrant; and *Malanai* is the Southeast quadrant.

Each house on the star compass is given a name. The corresponding houses in the east and west share the same name. Starting from the east or west and moving northwards and southwards, the first house on either side of *Hikina* (East) and *Komohana* (West) is called *Lā* (Sun). It is followed by *‘Āina* (Land), *Noio* (Tern), *Mamu* (Bird), *Nālanai* (Heavens), *Nā Leo* (Voices), and *Haka* (Empty). The 32 houses in the Hawaiian Star Compass correspond to the points of the 32-wind compass rose (Table 1.1.5).

**Table 1.1.5** The houses of the Hawaiian Star Compass and the corresponding points on the 32-wind compass rose.

Star Compass	32 Point Compass		Star Compass	32 Point Compass	
House	Symbol	Name	House	Symbol	Name
Hikina	E	East	Komohana	W	West
Lā Ko‘olau	EbN	East by North	Lā Kona	WbS	West by South
‘Āina Ko‘olau	ENE	East-northeast	‘Āina Kona	WSW	West-southwest
Noio Ko‘olau	NEbE	Northeast by East	Noio Kona	SWbW	Southwest by West
Manu Ko‘olau	NE	Northeast	Manu Kona	SW	Southwest
Nālani Ko‘olau	NEbN	Northeast by North	Nālani Kona	SWbS	Southwest by South
Nā Leo Ko‘olau	NNE	North-northeast	Nā Leo Kona	SSW	South-southwest
Haka Ko‘olau	NbE	North by east	Haka Kona	SbW	South by West
‘Ākau	N	North	Hema	S	South
Haka Ho‘olua	NbW	North by West	Haka Malanai	SbE	South by East
Nā Leo Ho‘olua	NNW	North-northwest	Nā Leo Malanai	SSE	South-southeast
Nālani Ho‘olua	NWbN	Northwest by North	Nālani Malanai	SEbS	Southeast by South
Manu Ho‘olua	NW	Northwest	Manu Malanai	SE	Southeast
Noio Ho‘olua	NWbW	Northwest by West	Noio Malanai	SEbE	Southeast by East
‘Āina Ho‘olua	WNW	West-northwest	‘Āina Malanai	ESE	East-southeast
Lā Ho‘olua	WbN	West by North	Lā Malanai	EbS	East by South

Celestial bodies move along parallel paths across the sky from East to West, rising and setting in the same house, remaining in its hemisphere. For example, if a star arrives in the Ko‘olau (northeastern) quadrant in the star house ‘Āina, it will arc overhead, staying in the northern hemisphere, and enter the horizon in the same house it arrived in, ‘Āina, but in the Ho‘olua (northwestern) quadrant (see Figure 1.1.6). Similarly, if a star arrives in the Malanai (southeastern) quadrant in the house Lā, it will remain in the southern hemisphere as it arcs overhead and enters the horizon in the house Lā in the Kona (southwestern) quadrant.




**Figure 1.1.6** In the celestial sphere, stars rise in the east, arc across the sky, and set in the west. Each star will both rise and set in the same house.

The star compass also serves as a guide for determining direction based on wind and ocean swells. As the wind and swells move, they intersect the star compass diagonally. For example, if a wind blows from the house Noio in the Ko‘olau (northeast) quadrant, it will blow in the direction of the Kona (southwest) quadrant and eventually exit in the same house, Noio.

Observations play a key role in determining direction using the star compass. At night, Thompson relies on approximately 220 stars, memorizing where they

rise and set on the horizon to navigate. During the day, we can use the sun's position on the horizon to gauge direction, but this method is only effective when the sun is near the horizon at sunrise and sunset. Alternatively, one can memorize the wind and wave directions, checking for any changes between sunrise and sunset to establish their current direction.

The canoe itself can serve as a compass, as shown in Figure 1.1.7. From the navigator's seat on either corner of the stern (back) of the deck, you can observe features like the rising sun and mark its position on the Star Compass located on the canoe. It's essential to note that the locations of the houses on this Star Compass are in relation to the canoe, not to a fixed map. For instance, only when the canoe is pointed towards the north will Hikina (East) be the house to the right. Depending on the canoe's orientation, at other times, Hikina may appear several houses further up the deck.



**Figure 1.1.7** The deck of a canoe can be used as a compass to help crew and navigators.

### 1.1.3 Marshall Islands


The Marshallese people use stick charts as navigational tools. These stick charts are constructed using a lattice-like structure made from curved and straight sticks, typically formed by tying together the midribs of coconut fronds. The curved sticks represent the islands and how they bend and refract the

ocean swells, while the straight sticks symbolize the major wave patterns in the surrounding waters.

The shells placed on the sticks indicate the relative locations of islands within the Marshall Islands archipelago. These shells serve as markers, helping navigators remember the positions of specific islands along their voyages.

Each stick chart is unique to its creator, reflecting their individual knowledge and experiences. The personalization of the stick charts allows navigators to develop a deep understanding of the ocean currents, wave patterns, and island locations in their specific region.

The stick charts serve as mental maps or navigational aids, allowing experienced navigators to visualize and recall the complex information while on their journeys. Navigators would memorize the stick charts, internalizing the knowledge embedded within them, enabling them to navigate the open ocean.




**Figure 1.1.8** Marshallese stick chart.

#### 1.1.4 Elsewhere in the Pacific


In addition to the star compass, many cultures across the Pacific use a *wind compass*. Similar to the star compass, the wind compass is also a mental construct.

Other Pacific Island cultures have also adapted the modern Hawaiian Star Compass to their languages, as illustrated in [Figure 1.1.9](#).



(a) Cook Islands.

(b) Māori.



(c) Sāmoan.

**Figure 1.1.9** Examples of Star Compasses across the Pacific.

**1.1.5 Exercises**

1. Who developed the Hawaiian Star Compass?  
**Answer.** Nainoa Thompson
2. The Hawaiian Star Compass was based on the Micronesian Star Compass, known as the *paafu*. Who shared the *paafu* with the Hawaiians?  
**Answer.** Mau Pailug or Papa Mau
3. According to the Hawaiian Star Compass, what is the name for
  - (a) North  
**Answer.** ‘Ākau
  - (b) East  
**Answer.** Hikina
  - (c) South  
**Answer.** Hema
  - (d) West  
**Answer.** Komohana

4. What is the Hawaiian name for winds in the

(a) Northeast quadrant

**Answer.** Ko'olau

(b) Southeast quadrant

**Answer.** Malanai

(c) Southwest quadrant

**Answer.** Kona

(d) Northwest quadrant

**Answer.** Ho'olua

**Exercise Group.** For each direction, identify the Hawaiian names of the corresponding house and quadrant in the Hawaiian Star Compass

5. Northwest by North

**Answer.** Nālani Ho'olua

7. North-northwest

**Answer.** Nā Leo Ho'olua

9. South by West

**Answer.** Haka Kona

11. Southwest by West

**Answer.** Noio Kona

13. South-southeast

**Answer.** Nā Leo Malanai

6. East-northeast

**Answer.** 'Āina Ko'olau

8. Southeast

**Answer.** Manu Malanai

10. East by South

**Answer.** Lā Malanai

12. Northeast by North

**Answer.** Nālani Ko'olau

14. West-southwest

**Answer.** 'Āina Kona

**Exercise Group.** Identify the corresponding point on the 32-wind compass for each house on the Hawaiian Star Compass

15. Nā Leo Kona

**Answer.** South-southwest

17. 'Āina Ho'olua

**Answer.** West-northwest

19. Manu Ko'olau

**Answer.** Northeast

21. Nālani Kona

**Answer.** Southwest by South

23. Noio Ho'olua

**Answer.** Northwest by West

16. 'Āina Malanai

**Answer.** East-southeast

18. Lā Ko'olau

**Answer.** East by North

20. Haka Malanai

**Answer.** South by East

22. Haka Ho'olua

**Answer.** North by West

24. Noio Ko'olau

**Answer.** Northeast by East

25. The winter solstice in the southern hemisphere occurs around June 22. It is the time when the sun is at its lowest elevation in the sky, resulting in the shortest daylight of the year. During the winter solstice, the sun rises from its northernmost position, 'Āina Ko'olau. In what house does the sun set during the winter solstice in the southern hemisphere?

**Answer.** 'Āina Ho'olua

26. The winter solstice in the northern hemisphere occurs around December 22 when the sun rises from its southernmost position, 'Āina Malanai. In which house does the sun set during the winter solstice in the northern hemisphere?

**Answer.** 'Āina Kona

- 27.** Wind is coming from Nā Leo Kona. In what direction is the wind blowing?


**Answer.** Nā Leo Ko'olau

- 28.** Current is coming from Noio Ho'olua. In what direction is the current heading?

**Answer.** Noio Malanai


## 1.2 Angles and Their Measure

One method people use to identify their position is by looking at the *latitude*. These imaginary lines form circles around the Earth and run parallel to the Equator. The latitude of a place is defined as the angle between a line drawn from the center of the Earth to that point and the equatorial plane. For any point in the Northern Hemisphere, a navigator can measure their latitude by determining the angle that *Hōkūpa‘a* (also known as *Kūmau*, *Wuli wulifasmughet*, *Fuesemagut*, North Star, or Polaris) makes with the horizon.




During voyages, knowing the correct angles can make the difference between reaching your destination or missing it. Navigators carefully observe angles on the Hawaiian Star Compass to determine the entry and exit points of celestial bodies in the sky, as well as the direction of wind and current. In this section, we will explore the properties of angles and their measure.

**Definition 1.2.1** A **ray** is a part of a line that begins at a point  $O$  and extends in one direction.



◇

**Definition 1.2.2** We can create an **angle**,  $\theta$ , by rotating rays. First, we begin with two rays lying on top of each other and beginning at  $O$ . We let one ray be fixed and will rotate the second ray about the point  $O$ . The ray that is fixed is called the **initial side** and the ray that is rotated is called the **terminal side**.



◇


**Remark 1.2.3** Angles are often measured using Greek letters. The commonly used Greek letters include  $\theta$ ,  $\phi$ ,  $\alpha$ ,  $\beta$ , and  $\gamma$ .

### 1.2.1 Degree

The **measure** of an angle is the amount of rotation from the initial side to the terminal side. One unit of measuring angles is the degree. One **degree**, denoted by  $1^\circ$ , is  $\frac{1}{360}$  of a complete circular revolution, so one full revolution is  $360^\circ$ .



The Hawaiian Star Compass consists of 32 houses, each spanning  $11.25^\circ$  ( $\frac{360^\circ}{32}$ ). Assuming due East corresponds to  $0^\circ$  and the center of the House of Hikina points due East, the border between Hikina and Lā Ko'olau will be half the angle of the house,  $5.625^\circ$  ( $\frac{11.25^\circ}{2}$ ). The angles for the other boundaries on the Hawaiian Star Compass are shown in Figure 1.2.4.



**Figure 1.2.4** The Star Compass with the angles indicating the boundaries for each House.

Although decimals are commonly used to represent fractional parts of a degree, traditionally, degrees were represented in minutes and seconds. One **minute** or **arc minute**, denoted as  $1'$ , is equal to  $\frac{1}{60}$  degrees, and one **second** or **arc second**, denoted as  $1''$ , is equal to  $\frac{1}{60}$  minutes.

**Remark 1.2.5** Conversion Between Degree, Minutes, and Seconds.

$$\begin{aligned}
 1^\circ &= 60' & 1' &= \left(\frac{1}{60}\right)^\circ & 1'' &= \left(\frac{1}{3600}\right)^\circ \\
 1^\circ &= 3600'' & 1' &= 60'' & 1'' &= \left(\frac{1}{60}\right)'
 \end{aligned}$$

**Example 1.2.6** Convert angle from decimal degrees to degrees/minutes/seconds. In the Star Compass (Figure 1.2.4), the angle between the houses Manu Ho'olua (northwest) and Noio Ho'olua (northwest by west) measures  $140.625^\circ$ . Represent this angle in degrees, minutes, and seconds.

**Solution.** First we will convert  $0.625^\circ$  to minutes using the conversion  $1^\circ =$

60',

$$0.625^\circ = 0.625^\circ \cdot \frac{60'}{1^\circ} = 37.5'$$

Since  $1' = 60''$ , we can convert 0.5' to seconds:  $0.5' = 0.5' \cdot \frac{60''}{1'} = 30''$ .

So  $140.625^\circ = 140^\circ 37' 30''$ .  $\square$

**Example 1.2.7 Convert angle from degrees/minutes/seconds to decimal degrees.** Convert  $263^\circ 24' 45''$  to decimal degrees.

**Solution.** We will first convert  $24'$  and  $45''$  to degrees.


$$24' = 24 \cdot 1' = 24 \cdot \left(\frac{1}{60}\right)^\circ = 0.4^\circ$$

and

$$45'' = 45 \cdot 1'' = 45 \cdot \left(\frac{1}{3600}\right)^\circ = 0.0125^\circ$$


So  $263^\circ 24' 45'' = 263^\circ + 24' + 45'' = 263^\circ + 0.4^\circ + 0.0125^\circ = 263.4125^\circ$   $\square$

**Definition 1.2.8** If an angle is drawn on the  $xy$ -plane, and the vertex is at the origin, and the initial side is on the positive  $x$ -axis, then that angle is said to be in **standard position**. If the angle is measured in a counterclockwise rotation, the angle is said to be a **positive angle**, and if the angle is measured in a clockwise rotation, the angle is said to be a **negative angle**.



$\diamond$

**Definition 1.2.9** When an angle is in standard position, the terminal side will either lie in a quadrant or it will lie on the  $x$ -axis or  $y$ -axis. An angle is called a **quadrantal angle** if the terminal side lies on  $x$ -axis or  $y$ -axis. The two axes divide the plane into four **quadrants**. In the Cartesian plane, the four quadrants are Quadrant I, II, III, and IV. The corresponding quadrants of Star Compass are Ko'olau (NE), Ho'olua (NW), Kona (SE), and Malanai (SW).



◇

**Definition 1.2.10 Coterminal angles** are angles in standard position that have the same initial side and the same terminal side. Any angle has infinitely many coterminal angles because each time we add or subtract  $360^\circ$  from it, the resulting angle has the same terminal side. ◇

**Example 1.2.11 Coterminal angles.**  $90^\circ$  and  $450^\circ$  are coterminal angles since  $450^\circ - 360^\circ = 90^\circ$ . □

To determine the quadrant in which an angle lies, add or subtract one revolution ( $360^\circ$ ) until you obtain a coterminal angle between  $0^\circ$  and  $360^\circ$ . The quadrant where the terminal side lies is the quadrant of the angle. Quadrantal angles do not lie in any quadrant.

**Example 1.2.12 Determine the corresponding house and quadrant in the Star Compass.** Determine the quadrant in which each angle lies and name the corresponding House and quadrant in the Star Compass (Figure 1.2.4).

(a)  $140^\circ$

**Solution.** Since  $90^\circ < 140^\circ < 180^\circ$ ,  $140^\circ$  lies in Quadrant II, or Manu Ho'olua.

(b)  $-770^\circ$

**Solution.** Since  $-770^\circ < 0^\circ$ , we first add  $3 \times 360$  to  $-770^\circ$  to obtain an angle  $0^\circ$  and  $360^\circ$ ,

$$-770^\circ + 3 \times 360^\circ = 310^\circ$$

So  $310^\circ$  and  $-770^\circ$  are coterminal. Since  $270^\circ < 310^\circ < 360^\circ$ ,  $310^\circ$  lies in Quadrant IV, or Manu Malanai.

(c)  $923^\circ$

**Solution.** Since  $923^\circ > 360^\circ$  we begin by subtracting  $2 \times 360^\circ$

$$923^\circ - 2 \times 360^\circ = 203^\circ$$


So  $203^\circ$  and  $923^\circ$  are coterminal. Since  $180^\circ < 203^\circ < 270^\circ$ ,  $923^\circ$  lies in Quadrant III or 'Aina Kona.

□

**Example 1.2.13** Determine the corresponding quadrant given its location in the Star Compass. What is the corresponding quadrant for Nālani Kona?


**Solution.** Locating Nālani Kona in the Star Compass, we see it is in Quadrant III.  $\square$

**Definition 1.2.14** A **central angle** is a positive angle formed at the center of a circle by two radii.




$\diamond$

**Remark 1.2.15** **Heading and Azimuth.** In navigation, the direction a wa'a is pointed towards is referred to as the **heading**. Unlike in trigonometry, where it is conventional to define an angle in standard position, i.e.,  $0^\circ$  lies along the positive  $x$ -axis, in navigation, North corresponds to a heading of  $0^\circ$  and positive angles are measured in a clockwise rotation (see [Figure 1.2.16](#)).



**Figure 1.2.16** The cardinal directions for headings are as follows:  $0^\circ$  (or  $360^\circ$ ) points north,  $90^\circ$  points east,  $180^\circ$  points south, and  $270^\circ$  points west.

The Star Compass can now be presented in terms of heading angles, as demonstrated in [Figure 1.2.17](#).



**Figure 1.2.17** The Star Compass is presented in terms of heading, with the angles indicating the boundaries for each House.

In astronomy and navigation, the position of a celestial body as it rises or sets on the horizon can be measured using the **azimuth**, which indicates the direction of celestial objects relative to an observer's position. Similar to heading, azimuth starts from the north and increases clockwise.

In navigation, "heading" typically denotes the direction an object like a canoe or wind is pointed, whereas "azimuth" pertains to the angular measurement of celestial bodies on the horizontal plane. Both "heading" and "azimuth" measure angles in degrees, beginning from north and progressing clockwise. Unless specified otherwise to use the heading or azimuth angle (in which case, refer to [Figure 1.2.17](#)), this book will use [Figure 1.2.4](#) for the angles of the Star Compass.


## 1.2.2 Radian

Another way to measure an angle is with *radians*, which measure the the arc of a circle that is formed from an angle.

**Definition 1.2.18 Definition of a Radian.** The **radian measure** of a central angle in a circle is the ratio of the length of the arc on a circle subtended by the angle to the radius. If  $r$  is the radius of the circle,  $\theta$  is the angle, and  $s$  is the arc length, then we have the following


$$\theta = \frac{s}{r}$$

A radian is abbreviated by **rad**.




◇

The measure of a central angle obtained when the length of the arc is also equal to the radius,  $r$ , is called *one radian* (1 rad). Similarly, if  $\theta = 2$  rad, then the arc length equals  $2r$ .



The circumference of a circle is  $C = 2\pi r$ . This means that the circumference is  $2\pi \approx 6.28$  times the radius. Consequently, if we were to use a piece of string with the length of the radius, we would need six pieces of string plus a fractional piece of the string, as shown in [Figure 1.2.19](#).



**Figure 1.2.19** One rotation of the unit circle is  $2\pi \approx 6.28$  radians.

**Remark 1.2.20 Relationships Between Degrees and Radians.** If a circle with radius 1 is drawn, it has  $360^\circ$ , and the full arc length is the circumference, which is  $2\pi$ . Therefore, the relationship between degrees and radians is:

$$360^\circ = 2\pi \text{ radian, or } 180^\circ = \pi \text{ radian}$$

$$1 \text{ radian} = \frac{180^\circ}{\pi}$$

$$1^\circ = \frac{\pi}{180} \text{ radian}$$

**Remark 1.2.21 Converting Between Degrees and Radians.**

1. To convert degree to radians, multiply by  $\frac{2\pi \text{ radians}}{360^\circ}$  or  $\frac{\pi \text{ radian}}{180^\circ}$
2. To convert radians to degrees, multiply by  $\frac{360^\circ}{2\pi \text{ radians}}$  or  $\frac{180^\circ}{\pi \text{ radian}}$

**Example 1.2.22** Express  $45^\circ$  in radians.

**Solution.**  $45^\circ = 45^\circ \left( \frac{2\pi \text{ radians}}{360^\circ} \right) = \frac{\pi}{4} \text{ radians}$  □

**Example 1.2.23** Express  $\frac{5\pi}{6}$  in degrees.

**Solution.**  $\frac{5\pi}{6} \text{ rad} = \frac{5\pi}{6} \text{ rad} \left( \frac{360^\circ}{2\pi \text{ rad}} \right) = 150^\circ$  □

Using this method, we can obtain [Table 1.2.24](#) of common angles used in trigonometry and the corresponding radian and degree measures.

**Table 1.2.24 Commonly Used Angles in Trigonometry: Degrees and Radians**

<i>Radians</i>	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$
<i>Degrees</i>	0°	30°	45°	60°	90°	120°	135°	150°	180°
<i>Radians</i>	$\pi$	$\frac{7\pi}{6}$	$\frac{5\pi}{4}$	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	$\frac{7\pi}{4}$	$\frac{11\pi}{6}$	$2\pi$
<i>Degrees</i>	180°	210°	225°	240°	270°	300°	315°	330°	360°

### 1.2.3 Arc Length

Recall that the definition of a radian is the ratio of the arc length to the radius of a circle,  $\theta = \frac{s}{r}$ . By rearranging this formula, we can obtain a formula for the arc length of a circle.

**Theorem 1.2.25** *In a circle of radius  $r$ , the arc length,  $s$ , subtended by a central angle (in radians),  $\theta$ , is*

$$s = r\theta$$


*If  $\theta$  is given in degrees, then  $s = 2\pi r \cdot \left(\frac{\theta}{360^\circ}\right)$ .*

**Example 1.2.26** Find the length of an arc of a circle with radius 10 cm subtended by an angle of 2 radians.

**Solution.** Using the arc formula we get  $s = 10\text{cm} \cdot 2\text{rad} = 20\text{cm}$ . □

**Example 1.2.27** Kiritimati, also known as Christmas Island, is an atoll in the Republic of Kiribati. Kiritimati's location west of the International Date Line makes it one of the first places in the world to welcome the New Year, while Hawai'i is one of the last places. Although Kiritimati and Moloka'i share the same longitude at  $157^\circ 12'$  west (meaning Moloka'i is directly north of Kiritimati), both islands are 24 hours apart. For example, if the time on O'ahu is 3:00 pm on Thursday, then at that same moment it is 3:00 pm on Friday in Kiritimati. Find the distance between Kiritimati ( $1^\circ 45'$  north latitude) and Moloka'i ( $21^\circ 08'$  north latitude). Assume the radius of Earth is 3,960 miles and that the central angle between the two islands is the difference in their latitudes.





**Solution.** The measure of the central angle between the two islands is

$$\begin{aligned}\theta &= 21^\circ 08' - 1^\circ 45' \\ &= 19^\circ 23' \\ &= 19^\circ + \frac{23}{60}^\circ \\ &\approx 19.3833^\circ\end{aligned}$$

To find the distance, we use [Theorem 1.2.25](#) to find the arc length:

$$s = 2\pi r \cdot \left(\frac{\theta}{360^\circ}\right) \approx 2\pi \cdot (3,960 \text{ miles}) \frac{19.3833^\circ}{360^\circ} \approx 1,340 \text{ miles}$$

So the distance between Kiritimati and Moloka'i is approximately 1,340 miles.  $\square$

### 1.2.4 Area of a Sector of a Circle

**Definition 1.2.28 Area of a Sector.** The **area of the sector** of a circle of radius  $r$  formed by a central angle of  $\theta$  is

$$\begin{aligned}A &= \frac{\theta}{360^\circ} \cdot \pi \cdot r^2, \text{ when } \theta \text{ is in degrees} \\ A &= \frac{1}{2}r^2\theta, \text{ when } \theta \text{ is in radians}\end{aligned}$$

$\diamond$

Notice the ratio  $\frac{\theta}{360^\circ}$  is the proportion of the angle  $\theta$  (in degrees) to one complete circle. Additionally, the circumference of a circle is given by  $2\pi r$ . Therefore, the arc length is simply the proportion of the central angle to the whole circle multiplied by the circumference of the circle.

$$s = \text{arc length} = (\text{proportion of circle}) \cdot (\text{circumference}) = \left(\frac{\theta}{360^\circ}\right) \cdot (2\pi r)$$

Similarly, the area of a circle is given by  $\pi r^2$ . So the area of sector is the

proportion of the central angle to the whole circle multiplied by the area of the circle.

$$A = \text{area of sector} = (\text{proportion of circle}) \cdot (\text{area of circle}) = \left(\frac{\theta}{360^\circ}\right) \cdot (\pi r^2)$$


**Theorem 1.2.29** *Given a circle of radius  $r$  formed by a central angle of  $\theta$ , then the arc length and area of the sector formed by  $\theta$  can be expressed as the proportion of the angle to the full circle multiplied by the circumference and area of the circle, respectively.*

$$s = (\text{proportion of circle}) \cdot (\text{circumference}) = \left(\frac{\theta}{360^\circ}\right) \cdot (2\pi r)$$

and

$$A = (\text{proportion of circle}) \cdot (\text{area of circle}) = \left(\frac{\theta}{360^\circ}\right) \cdot (\pi r^2)$$

**Example 1.2.30** When sailing, Hōkūle‘a cannot make headway by sailing directly into the wind. It can only sail beyond  $67^\circ$  in either direction from the wind (Figure 1.2.31). If Hōkūle‘a sails for 50 miles, what is the area of the sector that cannot be sailed? Round your answer to the nearest square mile.



**Figure 1.2.31** Hōkūle‘a cannot sail within  $67^\circ$  into the direction of the wind.


**Solution.** The angle is  $\theta = 2 \cdot 67^\circ = 134^\circ$  and the radius is  $r = 50$  miles. So the area is given by

$$A = \frac{\theta}{360^\circ} \pi \cdot r^2 = \frac{134}{360} \pi \cdot 50^2 \approx 2,923 \text{ square miles}$$

□

### 1.2.5 Angular Velocity and Linear Speed

Consider an object moving along a circle as shown below. There are two ways to describe the circular motion of this object: *linear speed* which measures the distance traveled; and *angular speed* which measures the rate at which the central angle changes.



**Definition 1.2.32 Linear Speed.** Suppose an object moves along a circle with radius  $r$  and  $\theta$  (measured in radians) is the angle transversed in time  $t$ . Let  $s$  be the distance the object traveled in time  $t$ . Then the **linear speed**,  $v$ , of the object is given by

$$v = \frac{s}{t}$$

◇

**Definition 1.2.33 Angular Speed.** Suppose an object moves along a circle. Let  $\theta$  (measured in radians) be the angle transversed by the object in time  $t$ . The **angular speed**,  $\omega$ , of the object is given by

$$\omega = \frac{\theta}{t}$$

◇

Notice that we can rearrange the angular speed to get  $\theta = \omega t$ . Since  $s$  is an arc length, we have  $s = r\theta$ , and thus we can write the linear speed as


$$v = \frac{s}{t} = \frac{r\theta}{t} = \frac{r\omega t}{t} = r\omega$$

**Definition 1.2.34 Linear Speed.** Suppose an object moves along a circle with radius  $r$  and an angular speed  $\omega$  (measured in radians per unit time). Then the **linear speed**,  $v$ , of the object is given by


$$v = r\omega$$

◇

**Example 1.2.35** Une.



One method a wa'a uses to change direction is with the *hoe uli*, or the steering paddle. When a sharp turn is needed for maneuvers such as tacking, the steersperson will turn the handle of the hoe uli in a circular motion, as a lever to scoop the paddle in the water and change the heading of a vessel. This move called *une* (pronounced oo-NAY although it is often mispronounced as oo-NEE) literally translates to “lever.”



**Figure 1.2.36** A wa'a (canoe) can change directions by rotating the hoe uli (steering sweep) in a process known as *une*.

If the steersperson is performing an *une* at a rate of 25 rotations per minute and the radius of the circular movement is 2 feet, calculate:

- (a) The angular speed measured in radians per minute.

**Solution.** We are given the angular speed is  $\omega = 25$  revolutions per minute. To convert our angular speed to radians per minute, we use the fact that one revolution is  $2\pi$  radians to get

$$\omega = 25 \frac{\text{revolution}}{\text{minute}} = 25 \frac{\text{rev}}{\text{minute}} \cdot 2\pi \frac{\text{radians}}{\text{revolution}} = 50\pi \frac{\text{radians}}{\text{minute}}$$

Thus the hoe uli is moving at an angular speed of  $50\pi$  radians per second.

- (b) The linear speed of the hoe uli in miles per hour (round your answer to two decimal places).

**Solution.** Since the radius is  $r = 2$  ft and the angular speed is  $50\pi$  radians per minute, we can use [Definition 1.2.34](#) to calculate the linear speed

$$v = r\omega = 2\text{ft} \cdot 50\pi \frac{\text{rad}}{\text{min}} \cdot \frac{\text{mile}}{5280\text{ft}} \cdot \frac{60\text{min}}{\text{hr}} \approx 3.57 \frac{\text{miles}}{\text{hour}}$$

Thus the steersperson is moving the hoe uli at a linear speed of 3.57 mph.

□

## 1.2.6 Exercises


**Exercise Group.** Given an angle,  $\theta$ , identify the house and quadrant on the Hawaiian Star Compass.

- |                                 |                                  |                                |
|---------------------------------|----------------------------------|--------------------------------|
| 1. $\theta = 336^\circ$         | 2. $\theta = 240^\circ$          | 3. $\theta = 35^\circ$         |
| <b>Answer.</b> 'Āina<br>Malanai | <b>Answer.</b> Nālani<br>Kona    | <b>Answer.</b> Noio<br>Ko'olau |
| 4. $\theta = 221^\circ$         | 5. $\theta = 108^\circ$          | 6. $\theta = 190^\circ$        |
| <b>Answer.</b> Manu<br>Kona     | <b>Answer.</b> Nā<br>Leo Ho'olua | <b>Answer.</b> Lā<br>Kona      |
| 7. $\theta = 323^\circ$         | 8. $\theta = 158^\circ$          | 9. $\theta = 172^\circ$        |
| <b>Answer.</b> Noio<br>Malanai  | <b>Answer.</b> 'Āina<br>Ho'olua  | <b>Answer.</b> Lā<br>Ho'olua   |

**Exercise Group.** Convert the given angle  $\theta$  to a decimal in degrees rounded to two decimal places.

- |                                   |                                   |                                   |
|-----------------------------------|-----------------------------------|-----------------------------------|
| 10. $\theta = 20^\circ 50' 30''$  | 11. $\theta = 80^\circ 25' 16''$  | 12. $\theta = 7^\circ 22' 38''$   |
| <b>Answer.</b> $20.84^\circ$      | <b>Answer.</b> $80.42^\circ$      | <b>Answer.</b> $7.38^\circ$       |
| 13. $\theta = 330^\circ 14' 12''$ | 14. $\theta = 168^\circ 22' 26''$ | 15. $\theta = 49^\circ 27' 12''$  |
| <b>Answer.</b> $330.24^\circ$     | <b>Answer.</b> $168.37^\circ$     | <b>Answer.</b> $49.45^\circ$      |
| 16. $\theta = 327^\circ 38' 58''$ | 17. $\theta = 281^\circ 48' 50''$ | 18. $\theta = 134^\circ 53' 47''$ |
| <b>Answer.</b> $327.65^\circ$     | <b>Answer.</b> $281.81^\circ$     | <b>Answer.</b> $134.90^\circ$     |
19. Sirius, the brightest star in the night sky, has been known by various names in different cultures and languages around the world. In Tahiti, it is called Taurere and is considered a zenith star as it passes directly overhead. Taurere has a **declination** of  $-16^\circ 42' 58''$ , representing its angular distance south of the celestial equator. Express Taurere's declination as a decimal rounded to two decimal places.
- Answer.**  $16.72^\circ$

**Exercise Group.** Recall the Star Compass with the boundaries for each House. Write the angles for the boundaries between the following houses in the Ko'olau quadrant in terms of degrees, minutes, and seconds.



20. Between Hikina and Lā ( $5.625^\circ$ )  
**Answer.**  $5^\circ 37' 30''$
21. Between Lā and 'Āina ( $16.875^\circ$ )  
**Answer.**  $16^\circ 52' 30''$
22. Between 'Āina and Noio ( $28.175^\circ$ )  
**Answer.**  $28^\circ 7' 30''$
23. Between Noio and Manu ( $39.375^\circ$ )  
**Answer.**  $39^\circ 22' 30''$

**Exercise Group.** Convert the given angle  $\theta$  to degrees/minutes/seconds rounded to the nearest second and identify the house and quadrant on the Hawaiian Star Compass.

- |  |   |
|--|---|
| 24. $\theta = 258.39^\circ$<br><b>Answer.</b> $\theta = 258^\circ 23' 24''$<br>Haka Malanai  | 25. $\theta = 212.43^\circ$<br><b>Answer.</b> $\theta = 212^\circ 25' 33''$<br>Noio Kona    |
| 26. $\theta = 244.97^\circ$<br><b>Answer.</b> $\theta = 244^\circ 57' 59''$ Nā<br>Leo Kona   | 27. $\theta = 93.95^\circ$<br><b>Answer.</b> $\theta = 93^\circ 57' 6''$ Haka<br>Ho'olua    |
| 28. $\theta = 135.625^\circ$<br><b>Answer.</b> $\theta = 135^\circ 37' 30''$<br>Manu Ho'olua | 29. $\theta = 162.52^\circ$<br><b>Answer.</b> $\theta = 162^\circ 59' 56''$<br>Manu Ho'olua |
| 30. $\theta = 328.21^\circ$<br><b>Answer.</b> $\theta = 328^\circ 12' 9''$<br>Noio Malanai   | 31. $\theta = 48.12^\circ$<br><b>Answer.</b> $\theta = 48^\circ 7' 17''$<br>Manu Ko'olau    |
| 32. $\theta = 241.27^\circ$<br><b>Answer.</b> $\theta = 241^\circ 16' 15''$ Nā<br>Lani Kona  |   |

**Exercise Group.** Convert the given angle  $\theta$  to radians and identify the house and quadrant on the Hawaiian Star Compass. Keep your answers in terms of  $\pi$ .

33.  $\theta = 102^\circ$

Answer.  $\theta = \frac{17\pi}{30}$  Haka  
Ho'olua

35.  $\theta = 190^\circ$

Answer.  $\theta = \frac{19\pi}{18}$  Lā Kona

37.  $\theta = 32^\circ$

Answer.  $\theta = \frac{8\pi}{45}$  Noio  
Ko'olau

39.  $\theta = 82^\circ$

Answer.  $\theta = \frac{41\pi}{90}$  Haka  
Ko'olau

41.  $\theta = 332^\circ$

Answer.  $\theta = \frac{83\pi}{45}$  Āina  
Ko'olau

34.  $\theta = 96^\circ$

Answer.  $\theta = \frac{8\pi}{15}$  Ākau

36.  $\theta = 122^\circ$

Answer.  $\theta = \frac{61\pi}{90}$  Nālani  
Ho'olua

38.  $\theta = 236^\circ$

Answer.  $\theta = \frac{59\pi}{45}$  Nālani  
Kona


40.  $\theta = 228^\circ$

Answer.  $\theta = \frac{19\pi}{15}$  Manu  
Kona

**Exercise Group.** Draw the angle in standard position.


42.  $60^\circ$

Answer.




43.  $135^\circ$

Answer.




44.  $-30^\circ$

Answer.




45.  $495^\circ$

Answer.




46.  $-240^\circ$

Answer.




47.  $800^\circ$

Answer.




48.  $\frac{\pi}{4}$

Answer.




49.  $\frac{5\pi}{6}$

Answer.




50.  $-\frac{\pi}{3}$

Answer.




51.  $\frac{10\pi}{3}$

Answer.




52.  $\frac{11\pi}{6}$

Answer.



53.  $-\frac{2\pi}{3}$

Answer.



**Exercise Group.** Convert the given angle from degrees to radians. Round your answer to two decimal places.

- |                        |                        |                        |
|------------------------|------------------------|------------------------|
| <b>54.</b> $27^\circ$  | <b>55.</b> $63^\circ$  | <b>56.</b> $-39^\circ$ |
| <b>Answer.</b> 0.47    | <b>Answer.</b> 1.10    | <b>Answer.</b> $-0.68$ |
| <b>57.</b> $200^\circ$ | <b>58.</b> $415^\circ$ | <b>59.</b> $105^\circ$ |
| <b>Answer.</b> 3.49    | <b>Answer.</b> 7.24    | <b>Answer.</b> 1.83    |

**Exercise Group.** Convert the given angle from radians to degrees. Round your answer to two decimal places.

- |                              |                               |                                |
|------------------------------|-------------------------------|--------------------------------|
| <b>60.</b> $\frac{3\pi}{5}$  | <b>61.</b> $\frac{8\pi}{3}$   | <b>62.</b> $-\frac{4\pi}{7}$   |
| <b>Answer.</b> $108^\circ$   | <b>Answer.</b> $480^\circ$    | <b>Answer.</b> $-102.86^\circ$ |
| <b>63.</b> $\frac{15\pi}{4}$ | <b>64.</b> $-1.5$             | <b>65.</b> $3$                 |
| <b>Answer.</b> $675^\circ$   | <b>Answer.</b> $-85.94^\circ$ | <b>Answer.</b> $171.89^\circ$  |

**Exercise Group.** Determine whether the two given angles in standard position are coterminal.

- |                                   |                                    |                                   |                                    |
|-----------------------------------|------------------------------------|-----------------------------------|------------------------------------|
| <b>66.</b> $50^\circ, 410^\circ$  | <b>67.</b> $-60^\circ, 330^\circ$  | <b>68.</b> $30^\circ, 1110^\circ$ | <b>69.</b> $-210^\circ, 150^\circ$ |
| <b>Answer.</b> Yes                | <b>Answer.</b> Yes                 | <b>Answer.</b> No                 | <b>Answer.</b> Yes                 |
| <b>70.</b> $-807^\circ, 93^\circ$ | <b>71.</b> $-757^\circ, 683^\circ$ |                                   |                                    |
| <b>Answer.</b> No                 | <b>Answer.</b> Yes                 |                                   |                                    |

**Exercise Group.** Find an angle between  $0^\circ$  and  $360^\circ$  that is coterminal with the given angle


- |                           |                            |                            |
|---------------------------|----------------------------|----------------------------|
| <b>72.</b> $405^\circ$    | <b>73.</b> $600^\circ$     | <b>74.</b> $1035^\circ$    |
| <b>Answer.</b> $45^\circ$ | <b>Answer.</b> $240^\circ$ | <b>Answer.</b> $315^\circ$ |
| <b>75.</b> $-300^\circ$   | <b>76.</b> $381^\circ$     | <b>77.</b> $-754^\circ$    |
| <b>Answer.</b> $60^\circ$ | <b>Answer.</b> $21^\circ$  | <b>Answer.</b> $326^\circ$ |

**Exercise Group.** Given a circle with radius  $r$ , calculate (a) the length of the arc subtended by a central angle  $\theta$ ; and (b) the area of a sector with central angle  $\theta$ . Round your answer to four decimal places.

- |   |   |
|---|---|
| <b>78.</b> $r = 10$ in, $\theta = 45^\circ$             | <b>79.</b> $r = 5$ m, $\theta = 120^\circ$  |
| <b>Answer.</b>  | <b>Answer.</b>                              |
| (a) 7.8540 in;  | (a) 2.0944 m;                               |
| (b) 39.2699 in <sup>2</sup>                             | (b) 26.1799 m <sup>2</sup>                  |
| <b>80.</b> $r = 3$ mi, $\theta = \frac{\pi}{3}$ radians | <b>81.</b> $r = 8$ cm, $\theta = 2$ radians |
| <b>Answer.</b>  | <b>Answer.</b>                              |
| (a) 3.1416 mi;  | (a) 16 cm;                                  |
| (b) 4.7124 mi <sup>2</sup>                              | (b) 64 cm <sup>2</sup>                      |

**Exercise Group.** At the start of this section, you learned that the **latitude** of a place is the angle between a line drawn from the center of the earth to that point and the equatorial plane. If the radius of the Earth is 3,959 miles, calculate the arc length,  $s$ , along the surface of the earth for each value of  $\theta$ :





82.  $\theta = 1^\circ$  of latitude (in miles, rounded to 2 decimals)  
**Answer.** 69.10 miles
83.  $\theta = 1'$  of latitude (in miles, rounded to 2 decimals). A *nautical mile*, frequently used in navigation, is slightly longer than a mile on land. One nautical mile was historically defined to be the arc length corresponding to one minute of latitude. Check your answer with the value of one nautical mile.  
**Answer.** 1.15 miles
84.  $\theta = 1''$  of latitude (in feet, rounded to the nearest integer).  
**Answer.** 101 feet
85. The *oeoe*, or Hawaiian bullroarer, is made by drilling holes into a kamani seed or coconut shell, then threading a long string through the holes to secure it. When the *oeoe* is swung by the string, a whistling sound is produced, similar to the sound of the wind on the top of mountains. If a girl is swinging an *oeoe* at the end of 3 foot long rope at a rate of 180 revolutions per minute, calculate
- (a) The angular speed measured in radians per minute.  
**Answer.**  $360\pi$  radians/minute
- (b) The linear speed of the shell in miles per hour (round to two decimal places).  
**Answer.** 38.56 miles/hour
86. Earth completes one rotation around the Sun approximately every 365.25 days. We will assume the orbit is a circle, and that the Earth is 92.9 million miles from the Sun.
- (a) How far does the Earth travel in one day, expressed as millions of miles?  
**Hint.** First determine the angle or proportion of a rotation that Earth travels in one day, then calculate the arc length of Earth's orbit.  
**Answer.** 1.6 million miles

- (b) How far does the Earth travel in 30 days, expressed as millions of miles?

**Answer.** 47.9 million miles

- (c) How far does the Earth travel in one rotation around the sun, expressed as millions of miles?

**Answer.** 583.7 million miles

- (d) What is the linear speed of Earth as it orbits the Sun? Express your answer in miles per hour.

**Answer.** 66,588 miles per hour

87. As the the moon orbits the Earth, different parts of its surface become illuminated by the Sun which we call moon phases. The moon completes one rotation about Earth in approximately 27.3 days. If we assume its orbit is circular and the moon is 239,000 miles from Earth, calculate the linear speed of the moon, expressed as miles per hour.

**Answer.** 2,292 miles per hour

88. At  $17.7^\circ$  S latitude, the city of Nadi, Fiji is 6,071 km from the Earth's axis of rotation. In 24 hours, Nadi will have traveled one rotation around Earth or  $2\pi \cdot (6,071)$  km. The city of Port Vila, Vanuatu lies 967 km directly to the west of Nadi, Fiji. As the Earth rotates, how many minutes sooner will the people of Nadi see the Sun rise than the people in Port Vila, rounded to one decimal?

**Hint.** The proportion of distance between the two cities to the distance traveled in one rotation is the same as the proportion of the time it takes to see the sun between the two cities to time it takes to complete one rotation.

**Answer.** 36.5 minutes

89. In [Example 1.2.30](#), we learned that wa'a cannot sail directly into the wind. For each of the following wa'a and distance traveled, determine the area of the sector that cannot be sailed? Recall that 1 house =  $11.25^\circ$ . Round your answer to the nearest square mile.

- (a) Makali'i sails for 25 miles and cannot sail within 4 houses from the direction of the wind.

**Answer.** 491 square miles

- (b) Alingano Maisu sails for 15 miles and cannot sail within 3 houses from the direction of the wind.

**Answer.** 133 square miles

90. **Navigating by the Sun: Using Solar Declination and Rising Sun to Orient on a Canoe.** The position of the rising or setting sun changes throughout the year. **Solar declination** (denoted as  $\delta$ ) is the angle between the direction where the Sun rises (or sets) and due east (or due west) on the horizon. It represents how far north or south the Sun is from the celestial equator, projected onto the Earth's equatorial plane. Solar declinations to the north are positive, while those to the south are negative. At the Equinoxes (around March 20th and September 22nd), the solar declination is  $0^\circ$  ( $\delta = 0^\circ$ ), as the Sun is directly above the equator. During the December solstice, around December 22, the Sun rises from its most southern position, 23.5 degrees south of due east ( $\delta = -23.5^\circ$ ), and during

the June solstice, around June 22, the Sun rises from its most northern position, 23.5 degrees north of due east ( $\delta = 23.5^\circ$ ).

A navigator can use their knowledge of the rising sun to help orient themselves. For example, on May 22, the solar declination is  $\delta = 20^\circ 16'$ . If the navigator identifies where the Sun rises on the equinox, she can measure  $20^\circ 16'$  south to identify East and can then orient herself accordingly.

- (a) What is the azimuth of the sun?

**Answer.**  $69^\circ 44'$

- (b) What house is the sun rising in?

**Answer.** 'Āina Ko'olau

- (c) What house does the sun set in?

**Hint.** Celestial bodies rise and set in the same house but different quadrants.

**Answer.** 'Āina Ho'olua

- (d) If the canoe is sailing with the rising sun on the port side (left) and the navigator measures the angle between the direction of the canoe and the sun as being  $90^\circ$ , what is the heading of the canoe?

**Answer.**  $159^\circ 44'$

- (e) What house is the canoe sailing in?

**Answer.** Nā Leo Malanai

91. Swells are one of the most consistent navigational tools used to keep on a course because they can remain constant over time. On 27 May 2023, while sailing on the vaka Paikea from Rarotonga to Apia, you finished your shift and are ready to take a nap. Before you lay down, you take note that the canoe has a heading of  $310^\circ$  and the swells are coming from the southwest (Manu Kona) and hitting the canoe at  $5^\circ$  above directly left of the canoe. When you wake, you noticed the swells are now hitting the canoe from  $25^\circ$  to the left of your heading. You are aware that the swells couldn't have changed this fast and conclude that while you were asleep, the canoe changed its heading. Assuming the swell was constant, determine your new heading.

**Hint.**

- (a) Start by drawing a diagram that represents the heading of the canoe before the nap. Mark the initial heading as  $^\circ >$ .
- (b) Next, draw the direction of the swells with respect to the canoe on the same diagram. The swells are coming from the southeast (Manu Kona) and hitting the canoe at  $^\circ >$  above straight from the left of the canoe.
- (c) Now, draw another diagram of the swells and the canoe, but this time, represent the swells hitting the canoe from  $^\circ >$  to the left of straight in front (Nā Leo Ho'olua).
- (d) Observe that the swells couldn't have changed direction so fast while you were asleep. Thus, the change in the direction of the swells must be due to the canoe changing its heading.

- (e) Use the relationship between the angle of the swell and the angle between the swell and the canoe to determine the angle by which the canoe's heading changed.
- (f) Finally, update the initial heading of 310 degrees with the angle of change to find the new heading of the canoe after the nap.

**Answer.**  $250^\circ$

- 92.** After spending 6 weeks in Samoa, vaka Paikea is making his way back from Apia to Rarotonga. On July 14, 2023, the canoe is sailing with a heading of  $50^\circ$ , and the wind is coming to the canoe from  $60^\circ$  to the right of the your heading. What is the heading and house from which the wind is coming?

**Answer.**  $110^\circ$ ; 'Āina Malanai

## 1.3 Unit Circle

In this section, we will introduce the trigonometric functions using the Unit Circle.


### 1.3.1 Unit Circle

**Definition 1.3.1 Unit Circle.** The **unit circle** is a circle whose radius is 1 and whose center is at the origin of a rectangular plane (or  $xy$ -plane). The equation for the unit circle is


$$x^2 + y^2 = 1$$

◇

Let  $t$  be a real number. Recall from [Definition 1.2.18](#) that a radian measure of a central angle,  $t$ , is defined as the ratio of the arc length  $s$  to the radius  $r$ . In other words,  $t = \frac{s}{r}$ . In the unit circle, the radius is  $r = 1$ , and the angle in radians is equal to the arc length,  $t = s$ . We will let  $t$  be in radians. The circumference of the unit circle is  $2\pi r = 2\pi \cdot 1 = 2\pi$ .



If  $t \geq 0$ , we can imagine wrapping a line around the unit circle, marking off a distance of  $t$  in a counterclockwise direction, and labeling that point  $P(x, y)$ , which becomes the terminal point. If  $t < 0$  then we would wrap in a clockwise direction.



If  $t > 2\pi$  or  $t < -2\pi$ , then the length is longer than the circumference of the unit circle and you will need to travel around the unit circle more than once before arrive at the point  $P(x, y)$ . Therefore, we can conclude that regardless of the value of  $t$ , we have a unique point  $P(x, y)$  that lies on the unit circle. We call  $P(x, y)$  the *point on the unit circle that corresponds to  $t$* .

### 1.3.2 Trigonometric Functions


The  $x$ - and  $y$ -coordinates for  $P(x, y)$  can then be used to define the six trigonometric functions of a real number  $t$ :

*sine cosine tangent cosecant secant cotangent*

which are abbreviated as *sin*, *cos*, *tan*, *csc*, *sec*, and *cot*, respectively.

**Definition 1.3.2 Definition of Trigonometric Functions.** Let  $t$  be any real number and let  $P(x, y)$  be the terminal point on the unit circle associated with  $t$ . Then

$$\begin{array}{lll} \sin t = y & \cos t = x & \tan t = \frac{y}{x}, (x \neq 0) \\ \csc t = \frac{1}{y}, (y \neq 0) & \sec t = \frac{1}{x}, (x \neq 0) & \cot t = \frac{x}{y}, (y \neq 0) \end{array}$$



Notice that  $\tan t$  and  $\sec t$  are undefined when  $x = 0$  and  $\csc t$  and  $\cot t$  are undefined when  $y = 0$ .  $\diamond$

**Example 1.3.3** Let  $t$  be the angle that corresponds to the point  $P(\frac{\sqrt{3}}{2}, -\frac{1}{2})$ . Find the exact values of the six trigonometric functions corresponding to  $t$ :  $\sin t$ ,  $\cos t$ ,  $\tan t$ ,  $\csc t$ ,  $\sec t$ ,  $\cot t$ .

**Solution.** The point  $P(\frac{\sqrt{3}}{2}, -\frac{1}{2})$  gives us  $x = \frac{\sqrt{3}}{2}$  and  $y = -\frac{1}{2}$ . Then we have

$$\begin{aligned} \sin \theta &= y = -\frac{1}{2}, & \csc \theta &= \frac{1}{y} = \frac{1}{-\frac{1}{2}} = -2, \\ \cos \theta &= x = \frac{\sqrt{3}}{2}, & \sec \theta &= \frac{1}{x} = \frac{1}{\frac{\sqrt{3}}{2}} = \frac{2}{\sqrt{3}} = \frac{2\sqrt{3}}{3}, \\ \tan \theta &= \frac{y}{x} = \frac{-\frac{1}{2}}{\frac{\sqrt{3}}{2}} = -\frac{1}{\sqrt{3}} = -\frac{\sqrt{3}}{3}, & \cot \theta &= \frac{x}{y} = \frac{\frac{\sqrt{3}}{2}}{-\frac{1}{2}} = -\sqrt{3}. \end{aligned}$$

□

### 1.3.3 Trigonometric Functions of an Angle

**Definition 1.3.4 Trigonometric Functions of an Angle.** If  $\theta$  is an angle with radian measure  $t$ , then the **six trigonometric functions** become


$$\begin{aligned} \sin \theta &= y & \cos \theta &= x & \tan \theta &= \frac{y}{x}, \quad (x \neq 0) \\ \csc \theta &= \frac{1}{y}, \quad (y \neq 0) & \sec \theta &= \frac{1}{x}, \quad (x \neq 0) & \cot \theta &= \frac{x}{y}, \quad (y \neq 0) \end{aligned}$$

◇

**Example 1.3.5** Find the exact values of the six trigonometric functions for

(a)  $\theta = 0$

**Solution.** When  $\theta = 0$  radians ( $0^\circ$ ), the point on the circle is  $P(1, 0)$ .




Then  $x = 1$  and  $y = 0$  gives us

$$\begin{array}{ll} \sin 0 = \sin 0^\circ = 0, & \csc 0 = \csc 0^\circ = \text{undefined}, \\ \cos 0 = \cos 0^\circ = 1, & \sec 0 = \sec 0^\circ = 1, \\ \tan 0 = \tan 0^\circ = 0, & \cot 0 = \cot 0^\circ = \text{undefined}. \end{array}$$

(b)  $\theta = \frac{3\pi}{2}$

**Solution.** When  $\theta = \frac{3\pi}{2}$  radians ( $270^\circ$ ), the point on the circle is  $P(0, -1)$ .



Then  $x = 0$  and  $y = -1$  gives us

$$\begin{array}{ll} \sin \frac{3\pi}{2} = \sin 270^\circ = -1, & \csc \frac{3\pi}{2} = \csc 270^\circ = -1, \\ \cos \frac{3\pi}{2} = \cos 270^\circ = 0, & \sec \frac{3\pi}{2} = \sec 270^\circ = \text{undefined}, \\ \tan \frac{3\pi}{2} = \tan 270^\circ = \text{undefined}, & \cot \frac{3\pi}{2} = \cot 270^\circ = 0 \end{array}$$

(c)  $\theta = 5\pi$

**Solution.** Since  $\theta = 5\pi > 2\pi$ , our angle is greater than one full rotation of a circle. We first subtract  $\theta$  by one rotation,  $2\pi$ , to get


$$5\pi - 2\pi = 3\pi$$



Once again, since we have completed more than one full rotation, we can repeat the previous step:

$$3\pi - 2\pi = \pi$$

The values of the six trigonometric functions when  $\theta = 5\pi$  are equal to those when  $\theta = \pi$ . Notice that  $5\pi$  and  $\pi$  are *coterminal angles*, both ending at the point  $P(-1, 0)$ .




Since  $x = -1$  and  $y = 0$  we have

$$\begin{array}{lll} \sin 5\pi = 0, & \cos 5\pi = -1, & \tan 5\pi = 0, \\ \csc 5\pi = \text{undefined}, & \sec 5\pi = -1, & \cot 5\pi = \text{undefined}. \end{array}$$

□

**Example 1.3.6 Finding the Exact Values of the Trigonometric Functions for  $\theta = 45^\circ$ .** Find the exact values of the six trigonometric functions for  $\theta = 45^\circ$ .

**Solution.** We begin by drawing a right triangle with a base angle of  $45^\circ$  in the unit circle.



Since the first quadrant has  $90^\circ$ , at  $\theta = 45^\circ$ , the point  $P$  lies on the line that bisects the first quadrant. This means the point  $P$  is on the line  $y = x$ . Since  $P(x, y)$  also lies on the unit circle, whose equation is  $x^2 + y^2 = 1$ , we get

$$x^2 + y^2 = 1$$

$$\begin{aligned}
 x^2 + x^2 &= 1 && \text{(since } y = x\text{)} \\
 2x^2 &= 1 \\
 x^2 &= \frac{1}{2} \\
 x &= \frac{1}{\sqrt{2}} \\
 y &= \frac{1}{\sqrt{2}} && \text{(since } y = x\text{)}
 \end{aligned}$$


Then

$$\begin{aligned}
 \sin 45^\circ &= \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2} && \csc 45^\circ = \frac{1}{\frac{\sqrt{2}}{2}} = \sqrt{2} \\
 \cos 45^\circ &= \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2} && \sec 45^\circ = \frac{1}{\frac{\sqrt{2}}{2}} = \sqrt{2} \\
 \tan 45^\circ &= \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = 1 && \cot 45^\circ = \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = 1
 \end{aligned}$$


□

**Example 1.3.7 Finding the Exact Values of the Trigonometric Functions for  $\theta = 30^\circ$ .** Find the exact values of the six trigonometric functions for  $\theta = 30^\circ$ .

**Solution.** First, we will draw a triangle in a circle with an angle of  $30^\circ$  and a second triangle with an angle of  $-30^\circ$ .




This gives us two 30-60-90 triangles. Notice this now gives us one larger triangle whose angles are all  $60^\circ$ . Thus we have an equilateral triangle, with each side of length 1.



We see that  $1 = 2y$  so  $y = \frac{1}{2}$ . Then by the Pythagorean Theorem,

$$\begin{aligned} x^2 + y^2 &= 1^2 \\ x^2 + \left(\frac{1}{2}\right)^2 &= 1 \\ x^2 + \frac{1}{4} &= 1 \\ x^2 &= \frac{3}{4} \\ x &= \frac{\sqrt{3}}{2} \end{aligned}$$

Giving us the following triangle




Then

$$\begin{aligned} \sin 30^\circ &= \frac{1}{2}, & \csc 30^\circ &= \frac{1}{\frac{1}{2}} = 2, \\ \cos 30^\circ &= \frac{\sqrt{3}}{2}, & \sec 30^\circ &= \frac{1}{\frac{\sqrt{3}}{2}} = \frac{2}{\sqrt{3}} = \frac{2\sqrt{3}}{3}, \end{aligned}$$

$$\tan 30^\circ = \frac{\frac{1}{2}}{\frac{\sqrt{3}}{2}} = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}, \quad \cot 30^\circ = \frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}} = \sqrt{3}.$$

□

**Remark 1.3.8 Finding the Exact Values of the Trigonometric Functions for  $\theta = 90^\circ$ .** Similarly, we can get the following for  $\theta = 60^\circ$ .



We now summarize what we know about the six trigonometric functions for special angles. Note the trigonometric functions for  $\theta = \frac{\pi}{2}$  and  $\theta = \frac{\pi}{3}$  are left as exercises.

**Table 1.3.9 Trigonometric functions for special angles**

$\theta$ (deg)	$\theta$ (rad)	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\csc \theta$	$\sec \theta$	$\cot \theta$
$0^\circ$	0	0	1	0	undef	1	undef
$30^\circ$	$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$	$\sqrt{3}$
$45^\circ$	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	$\sqrt{2}$	$\sqrt{2}$	1
$60^\circ$	$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2	$\frac{\sqrt{3}}{3}$
$90^\circ$	$\frac{\pi}{2}$	1	0	undef	1	undef	0


### 1.3.4 Symmetry on the Unit Circle

If the point  $P(x, y)$  lies on the unit circle, the following symmetric points also lie on the unit circle:

1.  $Q(-x, y)$ : Symmetry about the  $y$ -axis.
2.  $R(-x, -y)$ : Symmetry about the origin.


3.  $S(x, -y)$ : Symmetry about the  $x$ -axis.

This symmetry within the unit circle resembles the pattern observed in the Star Compass. When a star emerges in the eastern sky, it will eventually descend and set in the corresponding house of the western sky. For instance, if a star rises above the horizon in the Nālani house of the Ko‘olau quadrant (northeast), it will journey across the sky and set in the equivalent house within the Ho‘olua quadrant (northwest). This similarity aligns with the symmetry between points  $P(x, y)$  and  $Q(-x, y)$ . Additionally, if an ocean swell or wind originates from the Nālani house in the Malanai quadrant (southeast), it will pass the wa‘a and exit in the opposite direction toward the Ho‘olua quadrant (northwest), still within the Nālani house. This mirrors the symmetry between points  $S(x, -y)$  and  $Q(-x, y)$ .




A fourth form of symmetry involves reflecting points across the diagonal line  $y = x$ , where the  $x$ - and  $y$ -values are equal.


1.  $T(y, x)$ : Symmetry about the line  $y = x$ . This is accomplished by interchanging the  $x$ - and  $y$ -values.



Notice on the Unit Circle that the radius extending from the center at an angle of  $30^\circ$  to the point  $T(x, y) = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$  is symmetric about the line  $y = x$ , in relation to the radius extending from the center at an angle of  $60^\circ$  to the point  $P(x, y)$ .



Using symmetry about the  $x$ -axis, symmetry about the  $y$ -axis, and symmetry about the origin, we can complete the unit circle, as long as we remember that the  $x$ -values in Quadrants II and III are negative while the  $y$ -values in Quadrants III and IV are negative.



Finally, we tie everything together and look at the entire Unit Circle. At first glance it may seem intimidating, however, similar to the Star Compass, there is a lot of symmetry ( $x$ -axis,  $y$ -axis, origin, about the line  $y = x$ ) and it can help by focusing on one quadrant, and use symmetry to fill out the rest of the circle.




Figure 1.3.10 The Unit Circle for common angles in radians and degrees.

### 1.3.5 Trigonometric Functions on a Circle with Radius $r$

Until now, computing the exact values of trigonometric functions of an angle  $\theta$  required us to locate the corresponding point  $P(x, y)$  on the unit circle. However, we can use any circle with center at the origin, that is, any circle of the form  $x^2 + y^2 = r^2$ , where  $r > 0$  is the radius. Note that if  $r = 1$ , then it is the unit circle.

**Theorem 1.3.11** For an angle  $\theta$  in standard position, let  $P(x, y)$  be the point on the terminal side of  $\theta$  that is also on the circle  $x^2 + y^2 = r^2$ . Then

$$\begin{aligned} \sin \theta &= \frac{y}{r} & \cos \theta &= \frac{x}{r} & \tan \theta &= \frac{y}{x}, \quad (x \neq 0) \\ \csc \theta &= \frac{r}{y}, \quad (y \neq 0) & \sec \theta &= \frac{r}{x}, \quad (x \neq 0) & \cot \theta &= \frac{x}{y}, \quad (y \neq 0) \end{aligned}$$

### 1.3.6 Exercises

**Exercise Group.** Verify algebraically that the point  $P$  is on the unit circle ( $x^2 + y^2 = 1$ )

1.  $P\left(\frac{3}{5}, -\frac{4}{5}\right)$

**Answer.**  $\left(\frac{3}{5}\right)^2 + \left(-\frac{4}{5}\right)^2 = 1$

2.  $P\left(-\frac{\sqrt{39}}{8}, -\frac{5}{8}\right)$

**Answer.**  $\left(-\frac{\sqrt{39}}{8}\right)^2 + \left(-\frac{5}{8}\right)^2 = 1$

3.  $P\left(-\frac{\sqrt{55}}{8}, \frac{3}{8}\right)$

**Answer.**  $\left(-\frac{\sqrt{55}}{8}\right)^2 + \left(\frac{3}{8}\right)^2 = 1$

$$\begin{array}{lll}
 4. & P\left(-\frac{2}{3}, \frac{\sqrt{5}}{3}\right) & 5. & P\left(\frac{3}{4}, \frac{\sqrt{7}}{4}\right) & 6. & P\left(\frac{\sqrt{21}}{5}, -\frac{2}{5}\right) \\
 \text{Answer.} & \left(-\frac{2}{3}\right)^2 + \left(\frac{\sqrt{5}}{3}\right)^2 = 1 & \text{Answer.} & \left(\frac{3}{4}\right)^2 + \left(\frac{\sqrt{7}}{4}\right)^2 = 1 & \text{Answer.} & \left(\frac{\sqrt{21}}{5}\right)^2 + \left(-\frac{2}{5}\right)^2 = 1
 \end{array}$$

**Exercise Group.** Let the point  $P$  be on the unit circle. Given the quadrant that  $P$  lies in, determine the missing coordinate,  $a$

$$\begin{array}{ll}
 7. & \text{III; } P\left(-\frac{2}{3}, a\right) & 8. & \text{IV; } P\left(\frac{5}{8}, a\right) \\
 \text{Answer.} & -\frac{\sqrt{5}}{3} & \text{Answer.} & -\frac{\sqrt{39}}{8} \\
 9. & \text{III; } P\left(a, -\frac{2}{5}\right) & 10. & \text{II; } P\left(a, \frac{4}{9}\right) \\
 \text{Answer.} & -\frac{\sqrt{21}}{5} & \text{Answer.} & -\frac{\sqrt{65}}{9}
 \end{array}$$

**Exercise Group.** Given an angle  $\theta$  that corresponds to the point  $P$  on the unit circle, determine the coordinates of the point  $P(x, y)$ .

$$\begin{array}{llll}
 11. & \theta = \frac{\pi}{2} & 12. & \theta = \pi & 13. & \theta = \frac{5\pi}{3} & 14. & \theta = \frac{4\pi}{3} \\
 \text{Answer.} & (0, 1) & \text{Answer.} & (-1, 0) & \text{Answer.} & \left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right) & \text{Answer.} & \left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right) \\
 15. & \theta = -\frac{\pi}{4} & 16. & \theta = \frac{5\pi}{6} & 17. & \theta = 315^\circ & 18. & \theta = 720^\circ \\
 \text{Answer.} & \left(\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}\right) & \text{Answer.} & \left(-\frac{\sqrt{3}}{2}, \frac{1}{2}\right) & \text{Answer.} & \left(\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}\right) & \text{Answer.} & (1, 0) \\
 19. & \theta = 60^\circ & 20. & \theta = -180^\circ & 21. & \theta = 210^\circ & 22. & \theta = 120^\circ \\
 \text{Answer.} & \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right) & \text{Answer.} & (-1, 0) & \text{Answer.} & \left(-\frac{\sqrt{3}}{2}, -\frac{1}{2}\right) & \text{Answer.} & \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right)
 \end{array}$$

**Exercise Group.** For each angle  $\theta$  in Exercises 1.3.6.11–22, find the exact values of the six trigonometric functions. If any are not defined, say “undefined.”

$$\begin{array}{lll}
 23. & \theta = \frac{\pi}{2} & 24. & \theta = \pi & 25. & \theta = \frac{5\pi}{3} \\
 \text{Answer.} & \sin \frac{\pi}{2} = 1; \cos \frac{\pi}{2} = 0; \tan \frac{\pi}{2} \text{ is undefined; } \csc \frac{\pi}{2} = 1; \sec \frac{\pi}{2} \text{ is undefined; } \cot \frac{\pi}{2} = 0 & \text{Answer.} & \sin \pi = 0; \cos \pi = -1; \tan \pi = 0; \csc \pi \text{ is undefined; } \sec \pi = -1; \cot \pi \text{ is undefined} & \text{Answer.} & \sin \frac{5\pi}{3} = -\frac{\sqrt{3}}{2}; \cos \frac{5\pi}{3} = \frac{1}{2}; \tan \frac{5\pi}{3} = -\sqrt{3}; \csc \frac{5\pi}{3} = -\frac{2\sqrt{3}}{3}; \sec \frac{5\pi}{3} = 2; \cot \frac{5\pi}{3} = -\frac{\sqrt{3}}{3} \\
 26. & \theta = \frac{4\pi}{3} & 27. & \theta = -\frac{\pi}{4} & 28. & \theta = \frac{5\pi}{6} \\
 \text{Answer.} & \sin \frac{4\pi}{3} = -\frac{\sqrt{3}}{2}; \cos \frac{4\pi}{3} = -\frac{1}{2}; \tan \frac{4\pi}{3} = \sqrt{3}; \csc \frac{4\pi}{3} = -\frac{2\sqrt{3}}{3}; \sec \frac{4\pi}{3} = -2; \cot \frac{4\pi}{3} = \frac{\sqrt{3}}{3} & \text{Answer.} & \sin\left(-\frac{\pi}{4}\right) = -\frac{\sqrt{2}}{2}; \cos\left(-\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2}; \tan\left(-\frac{\pi}{4}\right) = -1; \csc\left(-\frac{\pi}{4}\right) = -\sqrt{2}; \sec\left(-\frac{\pi}{4}\right) = \sqrt{2}; \cot\left(-\frac{\pi}{4}\right) = -1 & \text{Answer.} & \sin \frac{5\pi}{6} = \frac{1}{2}; \cos \frac{5\pi}{6} = -\frac{\sqrt{3}}{2}; \tan \frac{5\pi}{6} = -\frac{\sqrt{3}}{3}; \csc \frac{5\pi}{6} = 2; \sec \frac{5\pi}{6} = -\frac{2\sqrt{3}}{3}; \cot \frac{5\pi}{6} = -\sqrt{3}
 \end{array}$$



29.  $\theta = 315^\circ$

**Answer.**  $\sin 315^\circ = -\frac{\sqrt{2}}{2}$ ;  
 $\cos 315^\circ = \frac{\sqrt{2}}{2}$ ;  
 $\tan 315^\circ = -1$ ;  
 $\csc 315^\circ = -\sqrt{2}$ ;  
 $\sec 315^\circ = \sqrt{2}$ ;  
 $\cot 315^\circ = -1$

30.  $\theta = 720^\circ$

**Answer.**  $\sin 720^\circ = 0$ ;  $\cos 720^\circ = 1$ ;  
 $\tan 720^\circ = 0$ ;  
 $\csc 720^\circ$  is undefined;  
 $\sec 720^\circ = 1$ ;  
 $\cot 720^\circ$  is undefined

31.  $\theta = 60^\circ$

**Answer.**  $\sin 60^\circ = \frac{\sqrt{3}}{2}$ ;  $\cos 60^\circ = \frac{1}{2}$ ;  
 $\tan 60^\circ = \sqrt{3}$ ;  
 $\csc 60^\circ = \frac{2\sqrt{3}}{3}$ ;  
 $\sec 60^\circ = 2$ ;  
 $\cot 60^\circ = \frac{\sqrt{3}}{3}$

32.  $\theta = -180^\circ$

**Answer.**  $\sin(-180^\circ) = 0$ ;  $\cos(-180^\circ) = -1$ ;  
 $\tan(-180^\circ) = 0$ ;  
 $\csc(-180^\circ)$  is undefined;  
 $\sec(-180^\circ) = -1$ ;  $\cot(-180^\circ)$  is undefined

33.  $\theta = 210^\circ$

**Answer.**  $\sin 210^\circ = -\frac{1}{2}$ ;  
 $\cos 210^\circ = -\frac{\sqrt{3}}{2}$ ;  
 $\tan 210^\circ = \frac{\sqrt{3}}{3}$ ;  
 $\csc 210^\circ = -2$ ;  
 $\sec 210^\circ = -\frac{2\sqrt{3}}{3}$ ;  
 $\cot 210^\circ = \sqrt{3}$

34.  $\theta = 120^\circ$

**Answer.**  $\sin 120^\circ = \frac{\sqrt{3}}{2}$ ;  
 $\cos 120^\circ = -\frac{1}{2}$ ;  
 $\tan 120^\circ = -\sqrt{3}$ ;  
 $\csc 120^\circ = \frac{2\sqrt{3}}{3}$ ;  
 $\sec 120^\circ = -2$ ;  
 $\cot 120^\circ = -\frac{\sqrt{3}}{3}$

**Exercise Group.** Let  $\theta$  be the angle that corresponds to the point  $P$ . Exercises 1.3.6.1–6 verified  $P$  is on the unit circle. Find the exact values of the six trigonometric functions of  $\theta$ .

35.  $P\left(\frac{3}{5}, -\frac{4}{5}\right)$

**Answer.**  $\sin \theta = -\frac{4}{5}$ ;  $\cos \theta = \frac{3}{5}$ ;  
 $\tan \theta = -\frac{4}{3}$ ;  
 $\csc \theta = -\frac{5}{4}$ ;  
 $\sec \theta = \frac{5}{3}$ ;  
 $\cot \theta = -\frac{3}{4}$

36.  $P\left(-\frac{\sqrt{39}}{8}, -\frac{5}{8}\right)$

**Answer.**  $\sin \theta = -\frac{5}{8}$ ;  
 $\cos \theta = -\frac{\sqrt{39}}{8}$ ;  
 $\tan \theta = \frac{5\sqrt{39}}{39}$ ;  
 $\csc \theta = -\frac{8}{5}$ ;  
 $\sec \theta = -\frac{8\sqrt{39}}{39}$ ;  
 $\cot \theta = \frac{\sqrt{39}}{5}$

37.  $P\left(-\frac{\sqrt{55}}{8}, \frac{3}{8}\right)$

**Answer.**  $\sin \theta = \frac{3}{8}$ ;  $\cos \theta = -\frac{\sqrt{55}}{8}$ ;  
 $\tan \theta = -\frac{3\sqrt{55}}{55}$ ;  
 $\csc \theta = \frac{8}{3}$ ;  
 $\sec \theta = -\frac{8\sqrt{55}}{55}$ ;  
 $\cot \theta = -\frac{\sqrt{55}}{3}$

38.  $P\left(-\frac{2}{3}, \frac{\sqrt{5}}{3}\right)$

**Answer.**  $\sin \theta = \frac{\sqrt{5}}{3}$ ;  $\cos \theta = -\frac{2}{3}$ ;  
 $\tan \theta = -\frac{\sqrt{5}}{2}$ ;  
 $\csc \theta = \frac{3\sqrt{5}}{5}$ ;  
 $\sec \theta = -\frac{3}{2}$ ;  
 $\cot \theta = -\frac{2\sqrt{5}}{5}$

39.  $P\left(\frac{3}{4}, \frac{\sqrt{7}}{4}\right)$

**Answer.**  $\sin \theta = \frac{\sqrt{7}}{4}$ ;  $\cos \theta = \frac{3}{4}$ ;  
 $\tan \theta = \frac{\sqrt{7}}{3}$ ;  
 $\csc \theta = \frac{4\sqrt{7}}{7}$ ;  
 $\sec \theta = \frac{4}{3}$ ;  
 $\cot \theta = \frac{3\sqrt{7}}{7}$

40.  $P\left(\frac{\sqrt{21}}{5}, -\frac{2}{5}\right)$

**Answer.**  $\sin \theta = -\frac{2}{5}$ ;  $\cos \theta = \frac{\sqrt{21}}{5}$ ;  
 $\tan \theta = -\frac{2\sqrt{21}}{21}$ ;  
 $\csc \theta = -\frac{5}{2}$ ;  
 $\sec \theta = \frac{5\sqrt{21}}{21}$ ;  
 $\cot \theta = -\frac{\sqrt{21}}{2}$

**Exercise Group.** Find the exact value of each expression.

41.  $\sin 30^\circ + \sin 150^\circ$

**Answer.** 1

42.  $\cos 30^\circ + \cos 150^\circ$

**Answer.** 0

43.  $\sin 60^\circ + \sin 120^\circ + \sin 240^\circ + \sin 300^\circ$

**Answer.** 0

44.  $\cos 60^\circ + \cos 120^\circ + \cos 240^\circ + \cos 300^\circ$

**Answer.** 0

45.  $\tan 45^\circ + \tan 135^\circ$

**Answer.** 0

46.  $\tan 135^\circ + \tan 225^\circ$

**Answer.** 0


47.  $\tan 225^\circ + \tan 315^\circ$

**Answer.** 0

48.  $\tan 45^\circ + \tan 225^\circ$

**Answer.** 2


## 1.4 Right Triangle Trigonometry



During a voyage, a navigator utilizes a *reference course*—a line connecting the starting point and destination—to monitor their position. When the wa'a (canoe) encounters winds that veer it off course, the navigator mentally plots their position relative to the reference course. To ensure the destination isn't missed, navigators must monitor their deviation from the intended course, involving measurement of the angle of deviation from the reference course (in units of houses) and determining the distance traveled. This section explores the calculation of trigonometric functions using right triangles, enabling us to assess how much the wa'a has strayed from its intended reference course.

### 1.4.1 Trigonometric Ratios

**Definition 1.4.1 Trigonometric Ratios.** Consider a right triangle with  $\theta$  as one of its acute angles. The trigonometric ratios are defined as follows:



$$\begin{array}{lll} \sin \theta = \frac{\text{opposite}}{\text{hypotenuse}} & \cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}} & \tan \theta = \frac{\text{opposite}}{\text{adjacent}} \\ \csc \theta = \frac{\text{hypotenuse}}{\text{opposite}} & \sec \theta = \frac{\text{hypotenuse}}{\text{adjacent}} & \cot \theta = \frac{\text{adjacent}}{\text{opposite}} \end{array}$$

A common mnemonic for remembering these relationships is SOHCAHTOA, formed from the first letters of “Sine is Opposite over Hypotenuse, Cosine is Adjacent over Hypotenuse, Tangent is Opposite over Adjacent.”  $\diamond$

Based on the definition of the six trigonometric functions, we have the following trigonometric identities.

**Definition 1.4.2 Reciprocal Identities.**

$$\begin{array}{lll} \sin \theta = \frac{1}{\csc \theta} & \cos \theta = \frac{1}{\sec \theta} & \tan \theta = \frac{1}{\cot \theta} \\ \csc \theta = \frac{1}{\sin \theta} & \sec \theta = \frac{1}{\cos \theta} & \cot \theta = \frac{1}{\tan \theta} \end{array}$$


$\diamond$

**Definition 1.4.3** Quotient Identities.

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta}$$

◇

**Example 1.4.4** Find the exact values of the six trigonometric ratios of the angle  $\theta$  in the given triangle.




**Solution.** By the definition of the trigonometric ratios, we have

$$\begin{array}{lll} \sin \theta = \frac{3}{5} & \cos \theta = \frac{4}{5} & \tan \theta = \frac{3}{4} \\ \csc \theta = \frac{5}{3} & \sec \theta = \frac{5}{4} & \cot \theta = \frac{4}{3} \end{array}$$

□

**Example 1.4.5** Find the exact values of the six trigonometric ratios of the angle  $\theta$  in the given triangle.



**Solution.** Notice that  $\theta$  is in a different position. Here, the adjacent side is 3 and the opposite side is 5. If we let  $h$  denote the hypotenuse, then we can use the Pythagorean Theorem to get

$$h = \sqrt{5^2 + 2^2} = \sqrt{29}$$


Then by the definition of the trigonometric ratios, we have

$$\begin{array}{lll} \sin \theta = \frac{5}{\sqrt{29}} = \frac{5\sqrt{29}}{29} & \cos \theta = \frac{2}{\sqrt{29}} = \frac{2\sqrt{29}}{29} & \tan \theta = \frac{5}{2} \\ \csc \theta = \frac{\sqrt{29}}{5} & \sec \theta = \frac{\sqrt{29}}{2} & \cot \theta = \frac{2}{5} \end{array}$$

□

### 1.4.2 Special Triangles

The angles  $30^\circ$ ,  $45^\circ$ ,  $60^\circ$  ( $\frac{\pi}{6}$ ,  $\frac{\pi}{4}$ ,  $\frac{\pi}{3}$ ) give special values for trigonometric functions. The following figures are used to calculate trigonometric values.



The trigonometric values for the special angles  $0^\circ$ ,  $30^\circ$ ,  $45^\circ$ ,  $60^\circ$ ,  $90^\circ$  ( $0$ ,  $\frac{\pi}{6}$ ,  $\frac{\pi}{4}$ ,  $\frac{\pi}{3}$ ,  $\frac{\pi}{2}$ ) are given in [Table 1.4.6](#).


**Table 1.4.6 Values of the trigonometric functions in Quadrant I**

$\theta$ (degrees)	$\theta$ (radians)	$\sin \theta$	$\cos \theta$	$\tan \theta$
$0^\circ$	$0$	$0$	$1$	$0$
$30^\circ$	$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$
$45^\circ$	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	$1$
$60^\circ$	$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$90^\circ$	$\frac{\pi}{2}$	$1$	$0$	undefined

### 1.4.3 Cofunctions

The symmetry between  $\sin \theta$  and  $\cos \theta$  becomes evident when reversing the order of sine and cosine values from  $90^\circ$  to  $0^\circ$ . This symmetry yields  $\sin 0^\circ = \cos 90^\circ$ ,  $\sin 30^\circ = \cos 60^\circ$ ,  $\sin 45^\circ = \cos 45^\circ$ ,  $\sin 60^\circ = \cos 30^\circ$ , and  $\sin 90^\circ = \cos 0^\circ$ .

This pattern between sine and cosine is no coincidence; it emerges because the three angles in a triangle add up to  $180^\circ$  or  $\pi$  radians. When considering a right triangle, the remaining two angles combine to form  $90^\circ$  or  $\frac{\pi}{2}$  radians, making them *complementary angles*.



Consider the right triangle in the figure above, where angles  $\alpha$  and  $\beta$  are complementary angles. Side  $a$  is opposite of angle  $\alpha$ , and side  $b$  is opposite of angle  $\beta$ . Notice that we can also describe side  $b$  as adjacent to angle  $\alpha$  and side  $a$  as adjacent to angle  $\beta$ . Therefore,

$$\sin \alpha = \frac{\text{opposite}}{\text{hypotenuse}} = \frac{a}{c} \quad \text{and} \quad \cos \beta = \frac{\text{adjacent}}{\text{hypotenuse}} = \frac{a}{c}$$

Thus we can conclude that

$$\sin \alpha = \frac{a}{c} = \cos \beta$$

Sine and cosine are called **cofunctions** because of this relationship between these functions and their complementary angles. We can obtain similar relationships for all trigonometric functions:

$$\begin{array}{lll} \sin \alpha = \frac{a}{c} = \cos \beta & \cos \alpha = \frac{b}{c} = \sin \beta & \tan \alpha = \frac{a}{b} = \cot \beta \\ \csc \alpha = \frac{c}{a} = \sec \beta & \sec \alpha = \frac{c}{b} = \csc \beta & \cot \alpha = \frac{b}{a} = \tan \beta \end{array}$$

Since  $\alpha$  and  $\beta$  are complementary angles,  $\alpha + \beta = 90^\circ$ . Rearranging, we get  $\beta = 90^\circ - \alpha$ . Substituting this into our cofunctions and replacing  $\alpha$  with  $\theta$ , we get our **cofunction identities**.

**Definition 1.4.7 Cofunction Identities.** The cofunction identities in degrees are

$$\begin{array}{lll} \sin \theta = \cos(90^\circ - \theta) & \cos \theta = \sin(90^\circ - \theta) & \tan \theta = \cot(90^\circ - \theta) \\ \csc \theta = \sec(90^\circ - \theta) & \sec \theta = \csc(90^\circ - \theta) & \cot \theta = \tan(90^\circ - \theta) \end{array}$$

The cofunction identities in radians are

$$\begin{array}{lll} \sin \theta = \cos\left(\frac{\pi}{2} - \theta\right) & \cos \theta = \sin\left(\frac{\pi}{2} - \theta\right) & \tan \theta = \cot\left(\frac{\pi}{2} - \theta\right) \\ \csc \theta = \sec\left(\frac{\pi}{2} - \theta\right) & \sec \theta = \csc\left(\frac{\pi}{2} - \theta\right) & \cot \theta = \tan\left(\frac{\pi}{2} - \theta\right) \end{array}$$

◇

**Example 1.4.8** The Cofunction Identities explains the symmetry in [Table 1.4.6](#)

$$\begin{array}{ll} \sin 0^\circ = \cos(90^\circ - 0^\circ) = \cos 90^\circ & \sin 60^\circ = \cos(90^\circ - 60^\circ) = \cos 30^\circ \\ \sin 30^\circ = \cos(90^\circ - 30^\circ) = \cos 60^\circ & \sin 90^\circ = \cos(90^\circ - 90^\circ) = \cos 0^\circ \\ \sin 45^\circ = \cos(90^\circ - 45^\circ) = \cos 45^\circ & \end{array}$$

□

**Remark 1.4.9 Patterns in the Trigonometric Table.** Learning the values of the trigonometric functions in this table can increase your confidence and efficiency in trigonometry. To help remember the values of sine and cosine, we utilize cofunctions and also write them in the form  $\sqrt{\cdot}/2$

$\theta$	$\theta$	$\sin \theta$	$\cos \theta$
0	$0^\circ$	$\sqrt{0}/2$	$\sqrt{4}/2$
$\pi/6$	$30^\circ$	$\sqrt{1}/2$	$\sqrt{3}/2$
$\pi/4$	$45^\circ$	$\sqrt{2}/2$	$\sqrt{2}/2$
$\pi/3$	$60^\circ$	$\sqrt{3}/2$	$\sqrt{1}/2$
$\pi/2$	$90^\circ$	$\sqrt{4}/2$	$\sqrt{0}/2$

which simplifies to the values in [Table 1.4.6](#).

### 1.4.4 Using a Calculator

Sometimes you may encounter an angle other than the special angles described above. In this case, you will have to use a calculator.

First, be sure that your angle is either in degrees or radians, depending on the problem, refer to your calculator's manual for instructions. Most calculators will have a special button for the sine, cosine, and tangent functions. Depending on your calculator, you may see the following keys for

Function	Calculator Key
sine	<i>SIN</i>
cosine	<i>COS</i>
tangent	<i>TAN</i>

To calculate cosecant, secant, and cotangent, you will need to use the identity

$$\csc \theta = \frac{1}{\sin \theta}, \quad \sec \theta = \frac{1}{\cos \theta}, \quad \cot \theta = \frac{1}{\tan \theta}$$

Answers produced by calculators are *estimates* and you should pay close attention to see if the question is asking for the exact solution or a decimal approximation. For example, if you need to calculate  $\sin 45^\circ = \frac{\sqrt{2}}{2}$ , the calculator may give the answer as  $\sin 45^\circ \approx 0.70710678$ , which is a decimal approximation since the actual solution goes on forever. *Unless stated otherwise, answers in the book should be exact*, e.g.  $\frac{\sqrt{2}}{2}$  and not 0.70710678.

**Example 1.4.10** Use a calculator to evaluate

(a)  $\sin 22^\circ$

**Solution.** Before proceeding, we confirm that our calculator is set to either degree or radian mode. Additionally, for the sake of simplicity, we will round our answers to four decimal places.


Input: *SIN* (22); Output: 0.3746

(b)  $\cos 5^\circ$ **Solution.** Input:  $COS (5)$ ; Output: 0.9962(c)  $\cot 53^\circ$ **Solution.** Since most calculators do not have a key for cotangent, we Input:  $(1/ TAN (53))$ ; Output:  $\frac{1}{1.3270} \approx 0.7536$ (d)  $\cos 5 \text{ rad}$ **Solution.** Since this problem uses radians, we must change the mode on our calculator then Input:  $COS (5)$ ; Output: 0.2837.

□

**Remark 1.4.11** Observe that  $\cos 5^\circ \neq \cos 5 \text{ rad}$ . This emphasizes the significance of verifying whether your calculator is in degree or radian mode.

### 1.4.5 Solving Right Triangles



Consider the following right triangle where side  $a$  is opposite angle  $\alpha$ , side  $b$  is opposite angle  $\beta$ , and side  $c$  is the hypotenuse. Since  $\alpha$  and  $\beta$  are complementary angles, we have

$$\alpha + \beta = 90^\circ$$

Additionally, by the Pythagorean Theorem, we have

$$a^2 + b^2 = c^2$$


**Definition 1.4.12** To **solve a triangle** is the process of determining the values for all three lengths of its sides and the measures of all three angles, based on provided information about the triangle. ◇

**Remark 1.4.13 Solving Right Triangles.** In solving a right triangle, the following relationships are useful:

$$\alpha + \beta = 90^\circ, \quad a^2 + b^2 = c^2$$

**Example 1.4.14** Solve the right triangle. Round your answer to two decimal places.





**Solution.** Given that this is a right triangle, we already know one angle is  $90^\circ$ , and we have an additional angle of  $50^\circ$  along with an adjacent side length of 16. To solve this triangle, we need to determine the values of sides  $a$ ,  $c$ , and  $\beta$ . We begin by finding the measure of angle  $\beta$ . Since  $50^\circ + \beta = 90^\circ$  we have

$$\beta = 90^\circ - 50^\circ = 40^\circ$$

Next, we will solve for side  $a$ . Using the angle  $50^\circ$ , where the adjacent side is 16 and side  $a$  is the side opposite to the angle, we can apply the tangent function, which relates the opposite and adjacent sides:

$$\tan 50^\circ = \frac{a}{16}$$

Multiplying both sides by 16 we get

$$a = 16 \cdot \tan 50^\circ \approx 19.07$$

Using the Pythagorean Theorem, we get

$$c^2 = 16^2 + 19.07^2 \approx 619.66$$


Thus

$$c \approx \sqrt{619.66} \approx 24.89$$


□

### 1.4.6 Solving Applied Problems

**Example 1.4.15 Deviation.** We are now ready to calculate the deviation example proposed at the start of this section. In an average day of sailing, a wa'a sails for 120 nautical miles (NM). If Hikianalia is supposed to sail in the direction of Hikina (East), but currents have deviated her course by one house so she actually sailed in the house Lā, how far off the course has Hikianalia deviated?



**Solution.** From the Star Compass (Figure 1.1.4), the house Lā is one house ( $11.25^\circ$ ) from Hikina. If we let  $y$  denote the distance deviated from the reference course, our right triangle becomes:



Since we know the hypotenuse of the triangle and want to find the side opposite of the angle, we will use the sine function:


$$\sin 11.25^\circ = \frac{\text{opposite}}{\text{hypotenuse}} = \frac{y}{120 \text{ NM}}$$

multiplying both sides by 120, we get


$$y = 120 \cdot \sin 11.25^\circ \text{ NM} \approx 23.4 \text{ NM}$$

So Hikianalia has deviated 23.4 nautical miles north from the reference course.  $\square$

**Example 1.4.16 Solar panels.** Solar panels harness the sun’s energy to generate electricity, and for optimal energy output, they should be oriented perpendicularly to the sun’s light. The sun’s angle of elevation varies based on latitude, and in Hawai’i, for instance, south-facing solar panels are recommended to have a pitch of  $21^\circ$  to align with the sun’s rays. When installing a solar panel, determining its pitch might pose challenges. Instead of measuring the angle directly, an alternative approach involves measuring the height of the panel’s top. What height should a south-facing solar panel, measuring 65 inches in length, be installed at to achieve the desired angle of  $21^\circ$ ? Round your answer to the nearest tenth of an inch.



**Solution.** We begin by drawing the triangle.



Since we know the desired angle of pitch of the solar panel and the length of the panel, we can set up the following equation


$$\begin{aligned}\sin 21^\circ &= \frac{\text{opposite}}{\text{hypotenuse}} = \frac{\text{height}}{65 \text{ in}} \\ \text{height} &= 65 \text{ in} \cdot \sin 21^\circ \approx 23.3 \text{ in}\end{aligned}$$

Thus, when installing a solar panel in Hawai'i, the top of the solar panel should be positioned 23.3 inches above the bottom to optimize energy output.  $\square$

### 1.4.7 Exercises


**Exercise Group.** Find the exact values of the six trigonometric functions of the angle  $\theta$  in each triangle.

1.




**Answer.**  $\sin \theta = \frac{4}{5}$ ,  
 $\cos \theta = \frac{3}{5}$ ,  $\tan \theta = \frac{4}{3}$ ,  
 $\csc \theta = \frac{5}{4}$ ,  $\sec \theta = \frac{5}{3}$ ,  $\cot \theta = \frac{3}{4}$

2.




**Answer.**  $\sin \theta = \frac{5\sqrt{74}}{74}$ ,  
 $\cos \theta = \frac{7\sqrt{74}}{74}$ ,  $\tan \theta = \frac{5}{7}$ ,  
 $\csc \theta = \frac{\sqrt{74}}{5}$ ,  $\sec \theta = \frac{\sqrt{74}}{7}$ ,  
 $\cot \theta = \frac{7}{5}$

3.




**Answer.**  $\sin \theta = \frac{3\sqrt{13}}{13}$ ,  
 $\cos \theta = \frac{2\sqrt{13}}{13}$ ,  $\tan \theta = \frac{3}{2}$ ,  
 $\csc \theta = \frac{\sqrt{13}}{3}$ ,  $\sec \theta = \frac{\sqrt{13}}{2}$ ,  
 $\cot \theta = \frac{2}{3}$

4.




**Answer.**  $\sin \theta = \frac{5}{13}$ ,  
 $\cos \theta = \frac{12}{13}$ ,  $\tan \theta = \frac{5}{12}$ ,  
 $\csc \theta = \frac{13}{5}$ ,  $\sec \theta = \frac{13}{12}$ ,  
 $\cot \theta = \frac{12}{5}$

5.




**Answer.**  $\sin \theta = \frac{2\sqrt{5}}{5}$ ,  
 $\cos \theta = \frac{\sqrt{5}}{5}$ ,  $\tan \theta = 2$ ,  
 $\csc \theta = \frac{\sqrt{5}}{2}$ ,  $\sec \theta = \sqrt{5}$ ,  
 $\cot \theta = \frac{1}{2}$

6.




**Answer.**  $\sin \theta = \frac{\sqrt{2}}{2}$ ,  
 $\cos \theta = \frac{\sqrt{2}}{2}$ ,  $\tan \theta = 1$ ,  
 $\csc \theta = \sqrt{2}$ ,  $\sec \theta = \sqrt{2}$ ,  
 $\cot \theta = 1$

7.



**Answer.**  $\sin \theta = \frac{\sqrt{15}}{5}$ ,  
 $\cos \theta = \frac{\sqrt{10}}{5}$ ,  $\tan \theta = \frac{\sqrt{6}}{2}$ ,  
 $\csc \theta = \frac{\sqrt{15}}{3}$ ,  $\sec \theta = \frac{\sqrt{10}}{2}$ ,  
 $\cot \theta = \frac{\sqrt{6}}{3}$

8.




**Answer.**  $\sin \theta = \frac{2\sqrt{13}}{13}$ ,  
 $\cos \theta = \frac{3\sqrt{13}}{13}$ ,  $\tan \theta = \frac{2}{3}$ ,  
 $\csc \theta = \frac{\sqrt{13}}{2}$ ,  $\sec \theta = \frac{\sqrt{13}}{3}$ ,  
 $\cot \theta = \frac{3}{2}$

**Exercise Group.** For each of the following problems, calculate

- (a)  $\cos \alpha$  and  $\sin \beta$ ;
- (b)  $\tan \alpha$  and  $\cot \beta$ ;
- (c)  $\csc \alpha$  and  $\sec \beta$


9.



**Answer.**

- (a)  $\cos \alpha = \sin \beta = \frac{5\sqrt{34}}{34}$ ;
- (b)  $\tan \alpha = \cot \beta = \frac{3}{5}$ ;
- (c)  $\csc \alpha = \sec \beta = \frac{\sqrt{34}}{3}$

10.




**Answer.**

- (a)  $\cos \alpha = \sin \beta = \frac{7}{8}$ ;
- (b)  $\tan \alpha = \cot \beta = \frac{\sqrt{15}}{7}$ ;
- (c)  $\csc \alpha = \sec \beta = \frac{8\sqrt{15}}{15}$

**Exercise Group.** Use the Cofunction Identities to determine the value of  $\theta$


- |                                    |   |  |
|------------------------------------|---|--|
| 11. $\sin 28^\circ = \cos \theta$  | 12. $\cos 74^\circ = \sin \theta$       | 13. $\tan 52^\circ = \cot \theta$        |
| <b>Answer.</b> $\theta = 62^\circ$ | <b>Answer.</b> $\theta = 16^\circ$      | <b>Answer.</b> $\theta = 62^\circ$       |
| 14. $\sec 87^\circ = \csc \theta$  | 15. $\sin \frac{3\pi}{8} = \cos \theta$ | 16. $\cot \frac{2\pi}{5} = \tan \theta$  |
| <b>Answer.</b> $\theta = 3^\circ$  | <b>Answer.</b> $\theta = \frac{\pi}{8}$ | <b>Answer.</b> $\theta = \frac{\pi}{10}$ |

**Exercise Group.** In [Example 1.4.15](#), we determined that when a wa'a sails for one day (120 nautical miles) and deviates from its course by 1 House, the resulting deviation from the reference course is 23.4 NM. Now, calculate the deviations ( $x$ ) for the remaining 7 angles. Round your answer to the nearest tenth of a nautical mile. Remember that one house corresponds to  $11.25^\circ$ .




17. 2 Houses ( $11.25^\circ \times 2 = 22.5^\circ$ );  $x_2$   
**Answer.**  $x_2 = 45.9$  NM
18. 3 Houses ( $11.25^\circ \times 3 = 33.75^\circ$ );  $x_3$   
**Answer.**  $x_3 = 66.7$  NM
19. 4 Houses ( $11.25^\circ \times 4 = 45^\circ$ );  $x_4$   
**Answer.**  $x_4 = 84.5$  NM
20. 5 Houses ( $11.25^\circ \times 5 = 56.25^\circ$ );  $x_5$   
**Answer.**  $x_5 = 99.8$  NM
21. 6 Houses ( $11.25^\circ \times 6 = 67.5^\circ$ );  $x_6$   
**Answer.**  $x_6 = 110.9$  NM
22. 7 Houses ( $11.25^\circ \times 7 = 78.75^\circ$ );  $x_7$   
**Answer.**  $x_7 = 117.7$  NM
23. 8 Houses ( $11.25^\circ \times 8 = 90^\circ$ );  $x_8$   
**Answer.**  $x_8 = 120$  NM

**Exercise Group.** In [Exercise 1.4.7.17–23](#) we determined the deviation of a wa'a following a day of sailing (120 nautical miles). Your task now is to calculate the distance the wa'a has progressed along the reference course (north) for each deviation, denoted as  $y$ . Round your answer to the nearest tenth of a nautical mile and remember that one house corresponds to  $11.25^\circ$ .



24. 1 House ( $11.25^\circ \times 1 = 11.25^\circ$ );  $y_1$   
**Answer.**  $y_1 = 117.7$  NM
25. 2 Houses ( $11.25^\circ \times 2 = 22.5^\circ$ );  $y_2$   
**Answer.**  $y_2 = 110.9$  NM
26. 3 Houses ( $11.25^\circ \times 3 = 33.75^\circ$ );  $y_3$   
**Answer.**  $y_3 = 99.8$  NM
27. 4 Houses ( $11.25^\circ \times 4 = 45^\circ$ );  $y_4$   
**Answer.**  $y_4 = 84.5$  NM
28. 5 Houses ( $11.25^\circ \times 5 = 56.25^\circ$ );  $y_5$   
**Answer.**  $y_5 = 66.7$  NM
29. 6 Houses ( $11.25^\circ \times 6 = 67.5^\circ$ );  $y_6$   
**Answer.**  $y_6 = 45.9$  NM
30. 7 Houses ( $11.25^\circ \times 7 = 78.75^\circ$ );  $y_7$   
**Answer.**  $y_7 = 23.4$  NM

**Exercise Group.** One way to determine your bearing on a canoe is by observing and comparing the positions of celestial and other markers relative to your canoe. To facilitate this, you can mark the locations of the Star Compass on the opposite railings from the navigator's seat in the back corner of the canoe. However, since the Star Compass is circular and the canoe is rectangular, accurately placing the markings can be challenging.



When the navigator occupies the port stern (back left) corner of the deck, markers indicating the boundaries between houses can be placed on the corresponding railings on the bow (front) and starboard (right) sides of the canoe. For each value of  $\theta$ , calculate the distance along the starboard railing ( $y$ ) or bow railing ( $x$ ) for a canoe with dimensions  $l = 50$  ft and  $w = 20$  ft. Round your answers to three decimal places.

**31.**  $\theta = 5.625^\circ$ ;  $y_1$

**Answer.**  $y_1 = 1.970$  ft

**33.**  $\theta = 28.125^\circ$ ;  $y_3$

**Answer.**  $y_3 = 10.69$  ft

**35.**  $\theta = 50.625^\circ$ ;  $y_5$

**Answer.**  $y_5 = 24.370$  ft

**37.**  $\theta = 73.125^\circ$ ;  $x_7$

**Answer.**  $x_7 = 15.167$  ft

**32.**  $\theta = 16.875^\circ$ ;  $y_2$

**Answer.**  $y_2 = 6.067$  ft

**34.**  $\theta = 39.375^\circ$ ;  $y_4$

**Answer.**  $y_4 = 16.414$  ft

**36.**  $\theta = 61.875^\circ$ ;  $y_6$


**Answer.**  $y_6 = 37.417$  ft

**38.**  $\theta = 84.375^\circ$ ;  $x_8$


**Answer.**  $x_8 = 4.925$  ft

**Exercise Group.** Use the right triangle (not drawn to scale) provided below to solve for the given information. Round your solutions to two decimal places.





39.  $a = 5$ ,  $\beta = 35^\circ$ . Find  $b$ ,  $c$ , and  $\alpha$   
**Answer.**  $b = 3.50$ ,  $c = 6.10$ ,  $\alpha = 55^\circ$
40.  $b = 12$ ,  $\beta = 23^\circ$ . Find  $a$ ,  $c$ , and  $\alpha$   
**Answer.**  $a = 28.27$ ,  $c = 30.71$ ,  $\alpha = 67^\circ$
41.  $b = 7$ ,  $\alpha = 75^\circ$ . Find  $a$ ,  $c$ , and  $\beta$   
**Answer.**  $a = 26.12$ ,  $c = 27.05$ ,  $\beta = 15^\circ$
42.  $c = 4$ ,  $\beta = 50^\circ$ . Find  $a$ ,  $b$ , and  $\alpha$   
**Answer.**  $a = 3.06$ ,  $b = 2.57$ ,  $\alpha = 40^\circ$
43.  $c = 10$ ,  $\alpha = 18^\circ$ . Find  $a$ ,  $b$ , and  $\beta$   
**Answer.**  $a = 3.09$ ,  $b = 9.51$ ,  $\beta = 72^\circ$
44.  $a = 6$ ,  $\alpha = 38^\circ$ . Find  $b$ ,  $c$ , and  $\beta$   
**Answer.**  $b = 7.68$ ,  $c = 9.75$ ,  $\beta = 52^\circ$
45. A wa'a sails in the direction of the house Nālani Ho'olua for one day, covering 120 nautical miles. How many nautical miles has the wa'a traveled north? How many miles has the wa'a traveled west? To calculate the angle  $\theta$ , refer to the Star Compass (Figure 1.1.4) to determine the number of houses, and use the fact that one house is  $11.25^\circ$ .




**Answer.** 66.7 NM west; 99.8 NM north

46. **Movement of Sand.** The movement of sand on a beach is a dynamic process influenced by various factors, such as waves. When waves approach the shore at an angle, they lead to the shifting of sand. During the *swash*

phase, as the wave crashes onto the shore, water and sediment move onto the beach following the wave's angle. Subsequently, gravity propels the water and sediment back into the ocean, perpendicular to the shoreline, in a process known as *backwash*. This interplay of swash and backwash creates a zig-zag pattern called *longshore drift*.

Certain beaches undergo seasonal changes in wave direction. Some experience waves from one direction in one season and from another direction in the next, while those predominantly receiving waves from a single direction might accumulate sand in specific areas.

Calculate how far along the shore a single grain of sand moves after a wave breaks at a  $60^\circ$  angle and travels onto the shore for 10 ft before receding back into the ocean.




**Answer.** 5ft

**Exercise Group.** Between 2013 and 2017, Hōkūle‘a completed a global circumnavigation with a mission *mālama honua* - “care for Island Earth” and to foster a sense of ‘*ohana* (“family”) for all people and places. This remarkable voyage spanned 40,000 nautical miles and made stops at over 150 ports across 18 nations.

Throughout this voyage, Earth’s rotation occurs around an axis that extends from the North Pole to the South Pole. The rotation imparts an angular speed and linear velocity to every point on Earth. Assuming Earth completes one rotation within 24 hours and treating Earth as a perfect sphere with a radius of  $R = 4,000$  miles, we can calculate the following parameters for each of the Mālama Honua voyage’s ports, given their latitudes ( $\phi$ ):

- Calculate  $r$ , the distance from the port to Earth’s Axis of Rotation (in miles, rounded to one decimal place).
- Calculate  $\omega$ , the angular velocity (in radians per hour, rounded to four decimal places).
- Calculate  $v$ , the linear speed (in miles per hour, rounded to the nearest whole number).



47. Hilo, Hawai'i ( $19.7216^\circ$  N)

**Answer.**

- (a)  $r = 3,8765.4$  miles;  
 (b)  $\omega = 0.2618$  rad/hr;  
 (c)  $v = 986$  mi/hr

49. Apia, Samoa ( $13.8507^\circ$  S)

**Answer.**

- (a)  $r = 3,883.7$  miles;  
 (b)  $\omega = 0.2618$  rad/hr;  
 (c)  $v = 1,017$  mi/hr

51. Sydney, Australia ( $33.8688^\circ$  S)

**Answer.**

- (a)  $r = 3,321.3$  miles;  
 (b)  $\omega = 0.2618$  rad/hr;  
 (c)  $v = 870$  mi/hr

53. Port Louis, Mauritius ( $20.1609^\circ$  S)

**Answer.**

- (a)  $r = 3,754.9$  miles;  
 (b)  $\omega = 0.2618$  rad/hr;  
 (c)  $v = 983$  mi/hr

48. Papeete, Tahiti ( $17.5325^\circ$  S)

**Answer.**

- (a)  $r = 3,814.2$  miles;  
 (b)  $\omega = 0.2618$  rad/hr;  
 (c)  $v = 999$  mi/hr

50. Waitangi, Aotearoa ( $35.2683^\circ$  S)

**Answer.**

- (a)  $r = 32,65.8$  miles;  
 (b)  $\omega = 0.2618$  rad/hr;  
 (c)  $v = 855$  mi/hr

52. Bali, Indonesia ( $8.4095^\circ$  S)

**Answer.**

- (a)  $r = 3,957.0$  miles;  
 (b)  $\omega = 0.2618$  rad/hr;  
 (c)  $v = 1,036$  mi/hr

54. Cape Town, South Africa ( $33.9249^\circ$  S)

**Answer.**

- (a)  $r = 3,319.1$  miles;  
 (b)  $\omega = 0.2618$  rad/hr;  
 (c)  $v = 869$  mi/hr

55. Natal, Brazil ( $5.7842^\circ$  S)

**Answer.**

- (a)  $r = 3,979.6$  miles;
- (b)  $\omega = 0.2618$  rad/hr;
- (c)  $v = 1,042$  mi/hr

57. Yarmouth, Nova Scotia ( $43.8379^\circ$  N)

**Answer.**

- (a)  $r = 3,792.7$  miles;
- (b)  $\omega = 0.2618$  rad/hr;
- (c)  $v = 755$  mi/hr

59. Galapagos Islands ( $0.9538^\circ$  S)

**Answer.**

- (a)  $r = 3,999.5$  miles;
- (b)  $\omega = 0.2618$  rad/hr;
- (c)  $v = 1,047$  mi/hr

56. Necker, British Virgin Islands ( $18.5268^\circ$  N)

**Answer.**

- (a)  $r = 3,792.7$  miles;
- (b)  $\omega = 0.2618$  rad/hr;
- (c)  $v = 993$  mi/hr

58. Balboa, Panama ( $8.9614^\circ$  N)

**Answer.**

- (a)  $r = 3,951.2$  miles;
- (b)  $\omega = 0.2618$  rad/hr;
- (c)  $v = 1,034$  mi/hr

60. Rapa Nui ( $27.1127^\circ$  S)

**Answer.**


- (a)  $r = 3,560.5$  miles;
- (b)  $\omega = 0.2618$  rad/hr;
- (c)  $v = 932$  mi/hr

## 1.5 Trigonometric Functions of Any Angles

Now that we have been introduced to the six trigonometric functions for special angles in the first quadrant, we can explore their properties across all quadrants.

### 1.5.1 Determine the Signs of the Trigonometric Functions Based on its Quadrant

Let  $P(x, y)$  be a point on the circle. The signs of the six trigonometric functions vary depending on the quadrant in which  $P(x, y)$  lies in.



**Example 1.5.1** Let  $P(x, y)$  is in Quadrant II. Determine the signs for each of the six trigonometric functions.

**Solution.** Since we are in Quadrant II,  $x < 0$  and  $y > 0$ . Note that  $r > 0$ . Then we have


$$\begin{aligned} \sin \theta &= \frac{y}{r} = \frac{(+)}{(+)} = (+) & \cos \theta &= \frac{x}{r} = \frac{(-)}{(+)} = (-) & \tan \theta &= \frac{y}{x} = \frac{(+)}{(-)} = (-) \\ \csc \theta &= \frac{r}{y} = \frac{(+)}{(+)} = (+) & \sec \theta &= \frac{r}{x} = \frac{(+)}{(-)} = (-) & \cot \theta &= \frac{x}{y} = \frac{(-)}{(+)} = (-) \end{aligned}$$

□

You can check the remaining quadrants using a similar approach. [Table 1.5.2](#) and [Figure 1.5.3](#) provide a list of the signs of the six trigonometric functions for each quadrant.

**Table 1.5.2** Signs of the trigonometric functions

Quadrant	Positive Functions	Negative Functions
I	all	none
II	sin, csc	cos, sec, tan, cot
III	tan, cot	sin, csc, cos, sec
IV	cos, sec	sin, csc, tan, cot



**Figure 1.5.3** Signs of trigonometric functions


**Example 1.5.4** If  $\sin \theta < 0$  and  $\cos \theta > 0$ , what quadrant does  $\theta$  lie in?

**Solution.** Since  $\sin \theta < 0$ , then  $\theta$  is either in Quadrant III or IV. However, we also have  $\cos \theta > 0$  which means that  $\theta$  is either in Quadrant I or IV. Thus the only quadrant that satisfied both conditions is Quadrant IV.  $\square$

Mnemonic devices for remembering the quadrants in which the trigonometric functions are positive are

- “A Smart Trig Class”
- “All Students Take Calculus”

which correspond to “All Sin Tan Cos.”



**Example 1.5.5** Let  $\sin \theta = -\frac{12}{13}$  and  $\cos \theta = -\frac{5}{13}$ . Compute the exact values of the remaining trigonometric functions of  $\theta$  using identities.

**Solution.** Since  $\sin \theta < 0$  and  $\cos \theta < 0$ , we refer to [Table 1.5.2](#) and see that  $\theta$  is in Quadrant III. From [Table 1.5.2](#) we know  $\tan \theta > 0$ ,  $\csc \theta < 0$ ,  $\sec \theta < 0$ ,

$\cot \theta > 0$ . From the Quotient Identity (Definition 1.4.3), we have

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\frac{12}{13}}{\frac{5}{13}} = \frac{12}{5}$$

Next, using the Reciprocal Identities (Definition 1.4.2), we get

$$\csc \theta = \frac{1}{\sin \theta} = \frac{1}{\frac{12}{13}} = \frac{13}{12}$$

$$\sec \theta = \frac{1}{\cos \theta} = \frac{1}{\frac{5}{13}} = \frac{13}{5}$$

$$\cot \theta = \frac{1}{\tan \theta} = \frac{1}{\frac{12}{5}} = \frac{5}{12}$$

□

### 1.5.2 Reference Angles

Now that we can determine the signs of trigonometric functions, we will demonstrate how the value of any trigonometric function at any angle can be found from its value in Quadrant I (between  $0^\circ$  and  $90^\circ$  or  $0$  and  $\frac{\pi}{2}$ ).


**Definition 1.5.6** Let  $t$  be a real number. A **reference angle**,  $t'$ , is the acute angle ( $< 90^\circ$ ) formed by the terminal side of angle  $t$  and the  $x$ -axis. In other words, it is the shortest distance along the unit circle measured between the terminal side and the  $x$ -axis. Angles in Quadrant I are their own reference angles.  $\diamond$

**Remark 1.5.7 Calculating the reference angle.** To calculate the reference angle  $t'$  for a given angle  $t$ :

- In radians, if  $t > 2\pi$  or if  $t < 0$ , add or subtract multiples of  $2\pi$  to obtain a coterminal angle between  $0$  and  $2\pi$ . Then, find the reference angle.
- In degrees, if  $t > 360^\circ$  or  $t < 0^\circ$ , add or subtract multiples of  $360^\circ$  to obtain a coterminal angle between  $0^\circ$  and  $360^\circ$ . Then, find the reference angle.

Quadrant I:


$$t' = t$$



Quadrant II:

$$t' = \pi - t$$


$$t' = 180^\circ - t$$



Quadrant III:

$$t' = t - \pi$$


$$t' = t - 180^\circ$$



Quadrant IV:


$$t' = 2\pi - t$$

$$t' = 360^\circ - t$$




**Example 1.5.8** Find the reference angle for each value of  $t$

(a)  $t = \frac{\pi}{3}$




**Solution.** The angle  $t = \frac{\pi}{3}$  is in the first quadrant and so it is its own reference angle:  $t = t' = \frac{\pi}{3}$

(b)  $t = \frac{3\pi}{4}$



**Solution.** From the figure, we see that the shortest distance to the  $x$ -axis is in the direction of  $\pi$ . We see that  $t' + \frac{3\pi}{4} = \pi$  so  $t' = \pi - \frac{3\pi}{4} = \frac{\pi}{4}$ .

(c)  $t = -\frac{3\pi}{4}$




**Solution 1.** Since  $t < 0$ , we can add  $2\pi$  to get  $-\frac{3\pi}{4} + 2\pi = \frac{5\pi}{4}$ . From the formula, we get  $t' = \frac{5\pi}{4} - \pi = \frac{\pi}{4}$ .




**Solution 2.** Since  $-\frac{3\pi}{4}$  spans only two quadrants counterclockwise, we can treat it similarly to an angle in Quadrant II. By the previous problem,  $t' + \frac{3\pi}{4} = \pi$  so  $t' = \pi - \frac{3\pi}{4} = \frac{\pi}{4}$ .

(d)  $t = 240^\circ$




**Solution.** From the figure we see the shortest distance to the  $x$ -axis is towards  $180^\circ$ . We observe that  $240^\circ - t' = 180^\circ$  so  $t' = 240^\circ - 180^\circ = 60^\circ$

(e)  $t = \frac{11\pi}{6}$



**Solution.** From the figure, we see the shortest distance to the  $x$ -axis is towards  $2\pi$ . We observe that  $t' + \frac{11\pi}{6} = 2\pi$  so  $t' = 2\pi - \frac{11\pi}{6} = \frac{\pi}{6}$ .

(f)  $t = \frac{2\pi}{3}$




**Solution.** From the figure, we see the shortest distance to the  $x$ -axis is towards  $\pi$ . We observe that  $t' + \frac{2\pi}{3} = \pi$  so  $t' = \pi - \frac{2\pi}{3} = \frac{\pi}{3}$ .

□

**Remark 1.5.9 Calculate an angle in standard position given its quadrant and reference angle.** To calculate an angle in standard position,  $t$ , given the quadrant that  $t$  lies in and the reference angle  $t'$ ,

Quadrant I:


$$t = t'$$



Quadrant II:

$$t = \pi - t'$$


$$t = 180^\circ - t'$$



Quadrant III:

$$t = t' - \pi$$


$$t = t' - 180^\circ$$




Quadrant IV:

$$t = 2\pi - t'$$

$$t = 360^\circ - t'$$




For radians only: If the reference angle (in radians) is of the form  $t' = \frac{a\pi}{b}$ , then the associated angle in standard position,  $t$ , can be calculated by




**Example 1.5.10 Calculate an angle given its reference angle and quadrant.** Given a reference angle,  $t$ , compute the associated angle in standard position for Quadrant II, III, and IV.

(a)  $t' = \frac{\pi}{6}$



(i) Quadrant II


**Solution 1.** In Quadrant II, the associated angle is  $t = \pi - \frac{\pi}{6} = \frac{6\pi}{6} - \frac{\pi}{6} = \frac{5\pi}{6}$



**Solution 2.** Since  $t' = \frac{\pi}{6} = \frac{1\pi}{6}$ , then  $t' = \frac{(6-1)\pi}{6} = \frac{5\pi}{6}$

(ii) Quadrant III


**Solution 1.** In Quadrant III, the associated angle is  $t = \pi + \frac{\pi}{6} = \frac{6\pi}{6} + \frac{\pi}{6} = \frac{7\pi}{6}$



**Solution 2.**  $t' = \frac{(6+1)\pi}{6} = \frac{7\pi}{6}$


(iii) Quadrant IV

**Solution 1.** In Quadrant IV, the associated angle is  $t = 2\pi - \frac{\pi}{6} = \frac{12\pi}{6} - \frac{\pi}{6} = \frac{11\pi}{6}$




**Solution 2.**  $t' = \frac{(2 \cdot 6 - 1)\pi}{6} = \frac{11\pi}{6}$

(b)  $t' = 45^\circ$




(i) Quadrant II

**Solution.** In Quadrant II, the associated angle is  $t = 180^\circ - 45^\circ = 135^\circ$




(ii) Quadrant III

**Solution.** In Quadrant III, the associated angle is  $t = 180^\circ + 45^\circ = 225^\circ$



(iii) Quadrant IV


**Solution.** In Quadrant IV, the associated angle is  $t = 360^\circ - 45^\circ = 315^\circ$



□

### 1.5.3 Evaluating Trigonometric Functions Using Reference Angles


To evaluate trigonometric functions in any quadrant using reference angles, we begin with an angle,  $\theta$ , that lies in Quadrant II. When evaluating  $\sin \theta$  and  $\cos \theta$ , we begin by plotting  $\theta$  in standard position and then proceed to determine and draw its corresponding reference angle,  $\theta'$ .



By definition we know that

$$\sin \theta = \frac{y}{r}; \quad \cos \theta = \frac{x}{r}$$

Next, we draw the reference angle,  $\theta'$  in standard position



We now have

$$\sin \theta' = \frac{y'}{r}; \quad \cos \theta' = \frac{x'}{r}.$$

Notice that the  $y$ -coordinates for  $P$  and  $P'$  share the same value, thus  $y = y'$

and we get

$$\sin \theta = \sin \theta'.$$

Similarly, we can see that the  $x$ -coordinates of  $P$  and  $P'$  have opposite values, thus  $x = -x'$  and

$$\cos \theta = -\cos \theta'.$$

You may have noticed that we have two similar triangles, differing only in their  $x$ -coordinates have opposite values. Consequently, the values of each trigonometric function for the two triangles will match, except for a potential distinction in signs. The sign of each function can be deduced by referring to [Table 1.5.2](#). This approach is applicable across all quadrants. To sum up, we now outline the steps for utilizing reference angles to evaluate trigonometric functions.

**Remark 1.5.11 Steps for Evaluating Trigonometric Functions Using Reference Angles.** The values of a trigonometric function for a specific angle are equivalent to the values of the same trigonometric function for the reference angle, with a potential distinction in sign. To compute the value of a trigonometric function for any angle, use the following steps


1. Draw the angle in standard position.
2. Determine the reference angle associated with the angle.
3. Evaluate the trigonometric function at the reference angle.
4. Use [Table 1.5.2](#) and the quadrant of the original angle to determine the appropriate sign for the function.

**Example 1.5.12** Use the reference angle associated with the given angle to find the exact value of

- (a)  $\cos 210^\circ$

**Solution.** We will use the steps for evaluating trigonometric functions using reference angles.

- (a) First we draw the angle



(b) The reference angle is

$$\theta' = 210^\circ - 180^\circ = 30^\circ$$

(c)  $\cos 30^\circ = \frac{\sqrt{3}}{2}$


(d) Since  $210^\circ$  lies in Quadrant III, we know that  $\cos \theta < 0$ , so

$$\cos 210^\circ = -\frac{\sqrt{3}}{2}$$

(b)  $\tan \frac{7\pi}{4}$

**Solution.** We will use the steps for evaluating trigonometric functions using reference angles.

(a) First we draw the angle



(b) The reference angle is

$$2\pi - \frac{7\pi}{4} = \frac{8\pi}{4} - \frac{7\pi}{4} = \frac{\pi}{4}$$

(c)  $\tan \frac{\pi}{4} = 1$

(d) Since  $\frac{7\pi}{4}$  lies in Quadrant IV, we know that  $\tan \theta < 0$ , so

$$\tan \frac{7\pi}{4} = -1$$


□

**Example 1.5.13** Calculate  $\sin \theta$  and  $\cos \theta$  if  $\theta = \frac{20\pi}{3}$

**Solution.**

1. First we draw the angle





**Figure 1.5.14** The angle  $\theta = \frac{20\pi}{3}$  makes three rotations before ending in Quadrant II.

2. To obtain the reference angle, we first subtract multiples of  $2\pi$  from  $\theta$  to obtain a coterminal angle between 0 and  $2\pi$ :

$$\begin{array}{rcl} \frac{20\pi}{3} - 2\pi = & \frac{20\pi}{3} - \frac{6\pi}{3} = & \frac{14\pi}{3} \\ \frac{20\pi}{3} - 4\pi = & \frac{20\pi}{3} - \frac{12\pi}{3} = & \frac{8\pi}{3} \\ \frac{20\pi}{3} - 6\pi = & \frac{20\pi}{3} - \frac{18\pi}{3} = & \frac{2\pi}{3} \end{array}$$

From [Example 1.5.8](#), the reference angle for  $\frac{2\pi}{3}$  is  $\theta' = \frac{\pi}{3}$

3.  $\sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}$  and  $\cos \frac{\pi}{3} = \frac{1}{2}$   
 4. Since  $\frac{20\pi}{3}$  lies in Quadrant II, we know that  $\sin \theta > 0$  and  $\cos \theta < 0$ , so

$$\sin \frac{20\pi}{3} = \frac{\sqrt{3}}{2}; \quad \cos \frac{20\pi}{3} = -\frac{1}{2}$$

□

### 1.5.4 Periodic Functions

In [Figure 1.5.14](#) of [Example 1.5.13](#), point  $P$  corresponds to the angle  $\frac{20\pi}{3}$ . To determine the reference angle, we subtracted multiples of  $2\pi$ . Each iteration of  $2\pi$  retraces the unit circle back to the point  $P$ , resulting in a coterminal angle. Therefore

$$\sin \frac{2\pi}{3} = \sin \frac{8\pi}{3} = \sin \frac{14\pi}{3} = \sin \frac{20\pi}{3}$$

Rewriting the angles we get

$$\sin \left( \frac{2\pi}{3} + 0 \cdot 2\pi \right) = \sin \left( \frac{2\pi}{3} + 1 \cdot 2\pi \right) = \sin \left( \frac{2\pi}{3} + 2 \cdot 2\pi \right) = \sin \left( \frac{2\pi}{3} + 3 \cdot 2\pi \right)$$

Similarly,

$$\cos\left(\frac{2\pi}{3} + 0 \cdot 2\pi\right) = \cos\left(\frac{2\pi}{3} + 1 \cdot 2\pi\right) = \cos\left(\frac{2\pi}{3} + 2 \cdot 2\pi\right) = \cos\left(\frac{2\pi}{3} + 3 \cdot 2\pi\right)$$

In general, consider an angle  $\theta$  measured in radians and its corresponding point  $P$  on the unit circle. Adding or subtracting integer multiples of  $2\pi$  to  $\theta$  will lead to a point on the unit circle that aligns with  $P$ . Thus, the values of sine and cosine for all angles corresponding to point  $P$  are equivalent. This leads us to the following periodic properties.

**Definition 1.5.15 Periodic Properties.**

$$\sin(\theta + 2\pi k) = \sin \theta \qquad \cos(\theta + 2\pi k) = \cos \theta$$

where  $k$  is any integer. ◇

Functions like these that repeats its values in regular cycles are called *periodic functions*.

**Definition 1.5.16** A function  $f$  is called **periodic** if there exists a positive number  $p$  such that

$$f(\theta + p) = f(\theta)$$

for every  $\theta$ . The smallest number  $p$  is called the **period** of  $f$ . ◇

Sine, cosine, cosecant, and secant repeat their values with a period of  $2\pi$  while tangent and cotangent have a period of  $\pi$ .

**Definition 1.5.17 Periodic Properties.**

$$\begin{array}{lll} \sin(\theta + 2\pi) = \sin \theta & \cos(\theta + 2\pi) = \cos \theta & \tan(\theta + \pi) = \tan \theta \\ \csc(\theta + 2\pi) = \csc \theta & \sec(\theta + 2\pi) = \sec \theta & \cot(\theta + \pi) = \cot \theta \end{array}$$

◇

### 1.5.5 Trigonometric Table

The Trigonometric Identities and reference angles give us the values of trigonometric functions in [Table 1.5.18](#).

**Table 1.5.18** Values of the six trigonometric functions for common angles

$\theta$ (deg)	$\theta$ (rad)	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\csc \theta$	$\sec \theta$	$\cot \theta$
$0^\circ$	0	0	1	0	undef	1	undef
$30^\circ$	$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$	$\sqrt{3}$
$45^\circ$	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	$\sqrt{2}$	$\sqrt{2}$	1
$60^\circ$	$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2	$\frac{\sqrt{3}}{3}$
$90^\circ$	$\frac{\pi}{2}$	1	0	undef	1	undef	0
$120^\circ$	$\frac{2\pi}{3}$	$\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$-\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	-2	$-\frac{\sqrt{3}}{3}$
$135^\circ$	$\frac{3\pi}{4}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	-1	$\sqrt{2}$	$-\sqrt{2}$	-1
$150^\circ$	$\frac{5\pi}{6}$	$\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{3}$	2	$-\frac{2\sqrt{3}}{3}$	$-\sqrt{3}$
$180^\circ$	$\pi$	0	-1	0	undef	-1	undef
$210^\circ$	$\frac{7\pi}{6}$	$-\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	-2	$-\frac{2\sqrt{3}}{3}$	$\sqrt{3}$
$225^\circ$	$\frac{5\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	1	$-\sqrt{2}$	$-\sqrt{2}$	1
$240^\circ$	$\frac{4\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$\sqrt{3}$	$-\frac{2\sqrt{3}}{3}$	-2	$\frac{\sqrt{3}}{3}$
$270^\circ$	$\frac{3\pi}{2}$	-1	0	undef	-1	undef	0
$300^\circ$	$\frac{5\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$-\sqrt{3}$	$-\frac{2\sqrt{3}}{3}$	2	$-\frac{\sqrt{3}}{3}$
$315^\circ$	$\frac{7\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	-1	$-\sqrt{2}$	$\sqrt{2}$	-1
$330^\circ$	$\frac{11\pi}{6}$	$-\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{3}$	-2	$\frac{2\sqrt{3}}{3}$	$-\sqrt{3}$

**Remark 1.5.19 Table Made Easy.** Table 1.5.18 may seem intimidating but if you recognize the symmetry about  $90^\circ$ ,  $180^\circ$ , and  $270^\circ$ , you will only need

to focus on the values for the first quadrant (Table 1.4.6). In fact, you need only produce the values of sine in Quadrant I. Use the Cofunction Identities (Definition 1.4.7) to find the values of cosine. Next, apply the trigonometric identity to find  $\tan \theta = \sin \theta / \cos \theta$ . Finally, use the the Reciprocal Identities (Definition 1.4.2) to produce  $\csc \theta$ ,  $\sec \theta$ , and  $\cot \theta$ .


### 1.5.6 Pythagorean Identities

**Definition 1.5.20 Pythagorean Identities.**

1.  $\sin^2 \theta + \cos^2 \theta = 1$
2.  $1 + \tan^2 \theta = \sec^2 \theta$
3.  $1 + \cot^2 \theta = \csc^2 \theta$

◇

*Proof.* We will use the Pythagorean Theorem to prove the reciprocal identities.



If the point  $P(x, y)$  is a point on the circle with radius  $r$ , then the formula for the circle is

$$x^2 + y^2 = r^2$$

By definition  $\frac{x}{r} = \cos \theta$  and  $\frac{y}{r} = \sin \theta$ . Thus we have

$$\sin^2 \theta + \cos^2 \theta = \left(\frac{y}{r}\right)^2 + \left(\frac{x}{r}\right)^2 = \frac{x^2 + y^2}{r^2} = \frac{r^2}{r^2} = 1$$

which is our first Pythagorean Identity. The proofs of the remaining identities are left as exercises. ■

**Example 1.5.21** Let  $\theta$  be an angle in Quadrant IV and let  $\cos \theta = \frac{3}{5}$ . Calculate the exact values of  $\sin \theta$  and  $\tan \theta$ .

**Solution.** Substituting our value of  $\cos \theta$  into the Pythagorean Identity,

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\begin{aligned}\sin^2 \theta + \left(\frac{3}{5}\right)^2 &= 1 \\ \sin^2 \theta + \frac{9}{25} &= 1 \\ \sin^2 \theta &= 1 - \frac{9}{25} \\ \sin^2 \theta &= \frac{16}{25}\end{aligned}$$

Taking the square root of both sides,

$$\sin \theta = \pm \sqrt{\frac{16}{25}} = \pm \frac{4}{5}$$

Since  $\theta$  is in Quadrant II, we have  $\sin \theta < 0$ . Thus we choose the negative answer to get

$$\sin \theta = -\frac{4}{5}$$

Next we use the Quotient Identity to get

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{-\frac{4}{5}}{\frac{3}{5}} = -\frac{4}{5} \cdot \frac{5}{3} = -\frac{4}{3}$$

□

### 1.5.7 Even and Odd Trigonometric Functions

Recall that a function  $f$  is even if  $f(-x) = f(x)$  for all values of  $x$ , and a function is odd if  $f(-x) = -f(x)$  for all values of  $x$ . With this understanding, we can now classify trigonometric functions as either even or odd.

**Definition 1.5.22 Even and Odd Trigonometric Properties.** The cosine and secant functions are **even**


$$\cos(-\theta) = \cos \theta \qquad \sec(-\theta) = \sec \theta$$

The sine, cosecant, tangent, and cotangent functions are **odd**

$$\begin{aligned}\sin(-\theta) &= -\sin \theta & \csc(-\theta) &= -\csc(\theta) \\ \tan(-\theta) &= -\tan \theta & \cot(-\theta) &= -\cot(\theta)\end{aligned}$$

◇

*Proof.* Let  $P$  be a point on the unit circle corresponding to the angle  $\theta$  with coordinates  $(x, y)$  and  $Q$  be the point corresponding to the angle  $-\theta$  with coordinates  $(x, -y)$ .



Using the [Definition 1.3.4](#) for the six trigonometric functions we have

$$\sin \theta = y, \quad \sin(-\theta) = -y, \quad \cos \theta = x, \quad \cos(-\theta) = x$$

So

$$\sin(-\theta) = -y = -\sin \theta, \quad \cos(-\theta) = x = \cos \theta$$

Thus we conclude that sine is an odd function and cosine is an even function. Next, using the Quotient ([Definition 1.4.3](#)) and Reciprocal Identities ([Definition 1.4.2](#)) we get

$$\begin{aligned} \tan(-\theta) &= \frac{\sin(-\theta)}{\cos(-\theta)} = \frac{-\sin \theta}{\cos \theta} = -\tan \theta, \\ \cot(-\theta) &= \frac{1}{\tan(-\theta)} = \frac{1}{-\tan \theta} = -\cot \theta, \\ \csc(-\theta) &= \frac{1}{\sin(-\theta)} = \frac{1}{-\sin \theta} = -\csc \theta, \\ \sec(-\theta) &= \frac{1}{\cos(-\theta)} = \frac{1}{-\cos \theta} = \sec \theta. \end{aligned}$$

Thus tangent, cotangent, cosecant are odd functions and secant is an even function. ■

**Example 1.5.23** Use the even-odd properties of trigonometric functions to determine the exact value of

(a)  $\csc(-30^\circ)$

**Solution.** Since cosecant is an odd function, the cosecant of a negative angle is the opposite sign of the cosecant of the positive angle. Thus,  $\csc(-30^\circ) = -\csc 30^\circ = -2$

(b)  $\cos(-\theta)$  if  $\cos \theta = 0.4$

**Solution.** Cosine is an even function so  $\cos(-\theta) = \cos \theta = 0.4$ .

□


## 1.5.8 Exercises

**Exercise Group.** Determine the quadrant containing  $\theta$  given the following

- |  |  |
|--|--|
| 1. $\cot \theta < 0$ and $\cos \theta < 0$ | 2. $\csc \theta > 0$ and $\tan \theta > 0$ |
| <b>Answer.</b> QII                         | <b>Answer.</b> QI                          |
| 3. $\cos \theta > 0$ and $\sin \theta < 0$ | 4. $\sec \theta > 0$ and $\tan \theta > 0$ |
| <b>Answer.</b> QIV                         | <b>Answer.</b> QI                          |
| 5. $\tan \theta < 0$ and $\csc \theta > 0$ | 6. $\cot \theta > 0$ and $\sin \theta < 0$ |
| <b>Answer.</b> QII                         | <b>Answer.</b> QIV                         |
| 7. $\sec \theta < 0$ and $\csc \theta < 0$ | 8. $\cos \theta < 0$ and $\tan \theta > 0$ |
| <b>Answer.</b> QIII                        | <b>Answer.</b> QIII                        |

**Exercise Group.** The point  $P(x, y)$  is on the terminal side of angle  $\theta$ . Determine the exact values of the six trigonometric functions at  $\theta$

9.



**Answer.**  $\sin \theta = \frac{3}{5}$ ,  
 $\cos \theta = -\frac{4}{5}$ ,  $\tan \theta = -\frac{3}{4}$ ,  
 $\csc \theta = \frac{5}{3}$ ,  $\sec \theta = -\frac{5}{4}$ ,  
 $\cot \theta = -\frac{4}{3}$

11.  $(3, -4)$ 

**Answer.**  $\sin \theta = -\frac{4}{5}$ ,  
 $\cos \theta = \frac{3}{5}$ ,  $\tan \theta = -\frac{4}{3}$ ,  
 $\csc \theta = -\frac{5}{4}$ ,  $\sec \theta = \frac{5}{3}$ ,  
 $\cot \theta = -\frac{3}{4}$


13.  $(-2, -3)$ 

**Answer.**  $\sin \theta = -\frac{3\sqrt{13}}{13}$ ,  
 $\cos \theta = -\frac{2\sqrt{13}}{13}$ ,  $\tan \theta = \frac{3}{2}$ ,  
 $\csc \theta = -\frac{\sqrt{13}}{3}$ ,  $\sec \theta = -\frac{\sqrt{13}}{2}$ ,  
 $\cot \theta = \frac{2}{3}$

15.  $(-24, 7)$ 

**Answer.**  $\sin \theta = \frac{7}{25}$ ,  
 $\cos \theta = -\frac{24}{25}$ ,  $\tan \theta = -\frac{7}{24}$ ,  
 $\csc \theta = \frac{25}{7}$ ,  $\sec \theta = -\frac{25}{24}$ ,  
 $\cot \theta = -\frac{24}{7}$

10.



**Answer.**  $\sin \theta = -\frac{7\sqrt{85}}{85}$ ,  
 $\cos \theta = \frac{6\sqrt{85}}{85}$ ,  $\tan \theta = -\frac{7}{6}$ ,  
 $\csc \theta = -\frac{\sqrt{85}}{7}$ ,  $\sec \theta = \frac{\sqrt{85}}{6}$ ,  
 $\cot \theta = -\frac{6}{7}$

12.  $(-12, -5)$ 

**Answer.**  $\sin \theta = -\frac{5}{13}$ ,  
 $\cos \theta = -\frac{12}{13}$ ,  $\tan \theta = \frac{5}{12}$ ,  
 $\csc \theta = -\frac{13}{5}$ ,  $\sec \theta = -\frac{13}{12}$ ,  
 $\cot \theta = \frac{12}{5}$

14.  $(-4, 4)$ 

**Answer.**  $\sin \theta = \frac{\sqrt{2}}{2}$ ,  
 $\cos \theta = -\frac{\sqrt{2}}{2}$ ,  $\tan \theta = -1$   
 $\csc \theta = \sqrt{2}$ ,  $\sec \theta = -\sqrt{2}$ ,  
 $\cot \theta = -1$

16.  $(9, -40)$ 

**Answer.**  $\sin \theta = -\frac{40}{41}$ ,  
 $\cos \theta = \frac{9}{41}$ ,  $\tan \theta = -\frac{40}{9}$ ,  
 $\csc \theta = -\frac{41}{40}$ ,  $\sec \theta = \frac{41}{9}$ ,  
 $\cot \theta = -\frac{9}{40}$

**Exercise Group.** Find the exact value of the remaining five trigonometric functions of  $\theta$  from the given information.

17.  $\tan \theta = -\frac{12}{5}$ ,  $\theta$  is Quadrant II  
**Answer.**  $\sin \theta = \frac{12}{13}$ ,  
 $\cos \theta = -\frac{5}{13}$ ,  $\csc \theta = \frac{13}{12}$ ,  
 $\sec \theta = -\frac{13}{5}$ ,  $\cot \theta = -\frac{5}{12}$
18.  $\cos \theta = \frac{3}{5}$ ,  $\theta$  is Quadrant IV  
**Answer.**  $\sin \theta = -\frac{4}{5}$ ,  
 $\tan \theta = -\frac{4}{3}$ ,  $\csc \theta = -\frac{5}{4}$ ,  
 $\sec \theta = \frac{5}{3}$ ,  $\cot \theta = -\frac{3}{4}$
19.  $\csc \theta = \frac{\sqrt{10}}{2}$ ,  $\theta$  is Quadrant II  
**Answer.**  $\sin \theta = \frac{\sqrt{10}}{5}$ ,  
 $\cos \theta = -\frac{\sqrt{15}}{5}$ ,  $\tan \theta = -\frac{\sqrt{6}}{3}$ ,  
 $\sec \theta = -\frac{\sqrt{15}}{3}$ ,  $\cot \theta = -\frac{6}{2}$
20.  $\cos \theta = -\frac{5}{8}$ ,  $\theta$  is Quadrant III  
**Answer.**  $\sin \theta = -\frac{\sqrt{39}}{8}$ ,  
 $\tan \theta = \frac{\sqrt{39}}{5}$ ,  $\csc \theta = -\frac{8\sqrt{39}}{39}$ ,  
 $\sec \theta = -\frac{8}{5}$ ,  $\cot \theta = \frac{5\sqrt{39}}{39}$
21.  $\sec \theta = -2$ ,  $\pi < \theta < \frac{3\pi}{2}$   
**Answer.**  $\sin \theta = -\frac{\sqrt{3}}{2}$ ,  
 $\cos \theta = -\frac{1}{2}$ ,  $\tan \theta = \sqrt{3}$ ,  
 $\csc \theta = -\frac{2\sqrt{3}}{3}$ ,  $\cot \theta = \frac{\sqrt{3}}{3}$
22.  $\cot \theta = -\frac{5}{3}$ ,  $\frac{3\pi}{2} < \theta < 2\pi$   
**Answer.**  $\sin \theta = -\frac{3\sqrt{34}}{34}$ ,  
 $\cos \theta = \frac{5\sqrt{34}}{34}$ ,  $\tan \theta = -\frac{3}{5}$ ,  
 $\csc \theta = -\frac{\sqrt{34}}{3}$ ,  $\sec \theta = \frac{\sqrt{34}}{5}$
23.  $\cos \theta = \frac{2}{3}$ ,  $0 < \theta < \pi$   
**Answer.**  $\sin \theta = \frac{\sqrt{5}}{3}$ ,  
 $\tan \theta = -\frac{\sqrt{5}}{2}$ ,  $\csc \theta = \frac{3\sqrt{5}}{5}$ ,  
 $\sec \theta = -\frac{3}{2}$ ,  $\cot \theta = -\frac{2\sqrt{5}}{5}$
24.  $\tan \theta = \frac{7}{4}$ ,  $0 < \theta < \frac{\pi}{2}$   
**Answer.**  $\sin \theta = \frac{7\sqrt{65}}{65}$ ,  
 $\cos \theta = \frac{4\sqrt{65}}{65}$ ,  $\csc \theta = \frac{\sqrt{65}}{7}$ ,  
 $\sec \theta = \frac{\sqrt{65}}{4}$ ,  $\cot \theta = \frac{4}{7}$
25.  $\csc \theta = \frac{3}{2}$ ,  $\tan \theta < 0$   
**Answer.**  $\sin \theta = \frac{2}{3}$ ,  
 $\cos \theta = -\frac{\sqrt{5}}{3}$ ,  $\tan \theta = -\frac{2\sqrt{5}}{5}$ ,  
 $\sec \theta = -\frac{3\sqrt{5}}{5}$ ,  $\cot \theta = -\frac{\sqrt{5}}{2}$
26.  $\sin \theta = \frac{5}{6}$ ,  $\cot \theta > 0$   
**Answer.**  $\cos \theta = \frac{\sqrt{11}}{6}$ ,  
 $\tan \theta = \frac{5\sqrt{11}}{11}$ ,  $\csc \theta = \frac{6}{5}$ ,  
 $\sec \theta = \frac{6\sqrt{11}}{11}$ ,  $\cot \theta = \frac{\sqrt{11}}{5}$
27.  $\sin \theta = -\frac{15}{17}$ ,  $\cos \theta < 0$   
**Answer.**  $\cos \theta = -\frac{8}{17}$ ,  
 $\tan \theta = \frac{15}{8}$ ,  $\csc \theta = -\frac{17}{15}$ ,  
 $\sec \theta = -\frac{17}{8}$ ,  $\cot \theta = \frac{8}{15}$
28.  $\cot \theta = -\frac{1}{3}$ ,  $\sin \theta > 0$   
**Answer.**  $\sin \theta = \frac{3\sqrt{10}}{10}$ ,  
 $\cos \theta = -\frac{\sqrt{10}}{10}$ ,  $\tan \theta = -3$ ,  
 $\csc \theta = \frac{\sqrt{10}}{3}$ ,  $\sec \theta = -\sqrt{10}$

**Exercise Group.** Given a reference angle,  $t'$ , calculate the corresponding angle,  $t$ , in standard position, along with the values of  $\sin t$ ,  $\cos t$ , and  $\tan t$  for

(a) Quadrant II

(b) Quadrant III

(c) Quadrant IV

29.  $t' = \frac{\pi}{4}$

**Answer 1.**  $t = \frac{3\pi}{4}$ ,  
 $\sin \frac{3\pi}{4} = \frac{\sqrt{2}}{2}$ ,  $\cos \frac{3\pi}{4} = -\frac{\sqrt{2}}{2}$ ,  
 $\tan \frac{3\pi}{4} = -1$

**Answer 2.**  $t = \frac{5\pi}{4}$ ,  
 $\sin \frac{5\pi}{4} = -\frac{\sqrt{2}}{2}$ ,  $\cos \frac{5\pi}{4} = -\frac{\sqrt{2}}{2}$ ,  
 $\tan \frac{5\pi}{4} = 1$

**Answer 3.**  $t = \frac{7\pi}{4}$ ,  
 $\sin \frac{7\pi}{4} = -\frac{\sqrt{2}}{2}$ ,  $\cos \frac{7\pi}{4} = \frac{\sqrt{2}}{2}$ ,  
 $\tan \frac{7\pi}{4} = -1$

30.  $t' = \frac{\pi}{3}$

**Answer 1.**  $t = \frac{2\pi}{3}$ ,  
 $\sin \frac{2\pi}{3} = \frac{\sqrt{3}}{2}$ ,  $\cos \frac{2\pi}{3} = -\frac{1}{2}$ ,  
 $\tan \frac{2\pi}{3} = -\sqrt{3}$

**Answer 2.**  $t = \frac{4\pi}{3}$ ,  
 $\sin \frac{4\pi}{3} = -\frac{\sqrt{3}}{2}$ ,  $\cos \frac{4\pi}{3} = -\frac{1}{2}$ ,  
 $\tan \frac{4\pi}{3} = \sqrt{3}$

**Answer 3.**  $t = \frac{5\pi}{3}$ ,  
 $\sin \frac{5\pi}{3} = -\frac{\sqrt{3}}{2}$ ,  $\cos \frac{5\pi}{3} = \frac{1}{2}$ ,  
 $\tan \frac{5\pi}{3} = -\sqrt{3}$



31.  $t' = 30^\circ$

**Answer 1.**  $t = 150^\circ$ ,  
 $\sin 150^\circ = \frac{1}{2}$ ,  $\cos 150^\circ = -\frac{\sqrt{3}}{2}$ ,  
 $\tan 150^\circ = -\frac{\sqrt{3}}{3}$

**Answer 2.**  $t = 210^\circ$ ,  
 $\sin 210^\circ = -\frac{1}{2}$ ,  
 $\cos 210^\circ = -\frac{\sqrt{3}}{2}$ ,  
 $\tan 210^\circ = \frac{\sqrt{3}}{3}$

**Answer 3.**  $t = 330^\circ$ ,  
 $\sin 330^\circ = -\frac{1}{2}$ ,  $\cos 330^\circ = \frac{\sqrt{3}}{2}$ ,  
 $\tan 330^\circ = -\frac{\sqrt{3}}{3}$

32.  $t' = 60^\circ$

**Answer 1.**  $t = 120^\circ$ ,  
 $\sin 120^\circ = \frac{\sqrt{3}}{2}$ ,  $\cos 120^\circ = -\frac{1}{2}$ ,  
 $\tan 120^\circ = -\sqrt{3}$

**Answer 2.**  $t = 240^\circ$ ,  
 $\sin 240^\circ = -\frac{\sqrt{3}}{2}$ ,  
 $\cos 240^\circ = -\frac{1}{2}$ ,  $\tan 240^\circ = \sqrt{3}$

**Answer 3.**  $t = 300^\circ$ ,  
 $\sin 300^\circ = -\frac{\sqrt{3}}{2}$ ,  $\cos 300^\circ = \frac{1}{2}$ ,  
 $\tan 300^\circ = -\sqrt{3}$

**Exercise Group.** For each angle  $\theta$ ,

- (a) Determine the quadrant in which  $\theta$  lies.  
 (b) Calculate the reference angle  $\theta'$   
 (c) Use the reference angle,  $\theta'$  to evaluate the exact values of the six trigonometric functions for  $\theta$

33.  $\theta = -\frac{3\pi}{4}$

**Answer**  
 1. QIII

**Answer**  
 2.  $\theta' = \frac{\pi}{4}$

**Answer**  
 3.  $\sin \theta = -\frac{\sqrt{2}}{2}$ ,  
 $\cos \theta = -\frac{\sqrt{2}}{2}$ ,  
 $\tan \theta = 1$ ,  
 $\csc \theta = -\sqrt{2}$ ,  
 $\sec \theta = -\sqrt{2}$ ,  
 $\cot \theta = 1$

34.  $\theta = \frac{4\pi}{3}$

**Answer**  
 1. QIII

**Answer**  
 2.  $\theta' = \frac{\pi}{3}$

**Answer**  
 3.  $\sin \theta = -\frac{\sqrt{3}}{2}$ ,  
 $\cos \theta = -\frac{1}{2}$ ,  
 $\tan \theta = \sqrt{3}$ ,  
 $\csc \theta = -\frac{2\sqrt{3}}{3}$ ,  
 $\sec \theta = -2$ ,  
 $\cot \theta = \frac{\sqrt{3}}{3}$

35.  $\theta = \frac{11\pi}{6}$

**Answer**  
 1. QIV

**Answer**  
 2.  $\theta' = \frac{\pi}{6}$

**Answer**  
 3.  $\sin \theta = -\frac{1}{2}$ ,  
 $\cos \theta = \frac{\sqrt{3}}{2}$ ,  
 $\tan \theta = -\frac{\sqrt{3}}{3}$ ,  
 $\csc \theta = -2$ ,  
 $\sec \theta = \frac{2\sqrt{3}}{3}$ ,  
 $\cot \theta = -\sqrt{3}$

36.  $\theta = \frac{7\pi}{3}$

**Answer 1.** QI

**Answer**  
 2.  $\theta' = \frac{\pi}{3}$

**Answer**  
 3.  $\sin \theta = \frac{\sqrt{3}}{2}$ ,  
 $\cos \theta = \frac{1}{2}$ ,  
 $\tan \theta = \sqrt{3}$ ,  
 $\csc \theta = \frac{2\sqrt{3}}{3}$ ,  
 $\sec \theta = 2$ ,  
 $\cot \theta = \frac{\sqrt{3}}{3}$

37.  $\theta = 120^\circ$

**Answer 1.** QII

**Answer**  
 2.  $\theta' = 60^\circ$

**Answer**  
 3.  $\sin \theta = \frac{\sqrt{3}}{2}$ ,  
 $\cos \theta = -\frac{1}{2}$ ,  
 $\tan \theta = -\sqrt{3}$ ,  
 $\csc \theta = \frac{2\sqrt{3}}{3}$ ,  
 $\sec \theta = -2$ ,  
 $\cot \theta = -\frac{\sqrt{3}}{3}$

38.  $\theta = 480^\circ$

**Answer 1.** QII

**Answer**  
 2.  $\theta' = 60^\circ$

**Answer**  
 3.  $\sin \theta = \frac{\sqrt{3}}{2}$ ,  
 $\cos \theta = -\frac{1}{2}$ ,  
 $\tan \theta = -\sqrt{3}$ ,  
 $\csc \theta = \frac{2\sqrt{3}}{3}$ ,  
 $\sec \theta = -2$ ,  
 $\cot \theta = -\frac{\sqrt{3}}{3}$

**Exercise Group.** Use the fact that the trigonometric functions are periodic to find the exact value for each expression.

39.  $\tan 420^\circ$

**Answer.**  $\sqrt{3}$

40.  $\csc 540^\circ$

**Answer.** Undefined

41.  $\sin 765^\circ$

**Answer.**  $\frac{\sqrt{2}}{2}$

42.  $\sec 1200^\circ$

**Answer.**  $-2$

$$43. \cot \frac{8\pi}{3} \quad 44. \cos \frac{21\pi}{4} \quad 45. \tan \frac{35\pi}{6} \quad 46. \sin \frac{39\pi}{4}$$

$$\text{Answer. } -\frac{\sqrt{3}}{3} \quad \text{Answer. } -\frac{\sqrt{2}}{2} \quad \text{Answer. } \frac{\sqrt{3}}{2} \quad \text{Answer. } -\frac{\sqrt{2}}{2}$$

47. Prove the second Pythagorean Identity (Definition 1.5.20):  $1 + \tan^2 \theta = \sec^2 \theta$ .

**Hint.** Begin with  $\sin^2 \theta + \cos^2 \theta = 1$  and divide both sides of the equation by  $\cos^2 \theta$ .

48. Prove the third Pythagorean Identity (Definition 1.5.20):  $1 + \cot^2 \theta + 1 = \csc^2 \theta$ .

**Hint.** Begin with  $\sin^2 \theta + \cos^2 \theta = 1$  and divide both sides of the equation by  $\sin^2 \theta$ .

**Exercise Group.** Use the Pythagorean Identity to find the exact value of the following

$$49. \sin^2 38^\circ + \cos^2 38^\circ \quad 50. \csc^2 13^\circ - \cot^2 13^\circ \quad 51. \cot^2 200^\circ - \csc^2 200^\circ$$

$$\text{Answer. } 1 \quad \text{Answer. } 1 \quad \text{Answer. } -1$$

$$52. \sec^2 \frac{6\pi}{5} - \tan^2 \frac{6\pi}{5} \quad 53. \tan^2 \frac{5\pi}{7} - \sec^2 \frac{5\pi}{7} \quad 54. \sin^2 \frac{14\pi}{13} + \cos^2 \frac{14\pi}{13}$$

$$\text{Answer. } 1 \quad \text{Answer. } -1 \quad \text{Answer. } 1$$

**Exercise Group.** Use the Pythagorean Identities to express the first trigonometric function of  $\theta$  in terms of the second function, given the quadrant.

$$55. \sin \theta, \cos \theta, \text{ Quadrant III} \quad 56. \cos \theta, \sin \theta, \text{ Quadrant II}$$

$$\text{Answer. } \sin \theta = -\sqrt{1 - \cos^2 \theta} \quad \text{Answer. } \cos \theta = -\sqrt{1 - \sin^2 \theta}$$

$$57. \tan, \sec \theta, \text{ Quadrant IV} \quad 58. \cot \theta, \csc \theta, \text{ Quadrant III}$$

$$\text{Answer. } \tan \theta = -\sqrt{\sec^2 \theta - 1} \quad \text{Answer. } \cot \theta = \sqrt{\csc^2 \theta - 1}$$

$$59. \tan \theta, \sin \theta, \text{ Quadrant III} \quad 60. \tan \theta, \cos \theta, \text{ Quadrant II}$$

$$\text{Answer. } \tan \theta = -\frac{\sin \theta}{\sqrt{1 - \sin^2 \theta}} \quad \text{Answer. } \tan \theta = \frac{\sqrt{1 - \cos^2 \theta}}{\cos \theta}$$

**Exercise Group.** Use the Pythagorean Identities to find the exact values of the remaining five trigonometric functions of  $\theta$  from the given information.

$$61. \tan \theta = -\frac{4}{3}, \theta \text{ is in Quadrant IV} \quad 62. \cos \theta = -\frac{1}{4}, \theta \text{ is in Quadrant II}$$

$$\text{Answer. } \sin \theta = -\frac{4}{5}, \cos \theta = \frac{3}{5}, \csc \theta = -\frac{5}{4}, \sec \theta = \frac{5}{3}, \cot \theta = -\frac{3}{4} \quad \text{Answer. } \sin \theta = \frac{\sqrt{15}}{4}, \tan \theta = -\sqrt{15}, \csc \theta = \frac{4\sqrt{15}}{15}, \sec \theta = -4, \cot \theta = -\frac{\sqrt{15}}{15}$$

$$63. \sin \theta = -\frac{2}{3}, \theta \text{ is in Quadrant III} \quad 64. \cos \theta = \frac{3}{5}, \theta \text{ is in Quadrant IV}$$

$$\text{Answer. } \cos \theta = -\frac{\sqrt{5}}{3}, \tan \theta = \frac{2\sqrt{5}}{5}, \csc \theta = -\frac{3}{2}, \sec \theta = -\frac{3\sqrt{5}}{5}, \cot \theta = \frac{\sqrt{5}}{2} \quad \text{Answer. } \sin \theta = -\frac{4}{5}, \tan \theta = -\frac{4}{3}, \csc \theta = -\frac{5}{4}, \sec \theta = \frac{5}{3}, \cot \theta = -\frac{3}{4}$$

**Exercise Group.** Use the even and odd properties to evaluate the following

$$65. \cos(-60^\circ) \quad 66. \tan(-225^\circ) \quad 67. \csc(-330^\circ)$$

$$\text{Answer. } \frac{1}{2} \quad \text{Answer. } -1 \quad \text{Answer. } 2$$

68.  $\sin(-90^\circ)$

**Answer.**  $-1$

69.  $\cot(-300^\circ)$

**Answer.**  $\frac{\sqrt{3}}{3}$

70.  $\sec(-150^\circ)$

**Answer.**  $-\frac{2\sqrt{3}}{3}$

71.  $\sin\left(-\frac{11\pi}{6}\right)$

**Answer.**  $\frac{1}{2}$

72.  $\tan\left(-\frac{5\pi}{4}\right)$

**Answer.**  $-1$

73.  $\cos\left(-\frac{4\pi}{3}\right)$

**Answer.**  $-\frac{1}{2}$

74.  $\tan(-\pi)$

**Answer.**  $0$


75.  $\sec\left(-\frac{\pi}{4}\right)$

**Answer.**  $\sqrt{2}$

76.  $\csc\left(-\frac{7\pi}{6}\right)$

**Answer.**  $2$

**Exercise Group.** The Makali'i is sailing along the Kohala Coast, maintaining a distance of two nautical miles from the shore. An observer at Mahukona is monitoring Makali'i's passage. Let  $d$  denote the length of the line connecting Makali'i to the Mahukona observer. Given  $\theta$  as the angle formed between  $d$  and the shore, determine Makali'i's distance,  $d$ , from the observer for each value of  $\theta$ , rounded to one decimal place.



77.  $\theta = 30^\circ$

**Answer.** 4  
NM

78.  $\theta = 45^\circ$

**Answer.** 2.8  
NM

79.  $\theta = 60^\circ$

**Answer.** 2.3  
NM

80.  $\theta = 90^\circ$

**Answer.** 2  
NM

81.  $\theta = 120^\circ$

**Answer.** 2.3  
NM

82.  $\theta = 135^\circ$

**Answer.** 2.8  
NM

83.  $\theta = 150^\circ$

**Answer.** 4  
NM

## Chapter 2

# Graphs of the Trigonometric Functions

### 2.1 Graphs of the Sine and Cosine Functions

Sunrise and sunset are vital markers for navigational orientation and course corrections. During sunrise, you can determine the direction and observe the origin of wind and waves. As the sun rises higher, steering by the sun becomes impractical, and you must instead depend on swells to keep your course. At sunset, you can reassess your position and take note of any changes in wind and swell patterns. Although the general pattern is for the sun to rise in the east and set in the west, its exact position on the horizon, known as **solar declination**, varies throughout the year.

To understand why the position changes, we must first learn about Earth's **axial tilt**, which represents the angle between the Earth's rotational axis and its orbital plane around the Sun. To conceptualize this tilt, envision a pole passing through the Earth's center, extending from the North to South Pole, with the Earth revolving around this axis. Each complete rotation of the Earth on this axis corresponds to one day. As the Earth travels along its orbit around the Sun with a constant tilt, the orientation of the Earth's axial tilt causes either the North Pole or the South Pole to tilt toward the Sun. This tilt varies depending on the Earth's location in its orbit relative to the Sun.

During the equinoxes, which mark the transition from winter to spring and from summer to fall, the Earth's axis is not tilted towards or away from the Sun. Consequently, on these days, the Sun rises due east and sets due west, and the duration of day and night is approximately equal.


Following the **fall equinox** in the Southern Hemisphere, typically occurring around March 20th, the Earth proceeds along its orbit around the Sun, leading the Southern Hemisphere to tilt away from the Sun. As a result, the Sun's rising position progressively moves northward each day. By the time of the **winter solstice**, around June 20th, the Sun rises from its northernmost position, resulting in the shortest day in the Southern Hemisphere and the longest day of the year in the Northern Hemisphere.

After the winter solstice, the Earth's axial tilt remains the same, but the Southern Hemisphere starts tilting toward the Sun. This causes the Sun's rising position to gradually shift southward each day. When the **spring equinox** arrives, approximately on September 22nd, the Sun rises due east once again, and day and night are once more of equal duration.

As the Southern Hemisphere continues to tilt towards the Sun, the Sun's

rising position moves even further south each day. By the time of the **summer solstice**, approximately on December 21st, the Sun rises from its southernmost position, resulting in the longest day of the year in the Southern Hemisphere and the shortest day in the Northern Hemisphere.

Following the summer solstice, the Southern Hemisphere begins tilting away from the Sun, causing the Sun's rising position to gradually shift northward each day. This movement continues until the next fall equinox, completing the annual cycle.



**Figure 2.1.1** Illustration shows the relative positions and timing of solstice, equinox and seasons in relation to the Earth's orbit around the Sun and the axial tilt. During the June solstice, the northern hemisphere is tilted towards the Sun, while the southern hemisphere is tilted away. Conversely, during the December solstice, the southern hemisphere is tilted towards the Sun, and the northern hemisphere is tilted away. During the March and September equinoxes, the Earth's axis is perpendicular to the line connecting the Earth to the Sun, causing the Sun to appear directly above the equator. As a result, day and night are approximately equal in length all over the world.

The position where the Sun rises throughout the year exhibits a repeating pattern, characterized as a **periodic function** with a cycle of about one year. This function can be mathematically represented as either a sine or cosine function. Graphically, it illustrates the sunrise position on the horizon relative to the east (Hikina) as time progresses. By examining this function, we can observe the gradual northward and southward movement of the sunrise position throughout the year. Key dates such as the equinoxes and solstices mark significant points in this pattern, providing insights into the changing seasons and variations in daylight hours.

In this section, we will explore the periodic nature of sine and cosine functions and study their transformations. Studying the graphs of sine and cosine functions provides valuable insights into the world around us. The graphs of the remaining trigonometric functions will be covered in [Section 2.2](#).

### 2.1.1 Domain and Range of Sine and Cosine

The domains for  $\sin(\theta)$  and  $\cos(\theta)$  consist of the set of the inputs of the functions. Since any angle  $\theta$  can be input into sine and cosine and still have these functions defined, the domain for both sine and cosine is all real numbers. Recall that in [Section 1.3](#), if  $P(x, y)$  is any point on the unit circle that corresponds to the angle  $\theta$ , we defined  $\sin \theta = y$  and  $\cos \theta = x$ . Given the constraints on the unit circle,  $-1 \leq x \leq 1$  and  $-1 \leq y \leq 1$ , and thus

$$-1 \leq \sin \theta \leq 1, \quad \text{and} \quad -1 \leq \cos \theta \leq 1.$$

Since the range of a function consists of all its outputs, we conclude that the range of both the sine and cosine functions spans all real numbers between -1 and 1.

**Remark 2.1.2 Domain and Range for the Sine and Cosine Functions.** [Table 2.1.3](#) summarizes the domains and ranges for the sine and cosine functions.

**Table 2.1.3 Domains and Ranges of Sine and Cosine**


Function	Domain	Range
$\sin \theta$	All real numbers, $(-\infty, \infty)$	All real numbers from -1 to 1, $[-1, 1]$
$\cos \theta$	All real numbers, $(-\infty, \infty)$	All real numbers from -1 to 1, $[-1, 1]$

### 2.1.2 The Sine Function

**Convention 2.1.4** In [Chapter 1](#), trigonometric functions typically use  $\theta$  or  $t$  as the variable in the domain, such as  $y = \cos \theta$  and  $y = \sin t$ . However, when graphing functions on the Cartesian plane ( $xy$ -coordinate system),  $x$  is conventionally used as the variable in the domain. Therefore, when graphing trigonometric functions, we will use  $x$  as the variable, for example  $y = \cos x$  and  $y = \sin x$ .

The sine function, as discussed in [Subsection 1.5.4](#), is a periodic functions with period  $2\pi$ . To graph  $y = \sin x$ , we can focus on the interval  $[0, 2\pi]$ . By plotting this interval, we can then repeat the values over the entire domain to complete the remaining graph.

Recall from [Definition 1.3.2](#) that on the unit circle,  $\sin \theta$  is defined to be the  $y$ -value of the terminal point  $P(x, y)$  on the unit circle associated with the angle  $\theta$ . As the angle increases from 0 to  $\frac{\pi}{2}$ , the  $y$ -value also increases from 0 to 1. When the angle continues from  $\frac{\pi}{2}$  to  $\frac{3\pi}{2}$ , the  $y$ -value decreases from 1 to  $-1$ . Finally, as the angle approaches  $2\pi$ , the  $y$ -value increases to 0. This behavior is shown in [Figure 2.1.5](#).




**Figure 2.1.5** As  $\theta$  moves from  $0^\circ$  to  $360^\circ$ , this figure plots the values of  $y = \sin \theta$ . Move the slider for  $\theta$  to see how changing the angle affects  $\sin \theta$ . Note that while we will generally be using radians when graphing trigonometric functions, this figure uses degrees to help visualize the angle. For example, you may not be aware that 2 radians ( $\approx 114.6^\circ$ ) lies in Quadrant II.

Next we recall known values for the sine function listed on [Table 2.1.6](#).

**Table 2.1.6** Values for  $y = \sin x$


$x$	$\sin x$	$(x, y)$	$x$	$\sin x$	$(x, y)$
0	0	$(0, 0)$	$\pi$	0	$(\pi, 0)$
$\frac{\pi}{6}$	$\frac{1}{2}$	$(\frac{\pi}{6}, \frac{1}{2})$	$\frac{7\pi}{6}$	$-\frac{1}{2}$	$(\frac{7\pi}{6}, -\frac{1}{2})$
$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$(\frac{\pi}{4}, \frac{\sqrt{2}}{2})$	$\frac{5\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$(\frac{5\pi}{4}, -\frac{\sqrt{2}}{2})$
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$(\frac{\pi}{3}, \frac{\sqrt{3}}{2})$	$\frac{4\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$(\frac{4\pi}{3}, -\frac{\sqrt{3}}{2})$
$\frac{\pi}{2}$	1	$(\frac{\pi}{2}, 1)$	$\frac{3\pi}{2}$	-1	$(\frac{3\pi}{2}, -1)$
$\frac{2\pi}{3}$	$\frac{\sqrt{3}}{2}$	$(\frac{2\pi}{3}, \frac{\sqrt{3}}{2})$	$\frac{5\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$(\frac{5\pi}{3}, -\frac{\sqrt{3}}{2})$
$\frac{3\pi}{4}$	$\frac{\sqrt{2}}{2}$	$(\frac{3\pi}{4}, \frac{\sqrt{2}}{2})$	$\frac{7\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$(\frac{7\pi}{4}, -\frac{\sqrt{2}}{2})$
$\frac{5\pi}{6}$	$\frac{1}{2}$	$(\frac{5\pi}{6}, \frac{1}{2})$	$\frac{11\pi}{6}$	$-\frac{1}{2}$	$(\frac{11\pi}{6}, -\frac{1}{2})$
			$2\pi$	0	$(2\pi, 0)$

Now that we have a visual understanding of the graph for  $y = \sin x$ , we can utilize the data from [Table 2.1.6](#) to map out the points. This process enables us to construct the graph illustrated in [Figure 2.1.7](#), representing one complete period of the sine function.



**Figure 2.1.7** The values in [Table 2.1.6](#) plotted on a graph with a smooth curve connecting the points to make the curve for  $y = \sin(x)$ .

Since the graph in [Figure 2.1.7](#) represents one period, we can now complete the graph of  $y = \sin x$  by extending the pattern in both directions to obtain [Figure 2.1.8](#).




**Figure 2.1.8** The plot for  $y = \sin(x)$ .

Notice the graph of the sine function's symmetry with respect to the origin, a characteristic supported in [Section 1.5.7](#), where we learned that sine is an odd function.




### 2.1.3 The Cosine Function

Similarly, we can construct a plot for the cosine function, shown in [Figure 2.1.9](#).



**Figure 2.1.9** As  $\theta$  moves from  $0^\circ$  to  $360^\circ$ , this figure plots the values of  $y = \cos \theta$ . Move the slider for  $\theta$  to see how changing the angle affects  $\cos \theta$ . Note that while we will generally be using radians when graphing trigonometric functions, this figure uses degrees to help visualize the angle.

By plotting points for  $y = \cos x$  and using the fact that the cosine function is periodic, we obtain the graph for cosine over the entire domain. This is shown in [Figure 2.1.10](#).



**Figure 2.1.10** The plot for  $y = \cos(x)$ .

In alignment with [Section 1.5.7](#), observe that the graph of cosine is symmetric about the  $y$ -axis, confirming that it is an even function.

**Definition 2.1.11** The graphs for the sine and cosine functions are commonly referred to as **sinusoidal graphs** or **sinus curves**.  $\diamond$

### 2.1.4 Graphing Transformations of Sine and Cosine

Now that we've become familiar with the graphs of the sine and cosine functions, let's apply algebraic graphing techniques to these functions. Recall that when  $D > 0$ , the graph of  $y = f(x) + D$  shifts the graph of  $y = f(x)$  upward by  $D$  units, and the graph of  $y = f(x) - D$  shifts the graph of  $y = f(x)$  downward by  $D$  units.

**Definition 2.1.12 Vertical Shift.** The graphs of the functions

$$y = \sin x + D \quad \text{and} \quad y = \cos x + D$$

represent an **upward vertical shift** of the graphs  $y = \sin(x)$  and  $y = \cos(x)$  by  $D$  units, respectively.

Similarly, the functions


$$y = \sin x - D \quad \text{and} \quad y = \cos x - D$$

depict the graphs of  $y = \sin(x)$  and  $y = \cos(x)$  with a **downward vertical shift** by  $D$  units, respectively.  $\diamond$


**Example 2.1.13 Vertical Shifts.** Graph each function

1.  $y = \sin(x) + 2$
2.  $y = \cos(x) - 1$

**Solution.**



**Figure 2.1.14** The graph of  $y = \sin(x) + 2$  is the same as the graph of  $y = \sin(x)$  but shifted up by 2 units.



**Figure 2.1.15** The graph of  $y = \cos(x) - 1$  is the same as the graph of  $y = \cos(x)$  but shifted down by 1 unit.

□

Additionally, remember that the graph  $y = -f(x)$  reflects the graph of  $y = f(x)$  about the  $x$ -axis.


**Definition 2.1.16 Reflection about the  $x$ -axis.** The functions

$$y = -\sin x \quad \text{and} \quad y = -\cos x$$

represent the graphs of  $y = \sin(x)$  and  $y = \cos(x)$  with a **reflection about the  $x$ -axis**, respectively. ◇

**Example 2.1.17 Reflections about the  $x$ -axis.** Graph  $y = -\cos(x)$

**Solution.**



**Figure 2.1.18** The graph  $y = -\cos(x)$  is obtained by multiplying every  $y$ -value of the  $y = \cos(x)$  graph by  $-1$ . This transformation reflects all points across the  $x$ -axis, turning positive values negative and negative values positive.

□

Similarly, recall that the graph of  $y = f(-x)$  reflects the graph of  $y = f(x)$

about the  $y$ -axis.


**Definition 2.1.19 Reflection about the  $y$ -axis.** The functions

$$y = \sin(-x) \quad \text{and} \quad y = \cos(-x)$$

represent the graphs of  $y = \sin(x)$  and  $y = \cos(x)$  with a **reflection about the  $y$ -axis**, respectively.  $\diamond$

**Example 2.1.20 Reflections about the  $y$ -axis.** Graph  $y = \sin(-x) + 1$

**Solution.**




**Figure 2.1.21** The graph  $y = \sin(-x) + 1$  is obtained by reflecting the graph of  $y = \sin(x)$  about the  $y$ -axis and then vertically shifting it upward by 1. This transformation turns positive values of  $x$  negative and negative values positive, and increases every  $y$ -value by 1.  $\square$

**Example 2.1.22 Vertical Stretches and Compressions.** Graph each function

1.  $y = 2 \cos(x)$
2.  $y = \frac{1}{2} \cos(x)$

**Solution.**



**Figure 2.1.23** The graph of  $y = 2 \cos(x)$  is achieved by vertically stretching the  $y$ -values of  $y = \cos(x)$  by a factor of 2. Similarly, the graph of  $y = \frac{1}{2} \cos(x)$  is obtained by vertically compressing the  $y$ -values of  $y = \cos(x)$  by a factor of  $\frac{1}{2}$ .

□

The factor multiplied at the front of the cosine function plays a crucial role in stretching and compressing the graph. This factor is known as the **amplitude**, measuring the maximum vertical distance from the midline to the peak or trough of a sinusoidal wave. In [Example 2.1.22](#), the amplitude of  $y = 2 \cos(x)$  is 2, indicating a **vertical stretch** by a factor of 2 compared to the standard cosine function. Conversely, for  $y = \frac{1}{2} \cos(x)$ , the amplitude is  $\frac{1}{2}$ , which represented a **vertical compression** by a factor of  $\frac{1}{2}$ .

**Definition 2.1.24 Amplitude.** For a sinusoidal function, the **amplitude**, denoted as  $|A|$ , is the height of the function, representing half the distance between its maximum and minimum values:

$$|A| = \text{amplitude} = \frac{\text{maximum} - \text{minimum}}{2}.$$

In other words, the amplitude is the vertical distance from the midline to the maximum or minimum value of the function. The **midline** is a horizontal line representing the average value of the function. It can be calculated by:

$$y = \frac{\text{maximum} + \text{minimum}}{2}.$$

For a graph centered about the  $x$ -axis, the amplitude is simply the maximum value of the function.

In general, for

$$y = A \cdot \sin(x) \quad \text{or} \quad y = A \cdot \cos(x),$$

the amplitude is given by  $|A|$ . This absolute value ensures that amplitude is always a positive value, representing the magnitude of the vertical stretching or compression. ◇

**Definition 2.1.25 Vertical Stretch/Compression.** The functions

$$y = A \sin x \quad \text{and} \quad y = A \cos x$$


represents a sine and cosine function, respectively with an amplitude of  $|A|$ . The amplitude determines the vertical stretch or compression of the graph.

If  $|A| > 1$ , the graph undergoes a vertical stretch, making the peaks and troughs higher.

If  $0 < |A| < 1$ , the graph undergoes a vertical compression, reducing the distance between the peaks and troughs.  $\diamond$

**Example 2.1.26** Graph  $y = -4 \sin x$  and identify the amplitude.

**Solution.** The amplitude is  $|-4| = 4$ .



**Figure 2.1.27** Since the amplitude of  $y = -4 \sin(x)$  is 4, the graph is stretched by a factor of 4 and will oscillate between  $-4$  and  $4$ . Additionally, the negative sign indicates that the graph is reflected about the  $x$ -axis.  $\square$

Next we will look at functions of the form

$$y = \sin Bx \quad \text{and} \quad y = \cos Bx$$

You may recall from algebra that for functions of the form  $y = f(Bx)$ , a key factor emerges: when  $|B| > 1$ , the graph undergoes horizontal compression by a factor of  $\frac{1}{|B|}$ ; conversely, when  $0 < |B| < 1$ , the graph is horizontally stretched by a factor of  $\frac{1}{|B|}$ . Given that sine and cosine complete one period in  $2\pi$ , the horizontal stretching or compressing of a period will be by a factor of  $\frac{1}{|B|}$ .

**Definition 2.1.28 Period.** For sine and cosine functions of the form

$$y = \sin Bx \quad \text{and} \quad y = \cos Bx$$

the **period** is defined as

$$\text{period} = \frac{2\pi}{|B|}$$

Thus, if  $|B| > 1$ , the period is compressed; if  $0 < |B| < 1$ , the period is stretched.  $\diamond$

**Definition 2.1.29 Horizontal Stretch/Compression.** The graphs functions

$$y = \sin Bx \quad \text{and} \quad y = \cos Bx,$$

undergo a horizontal stretch or compression by a value of  $\frac{1}{|B|}$ .

If  $|B| > 1$ , the graph undergoes a horizontal compression, making the period shorter.

If  $0 < |B| < 1$ , the graph undergoes a horizontal stretch, making the period longer.  $\diamond$

**Example 2.1.30 Horizontal Stretches and Compressions.** Identify the period and graph one period for each of the following functions:

1.  $y = \sin(2x)$

2.  $y = \sin\left(\frac{1}{2}x\right)$

3.  $y = \sin\left(\frac{1}{3}x\right)$

**Solution.**

1. The period for  $y = \sin(2x)$  is


$$\text{period} = \frac{2\pi}{2} = \pi$$

2. The period for  $y = \sin\left(\frac{1}{2}x\right)$  is

$$\text{period} = \frac{2\pi}{\frac{1}{2}} = 2\pi \cdot \frac{2}{1} = 4\pi$$

3. The period for  $y = \sin\left(\frac{1}{3}x\right)$  is

$$\text{period} = \frac{2\pi}{\frac{1}{3}} = 2\pi \cdot \frac{3}{1} = 6\pi$$



**Figure 2.1.31** One period each of  $y = \sin(2x)$ ,  $y = \sin(\frac{1}{2}x)$ , and  $y = \sin(\frac{1}{3}x)$  compared to the standard  $y = \sin(x)$  graph. Observe the distinct effects of horizontal compression (when  $B = 2$ , reducing the period to  $\frac{\pi}{2}$ ) and stretching (when  $B = \frac{1}{2}$  and  $B = \frac{1}{3}$ , increasing the period to  $4\pi$  and  $6\pi$ , respectively).

□

Our final transformation involves functions of the form  $y = f(x - c)$ . When  $c > 0$ , the graph of  $y = f(x)$  is shifted  $c$  units to the right; when  $c < 0$ , it is shifted  $|c|$  units to the left.

**Definition 2.1.32 Phase Shift.** The functions

$$y = \sin(B(x - C)) \quad \text{and} \quad y = \cos(B(x - C)) \quad (2.1.1)$$

undergo a **horizontal shift**, known as **phase shift**, of  $C$  units. If  $C > 0$ , the phase shift is to the right; if  $C < 0$ , it is to the left. ◇

**Remark 2.1.33 Functions of the form  $y = \sin(Bx - E)$  and  $y = \cos(Bx - E)$ .** Note that you may see functions written in the form

$$y = \sin(Bx - E) \quad \text{and} \quad y = \cos(Bx - E). \quad (2.1.2)$$

There is a subtle yet important difference between (2.1.1) and (2.1.2). In (2.1.1), the term  $B$ , affecting the period, is multiplied by both  $x$  and  $C$ , the phase shift. In (2.1.2),  $B$  is only multiplied by  $x$ . We can rewrite (2.1.2) by factoring out  $B$  as

$$y = \sin\left(B\left(x - \frac{E}{B}\right)\right) \quad \text{and} \quad y = \cos\left(B\left(x - \frac{E}{B}\right)\right).$$

This form aligns with (2.1.1). Therefore, for equations of the form

$$y = \sin(Bx - E) \quad \text{and} \quad y = \cos(Bx - E)$$

the phase shift is  $\frac{E}{B}$  units.



**Example 2.1.34 Phase Shift.** Identify the period and phase shift of each function, and graph the function

1.  $y = \cos(x - \pi)$

2.  $y = \sin\left(\frac{\pi}{6}(x + 2)\right)$

**Solution.**


1. Since this equation is of the form  $y = \cos(Bx - E)$ , we have  $B = 1$  and  $E = \pi$ . Therefore,

$$\text{period} = \frac{2\pi}{|B|} = \frac{2\pi}{1} = 2\pi$$

and

$$\text{phase shift} = \frac{E}{B} = \frac{\pi}{1} = \pi \text{ (positive value indicates a shift to the right).}$$

A positive value for the phase shift indicates a shift to the right. It's important to note that, given  $B = 1$ , the phase shift is simply  $E = \pi$ .



**Figure 2.1.35** The graph of  $y = \cos(x - \pi)$  is the graph of  $y = \cos x$  with a phase shift of  $\pi$  units to the right.

2. Since the given equation is in the form  $y = \sin\left(\frac{\pi}{6}(x + 2)\right)$ , we can identify  $B = \frac{\pi}{6}$  and  $C = -2$ . Consequently,


$$\text{period} = \frac{2\pi}{|B|} = \frac{2\pi}{\frac{\pi}{6}} = 2\pi \cdot \frac{6}{\pi} = 12$$

and


$$\text{phase shift} = C = -2.$$

The negative sign indicates a phase shift to the left by 2 units.

To graph  $y = \sin\left(\frac{\pi}{6}(x+2)\right)$ , begin by graphing the sine function  $y = \sin\left(\frac{\pi}{6}x\right)$  with a period of 12. Then, apply a phase shift of 2 units to the left on the resulting graph.



**Figure 2.1.36** The graph of  $y = \sin\left(\frac{\pi}{6}x\right)$  represents the sine function  $y = \sin x$  with a horizontal stretch, resulting in a period of 12.



**Figure 2.1.37** Shifting the graph of  $y = \sin\left(\frac{\pi}{6}x\right)$  2 units to the left results in the graph of  $y = \sin\left(\frac{\pi}{6}(x+2)\right)$ .

□

We will now summarize the transformations by consolidating them into a single equation.

**Remark 2.1.38 Transformations of Sine and Cosine.** When dealing with functions in the form

$$y = A \cdot \sin(B(x - C)) + D \quad \text{and} \quad y = A \cdot \cos(B(x - C)) + D$$

we can express the transformations as follows:


- Amplitude and Vertical Compression/Stretch:  $|A|$ 
  - $|A|$  is the value of the amplitude.
  - If  $|A| > 1$ , there is vertical stretching.
  - If  $0 < |A| < 1$ , there is vertical compression.
- Period and Horizontal Stretch/Compression:  $|B|$ 
  - The period is  $\frac{2\pi}{|B|}$ .
  - If  $|B| > 1$ , there is horizontal compression and the period is shortened.
  - If  $0 < |B| < 1$ , there is horizontal stretching and the period is lengthened.
- Phase Shift:  $C$ 
  - If  $C$  is positive, there is a shift to the right.
  - If  $C$  is negative, there is a shift to the left.
- Vertical Shift:  $D$ 
  - If  $D$  is positive, there is a shift upward.
  - If  $D$  is negative, there is a shift downward.
- Reflection about the  $x$ -axis:
  - If  $A$  is negative ( $A < 0$ ), there is a reflection about the  $x$ -axis.
- Reflection about the  $y$ -axis:
  - If  $B$  is negative ( $B < 0$ ), there is a reflection about the  $y$ -axis.

**Remark 2.1.39 Transformations of the form  $y = A \cdot \sin(Bx - E) + D$  and  $y = A \cdot \cos(Bx - E) + D$ .** For functions of the form

$$y = A \cdot \sin(Bx - E) + D \quad \text{and} \quad y = A \cdot \cos(Bx - E) + D$$

the transformations are the same as above, except for the phase shift where you replace  $C$  with  $\frac{E}{B}$ . If  $\frac{E}{B} < 0$  the phase shift is to the right, and if  $\frac{E}{B} > 0$  it is to the left.

Explore the effects of various transformations using the interactive features in [Figure 2.1.40](#).




**Figure 2.1.40** Manipulate the graphs of sine and cosine by adjusting the sliders for  $A$ ,  $B$ ,  $C$ , and  $D$ . Observe the effects on amplitude, period, phase and vertical shifts, as well as reflections about the  $x$ - and  $y$ -axes. Additionally, you can toggle between the sine and cosine graphs by selecting the corresponding function.

### 2.1.5 Exercises

**Exercise Group.** Graph the function.


1.  $y = \sin(x) - 3$

**Answer.**




2.  $y = -2 \sin(x)$

**Answer.**




3.  $y = 4 \cos(x)$

**Answer.**




4.  $y = \cos(x) + 1$

**Answer.**




5.  $y = \frac{1}{2} \sin(x)$

**Answer.**




6.  $y = 5 \cos(x)$

**Answer.**




7.  $y = \sin(x) + 2$

Answer.




8.  $y = -4 \cos(x) - 2$

Answer.




9.  $y = -3 \sin(x) + 2$

Answer.




10.  $y = 4 \cos(x) - 1$

Answer.




11.  $y = 2 \sin(x) - 4$

Answer.



12.  $y = -\frac{1}{2} \cos(x) - 1$


Answer.



**Exercise Group.** Determine the amplitude and period for each function, and sketch the graph.


13.  $y = -2 \sin(3x)$

Answer. Amplitude: 2;  
Period:  $\frac{2\pi}{3}$ ;




14.  $y = 3 \cos(\frac{\pi}{4}x)$

Answer. Amplitude: 3;  
Period: 8;




15.  $y = \frac{1}{2} \sin\left(\frac{\pi}{6}x\right)$

**Answer.** Amplitude:  $\frac{1}{2}$ ;  
Period: 12;




16.  $y = -4 \cos(2x)$

**Answer.** Amplitude: 4;  
Period:  $\pi$ ;




17.  $y = -\frac{2}{3} \sin\left(\frac{\pi}{2}x\right)$

**Answer.** Amplitude:  $\frac{2}{3}$ ;  
Period: 4;




18.  $y = \frac{5}{4} \cos\left(\frac{\pi}{3}x\right)$

**Answer.** Amplitude:  $\frac{5}{4}$ ;  
Period: 6;




19.  $y = \frac{2}{7} \sin\left(\frac{1}{2}x\right)$

**Answer.** Amplitude:  $\frac{2}{7}$ ;  
Period:  $4\pi$ ;









20.  $y = -\frac{3}{5} \cos\left(\frac{\pi}{8}x\right)$

**Answer.** Amplitude:  $\frac{3}{5}$ ;  
Period: 16;



**Exercise Group.** Match the given function to one of the graphs below.





21.  $y = 2 \sin(x) + 1$

Answer. D

23.  $y = -2 \sin(2x)$

Answer. B

25.  $y = -\cos(x - \frac{\pi}{4})$

Answer. C

22.  $y = 3 \cos(4x)$

Answer. E

24.  $y = 3 \sin(\frac{1}{2}x)$

Answer. A


26.  $y = \cos(\frac{1}{2}x) - 1$

Answer. F

**Exercise Group.** Find the amplitude, period, phase shift, and vertical shift of each function and sketch the graph.


27.  $y = \frac{3}{4} \sin(\frac{\pi}{3}(x - 2))$

Answer. Amplitude:  $\frac{3}{4}$ ;  
 Period: 6; Phase Shift: 2;  
 Vertical Shift: 0




28.  $y = -2 \cos(\frac{\pi}{2}(x - \frac{1}{2}))$

Answer. Amplitude: 2;  
 Period: 1; Phase Shift: 0.5;  
 Vertical Shift: 0




29.  $y = 2 \sin(3x - \pi)$

**Answer.** Amplitude: 2;  
 Period:  $\frac{2\pi}{3}$ ; Phase Shift:  $\frac{\pi}{3}$ ;  
 Vertical Shift: 0




30.  $y = -3 \cos(2x + \frac{\pi}{3})$

**Answer.** Amplitude: 2;  
 Period:  $\pi$ ; Phase Shift:  $-\frac{\pi}{6}$ ;  
 Vertical Shift: 0




31.  $y = \frac{1}{2} \sin(\pi(x - 1) + \frac{3}{2})$

**Answer.** Amplitude:  $\frac{1}{2}$ ;  
 Period: 2; Phase Shift: 1;  
 Vertical Shift:  $\frac{3}{2}$




32.  $y = 3 \cos(\frac{\pi}{4}(x + 2)) - 2$

**Answer.** Amplitude:  $\frac{1}{2}$ ;  
 Period: 8; Phase Shift: -2;  
 Vertical Shift: -2




33.  $y = -\frac{2}{3} \sin(2x - \frac{\pi}{3}) + \frac{2}{3}$

**Answer.** Amplitude:  $\frac{2}{3}$ ;  
 Period:  $\pi$ ; Phase Shift:  $\frac{\pi}{6}$ ;  
 Vertical Shift:  $\frac{2}{3}$



34.  $y = \frac{5}{3} \cos(\frac{4\pi}{5}(x + 2))$


**Answer.** Amplitude:  $\frac{5}{3}$ ;  
 Period: 2.5; Phase Shift: -2;  
 Vertical Shift: 0






35.  $y = -\sin\left(\frac{1}{2}x - \frac{\pi}{4}\right) - 1$

**Answer.** Amplitude: 1;  
 Period:  $4\pi$ ; Phase Shift:  $\frac{\pi}{2}$ ;  
 Vertical Shift:  $-1$



36.  $y = 4\cos\left(\frac{1}{2}\left(x + \frac{2\pi}{3}\right)\right) + 3$

**Answer.** Amplitude: 4;  
 Period:  $4\pi$ ; Phase Shift:  $-\frac{2\pi}{3}$ ;  
 Vertical Shift: 3



**Solar Declination.** At the start of this section, we explored the effects of axial tilt on Earth's seasons, considering the sun's declination—the angle between the equator and a line drawn from the center of the Earth to the center of the Sun. When observing the sunrise, the sun's declination is the angle from the sunrise to due east.

During the June Solstice, the declination is  $\delta = 23.45^\circ$ , causing the sun to rise  $23.45^\circ$  to the north of east. On the December Solstice, the declination is  $\delta = -23.45^\circ$ , resulting in the sun rising  $23.45^\circ$  to the south of east.

Conversely, during the spring and fall equinox when the declination is  $\delta = 0^\circ$ , the sun rises precisely at due east. This period holds significant historical importance, as observers, both in the past and present, have utilized this time to precisely determine the eastward direction from their positions. This practice is widespread across various cultures, with individuals using the equinox to establish directional markers. Homes, ceremonial sites, and other notable locations are often intentionally oriented based on the equinox.

In terms of navigation, knowing the solar declination angle for a specific time of year allows observers to measure the same angle down or up from where the sun rose during sunrise, thereby determining the direction of east.

To approximate the solar declination angle  $\delta$  in degrees, we can use the following equation derived in [2.1.6.1]

$$\delta = -23.45^\circ \cdot \cos\left(\frac{360}{365} \cdot (N + 10)\right)$$

where  $N$  represents the day of the year, with January 1 denoted as  $N = 1$ , and December 31 as  $N = 365$ .

For each of the following problems, calculate the solar declination for the given day, assuming a 365-day year. Round your answer to two decimals.

37. March 22nd (81st day of the year).

**Answer.**  $\delta = -0.10^\circ$

39. September 21st (264th day of the year).

**Answer.**  $\delta = -0.10^\circ$

41. April 1st (91st day of the year).

**Answer.**  $\delta = 3.92^\circ$

43. May 28th (148th day of the year).

**Answer.**  $\delta = 21.40^\circ$

38. June 21st (172nd day of the year).

**Answer.**  $\delta = 23.45^\circ$

40. December 21st (355th day of the year).

**Answer.**  $\delta = -23.45^\circ$

42. September 3rd (246th day of the year).

**Answer.**  $\delta = 7.05^\circ$

44. November 23rd (327th day of the year).


**Answer.**  $\delta = -20.8^\circ$

45. What are significant about March 22nd, June 21st, September 21st, and December 21st?


**Answer.** They are the equinoxes and soltices.

46. Graph the solar declination angle over time. Use the horizontal axis for  $N$ , the day of the year, and the vertical axis for  $\delta$ , representing the solar declination angle in degrees.

**Answer.**




**Rough Seas.** **Wave heights**, defined as the vertical distance between the crest and the trough of a wave, can vary in the open ocean. However, the height of the wave alone does not necessarily indicate a calm or choppy sailing conditions. Another important factor is the **wave period**, representing the time between waves, which affects the smoothness of sailing. For each given equation, where  $w(t)$  is the number of feet the wave is above the mean sea level at  $t$  seconds, calculate: a) the wave height; b) the wave period; c) Plot the wave height for two periods.




47.  $w(t) = 4 \cos\left(\frac{\pi}{8}t\right)$

**Answer.** a) 8 feet; b) 16 seconds




48.  $w(t) = 9 \cos\left(\frac{\pi}{8}t\right)$

**Answer.** a) 18 feet; b) 16 seconds



49.  $w(t) = 4 \cos\left(\frac{\pi}{4}t\right)$

**Answer.** a) 18 feet; b) 8 seconds



50. Which of the three waves above would give the smoothest sailing?

**Answer.**  $w(t) = 4 \cos\left(\frac{\pi}{8}t\right)$

### 2.1.6 References

- [1] A. E. Dixon and J. D. Leslie, *Solar Energy Conversio*, Pergamon; (1979)

## 2.2 Graphs of Other Trigonometric Functions

This section explores the graphs of tangent, cotangent, cosecant, and secant, including their periodic behaviors and transformations.

### 2.2.1 Domains of the Tangent and Cotangent Functions

Recall the Quotient Identity for tangent ([Definition 1.4.3](#)):

$$\tan x = \frac{\sin x}{\cos x}$$

Issues arise when the denominator is zero, i.e., when  $\cos x = 0$ . This leads to undefined points at  $x = \dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \dots$ . In general, any angle of the form  $n\frac{\pi}{2}$ , where  $n$  is an odd integer, should be excluded from the domain since tangent is undefined at these values.

Similarly, we defined the cotangent function as

$$\cot x = \frac{\cos x}{\sin x}$$

The denominator becomes zero when  $\sin x = 0$ , corresponding to  $x = \dots, -\pi, 0, \pi, 2\pi, \dots$ . In general,  $\cot x$  is undefined for angles of the form  $n\pi$ , where  $n$  is an integer. These angles should be excluded from the domain of cotangent.

**Remark 2.2.1 Domains for Tangent and Cotangent.** The domains for tangent and cotangent functions are given in [Table 2.2.2](#)

**Table 2.2.2 Domains of the tangent and cotangent functions**

Function	Domain
$\tan \theta$	All real numbers except odd multiples of $\frac{\pi}{2}$ ( $90^\circ$ )
$\cot \theta$	All real numbers except integer multiples of $\pi$ ( $180^\circ$ )

### 2.2.2 Ranges of the Tangent and Cotangent Functions

To determine the range of the tangent function, consider the point  $P(x, y)$  on the unit circle corresponding to the angle  $\theta$ , and let  $a$  be a real number such that  $a = \tan \theta = \frac{y}{x}$ .

Multiplying both sides by  $x$ , we obtain:

$$y = ax$$

Squaring both sides yields:

$$y^2 = a^2x^2$$

Substituting into the Pythagorean Identity ([Definition 1.5.20](#)), we have:

$$1 = x^2 + y^2 = x^2 + a^2x^2 = x^2(1 + a^2)$$

Dividing both sides by  $1 + a^2$  and taking the square root gives:

$$x = \pm \frac{1}{\sqrt{1+a^2}}$$

Similarly, we obtain:

$$y = \pm \frac{a}{\sqrt{1+a^2}}$$

Thus, we conclude:

$$\tan \theta = \frac{y}{x} = \frac{\frac{a}{\sqrt{1+a^2}}}{\frac{1}{\sqrt{1+a^2}}} = a$$

In other words, since  $a$  can be any real number and  $\tan \theta = a$ , the range of the tangent function consists of all real numbers. A similar method can be used to show that the range of the cotangent function is also the set of all real numbers.

**Remark 2.2.3 Ranges for Tangent and Cotangent.** The ranges for tangent and cotangent functions are given in [Table 2.2.4](#)

**Table 2.2.4 Ranges of the tangent and cotangent functions**

Function	Range
$\tan \theta$	All real numbers
$\cot \theta$	All real numbers

### 2.2.3 Domains of the Cosecant and Secant Functions

Consider the Reciprocal Identity for the cosecant function ([Definition 1.4.2](#)):

$$\csc x = \frac{1}{\sin x}$$

When  $\sin x = 0$ , corresponding to  $x = \dots, -\pi, 0, \pi, 2\pi, \dots$ , the denominator becomes zero. In general,  $\csc x$  is undefined for angles of the form  $n\pi$ , where  $n$  is an integer, and these values should be excluded from the domain.

Similarly, since the cosecant function is defined as

$$\sec x = \frac{1}{\cos x}$$

we see that  $\sec x$  is undefined when  $\cos x = 0$ . This occurs at  $x = \dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \dots$ , and thus any angle of the general form  $n\frac{\pi}{2}$ , where  $n$  is an odd integer, should be excluded from the domain of  $\sec x$ .

**Remark 2.2.5 Domains for Cosecant and Secant.** The domains for cosecant and secant functions are given in [Table 2.2.6](#)

**Table 2.2.6** Domains of the trigonometric functions

Function	Domain
$\csc \theta$	All real numbers except integer multiples of $\pi$ ( $180^\circ$ )
$\sec \theta$	All real numbers except integer multiples of $\frac{\pi}{2}$ ( $90^\circ$ )

### 2.2.4 Ranges of the Cosecant and Secant Functions

If the angle is not an integer multiple of  $\pi$ , i.e.  $x \neq n\pi$ , where  $n$  is an integer, then the Reciprocal Identity allows us to define cosecant as

$$\csc x = \frac{1}{\sin x}.$$

In [Subsection 2.1.1](#), we learned that the function  $y = \sin x$  has a range of

$$-1 \leq \sin x \leq 1.$$

Therefore, taking the reciprocal of the range of sine, we get

$$\csc x \leq -1 \text{ or } \csc x \geq 1.$$

In other words, the range of the cosecant function is all real numbers less than or equal to  $-1$  or greater than or equal to  $1$ .

Similarly, since  $-1 \leq \cos x \leq 1$ , we can get the range for the secant function as

$$\sec x \leq -1 \text{ or } \sec x \geq 1$$

or all real numbers less than or equal to  $-1$  or greater than or equal to  $1$ .

**Remark 2.2.7 Ranges for Cosecant and Secant.** The ranges for cosecant and secant functions are given in [Table 2.2.8](#)

**Table 2.2.8** Ranges of the cosecant and secant functions

Function	Range
$\csc \theta$	All real numbers greater than or equal to $1$ or less than or equal to $-1$
$\sec \theta$	All real numbers greater than or equal to $1$ or less than or equal to $-1$

### 2.2.5 The Tangent Function

In [Subsection 1.5.4](#), we learned that the tangent function is periodic with a period of  $\pi$ . To graph  $y = \tan x$ , we focus on plotting one period and then repeat those values to complete the graph.


We also know that the domain of tangent includes all real numbers except any angle of the form  $n\frac{\pi}{2}$ , where  $n$  is an odd integer. These values are excluded since tangent is undefined there. In fact, any line of the form  $x = n\frac{\pi}{2}$  (e.g.  $x = -\frac{\pi}{2}$  and  $x = \frac{\pi}{2}$ ) is a **vertical asymptote**

Knowing the location of the vertical asymptotes, we choose the interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$  to plot our points for tangent. This interval has a length of  $\pi$  (one

period), allowing us to repeat the values to complete the graph of tangent over the entire domain.

Recall from [Definition 1.3.2](#) that on the unit circle, the expression  $\tan \theta = \frac{y}{x}$  denotes the ratio of the  $y$ -coordinate to the  $x$ -coordinate of a point  $P(x, y)$  associated with an angle  $\theta$ . This ratio changes dynamically as  $\theta$  varies from zero to  $\frac{\pi}{2}$ .

As  $\theta$  approaches zero, the  $y$ -value tends to zero, and  $x$  approaches 1, resulting in  $\tan \theta$  being a small fraction. Conversely, as  $\theta$  approaches  $\frac{\pi}{2}$ , the  $y$ -value approaches 1, while  $x$  becomes extremely small and near zero. This causes  $\tan \theta$  to evaluate as a fraction divided by a very small number, producing a large number. A similar effect occurs when  $\theta$  is between  $-\frac{\pi}{2}$  and zero, except  $\tan \theta$  is negative in this range. This behavior is shown in [Figure 2.2.9](#).




**Figure 2.2.9** As  $\theta$  moves from  $-90^\circ$  to  $90^\circ$ , this figure plots the values of  $y = \tan \theta$ . Move the slider for  $\theta$  to see how changing the angle affects  $\tan \theta$ . Note that while we will generally be using radians when graphing trigonometric functions, this figure uses degrees to help visualize the angle.

Next we recall values of tangent for known angles, which are listed in [Table 2.2.10](#) and plotted in [Figure 2.2.11](#).

**Table 2.2.10** Values for  $y = \tan x$

$x$	$\tan x$	$(x, y)$
$-\frac{\pi}{3}$	$-\sqrt{3}$	$(-\frac{\pi}{3}, -\sqrt{3})$
$-\frac{\pi}{4}$	$-1$	$(-\frac{\pi}{4}, -1)$
$-\frac{\pi}{6}$	$-\frac{\sqrt{3}}{3}$	$(-\frac{\pi}{6}, -\frac{\sqrt{3}}{3})$
$0$	$0$	$(0, 0)$
$\frac{\pi}{6}$	$\frac{\sqrt{3}}{3}$	$(\frac{\pi}{6}, \frac{\sqrt{3}}{3})$
$\frac{\pi}{4}$	$1$	$(\frac{\pi}{4}, 1)$
$\frac{\pi}{3}$	$\sqrt{3}$	$(\frac{\pi}{3}, \sqrt{3})$





**Figure 2.2.11** The values in Table 2.2.10 plotted on a graph with a smooth curve connecting the points to make the curve for  $y = \tan(x)$ .

Notice the symmetrical nature of the tangent function's graph with respect to the origin, a feature explained in Section 1.5.7, where we learned that tangent is an odd function.

Since the graph in Figure 2.2.11 represents one period, we can complete the graph of  $y = \tan x$  by extending the pattern in both directions to obtain Figure 2.2.12.




Figure 2.2.12 The plot for  $y = \tan(x)$

### 2.2.6 The Cotangent Function

The domain of the cotangent function is defined for all angles except those of the form  $n\pi$ , where  $n$  is an integer. These excluded values correspond to vertical asymptotes. In fact, any line of the form  $x = n\pi$ , where  $n$  is an integer, serves as a vertical asymptote. Additionally, as we learned in Subsection 1.5.4, the cotangent function has a period of  $\pi$ . We can construct a plot for it in a manner similar to how we constructed the tangent function, as illustrated in Figure 2.2.13.





Figure 2.2.13 As  $\theta$  moves from  $0^\circ$  to  $180^\circ$ , this figure plots the values of  $y = \cot \theta$ . Move the slider for  $\theta$  to see how changing the angle affects  $\cot \theta$ . Note that while we will generally be using radians when graphing trigonometric functions, this figure uses degrees to help visualize the angle.

Plotting points for  $y = \cot x$  and using the fact that the cotangent function is periodic, we obtain the graph for cotangent in Figure 2.2.14



**Figure 2.2.14** The plot for  $y = \cot(x)$ .

The symmetry of the cotangent function about the origin is evident, confirming its nature as an odd function, as explained in [Section 1.5.7](#).

### 2.2.7 The Cosecant Function


The cosecant function has vertical asymptotes at points  $n\pi$ , where  $n$  is an integer, corresponding to the values where the function is undefined. These points are the same ones excluded from the domain of  $\csc x$ . As discussed in [Subsection 1.5.4](#), the cosecant function has a period of  $2\pi$ . Given this, we choose to examine its behavior within one period, specifically from 0 to  $2\pi$ , since this interval spans one complete period of the cosecant function.

Consider the Reciprocal Identity for the cosecant function ([Definition 1.4.2](#)):

$$\csc x = \frac{1}{\sin x}.$$

As  $x$  approaches zero, sine decreases to zero, making cosecant approach positive infinity. Increasing  $x$  towards  $\frac{\pi}{2}$ ,  $\sin(x)$  increases to 1, and cosecant decreases to 1. As  $x$  moves from  $\frac{\pi}{2}$  to  $\pi$ , sine approaches zero, causing  $\csc(x)$  to approach infinity.

Similarly, for  $x > \pi$  nearing  $\pi$ ,  $\sin(x)$  becomes a small, negative number near zero, resulting in  $\csc(x)$  approaching negative infinity. As  $x$  increases to  $\frac{3\pi}{2}$ ,  $\sin(x)$  decreases to  $-1$ , and  $\csc(x)$  increases to  $-1$ . Finally, from  $\frac{3\pi}{2}$  to  $2\pi$ ,  $\sin(x)$  approaches a small negative number, causing cosecant to approach negative infinity. This behavior is shown in [Figure 2.2.15](#).




**Figure 2.2.15** As  $\theta$  moves from  $0^\circ$  to  $360^\circ$ , this figure plots the values of  $y = \sin x$  and  $y = \csc \theta$ . Move the slider for  $\theta$  to see how changing the angle affects  $\csc \theta$ . Note that while we will generally be using radians when graphing trigonometric functions, this figure uses degrees to help visualize the angle.

Next we recall values of sine and cosecant for known angles, which are listed in [Table 2.2.16](#) and plotted in [Figure 2.2.17](#).

**Table 2.2.16** Values for  $y = \csc x$

$x$	$\sin x$	$\csc x$	$x$	$\sin x$	$\csc x$
0	0	Undefined	$\pi$	0	Undefined
$\frac{\pi}{6}$	$\frac{1}{2}$	2	$\frac{7\pi}{6}$	$-\frac{1}{2}$	-2
$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$	$\frac{5\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$-\sqrt{2}$
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{2\sqrt{3}}{3}$	$\frac{4\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$-\frac{2\sqrt{3}}{3}$
$\frac{\pi}{2}$	1	1	$\frac{3\pi}{2}$	-1	-1
$\frac{2\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{2\sqrt{3}}{3}$	$\frac{5\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$-\frac{2\sqrt{3}}{3}$
$\frac{3\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$	$\frac{7\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$-\sqrt{2}$
$\frac{5\pi}{6}$	$\frac{1}{2}$	2	$\frac{11\pi}{6}$	$-\frac{1}{2}$	-2
			$2\pi$	0	Undefined



**Figure 2.2.17** The values in [Table 2.2.16](#) plotted on a graph with a smooth curve connecting the points to make the curve for  $y = \csc(x)$ .

Since the graph in [Figure 2.2.17](#) represents one period, we can complete the graph of  $y = \csc x$  by extending the pattern in both directions to obtain [Figure 2.2.18](#).




Figure 2.2.18 The plot for  $y = \sin(x)$  and  $y = \csc(x)$ .

Notice the graph of the cosecant function is symmetric with respect to the origin, confirming what we learned in Section 1.5.7, that cosecant is an odd function.

### 2.2.8 The Secant Function

As discussed earlier in this section, the secant function has a domain for all real numbers except angles of  $n\frac{\pi}{2}$ , where  $n$  is an integer. These excluded values correspond to the vertical asymptotes of the secant function. With a period of  $2\pi$ , we can focus on the interval 0 to  $2\pi$ . The construction of the secant function plot follows a similar approach to that used for the cosecant function, as illustrated in Figure 2.2.19.





Figure 2.2.19 As  $\theta$  moves from  $0^\circ$  to  $360^\circ$ , this figure plots the values of  $y = \cos x$  and  $y = \sec \theta$ . Move the slider for  $\theta$  to see how changing the angle affects  $\sec \theta$ . Note that while we will generally be using radians when graphing trigonometric functions, this figure uses degrees to help visualize the angle.

Plotting points for  $y = \sec x$  and using the fact that the secant function is periodic, we obtain the graph for secant in Figure 2.2.20.



**Figure 2.2.20** The plot for  $y = \cos(x)$  and  $y = \sec(x)$ .

Notice the graph of the secant function is symmetric about the  $y$ -axis, and thus secant is an even function, confirming what we learned in [Section 1.5.7](#).

## 2.2.9 Graphing Transformations of Other Trigonometric Functions

Similar to the graphs of sine and cosine, the graphs of the other trigonometric functions can undergo vertical stretching and compressing, horizontal stretching and compressing, phase shifts, vertical shift transformations, and reflections about the  $x$ - and  $y$ -axes. However, unlike the sine and cosine functions, there is no amplitude for the other trigonometric functions. These transformations are listed in [Definition 2.2.21](#).

**Definition 2.2.21 Transformations of the Tangent, Cotangent, Cosecant, and Secant Functions.** For functions of the form

$$y = A \cdot \tan(B(x - C)) + D, \quad y = A \cdot \cot(B(x - C)) + D,$$

$$y = A \cdot \csc(B(x - C)) + D, \quad \text{and} \quad y = A \cdot \sec(B(x - C)) + D,$$

we can express the transformations as follows:

- Vertical Compression/Stretch:  $|A|$ 
  - $|A|$  is the value of the vertical stretch/compression.
  - If  $|A| > 1$ , there is vertical stretching.
  - If  $0 < |A| < 1$ , there is vertical compression.
- Period and Horizontal Stretch/Compression:  $|B|$ 
  - The period is  $\frac{\pi}{|B|}$  for tangent and cotangent, and  $\frac{2\pi}{|B|}$  for cosecant and secant.

- If  $|B| > 1$ , there is horizontal compression, and the period is shortened.
- If  $0 < |B| < 1$ , there is horizontal stretching, and the period is lengthened.
- Phase Shift:  $C$ 
  - If  $C$  is positive, there is a shift to the right.
  - If  $C$  is negative, there is a shift to the left.
- Vertical Shift:  $D$ 
  - If  $D$  is positive, there is a shift upward.
  - If  $D$  is negative, there is a shift downward.
- Reflection about the  $x$ -axis:
  - If  $A$  is negative ( $A < 0$ ), there is a reflection about the  $x$ -axis.
- Reflection about the  $y$ -axis:
  - If  $B$  is negative ( $B < 0$ ), there is a reflection about the  $y$ -axis.
- Vertical Asymptotes:
  - For tangent and cotangent, vertical asymptotes occur at

$$x = C + \frac{\pi}{|B|}n,$$

where  $n$  is an integer.

- For cosecant and secant, vertical asymptotes occur at

$$x = C + \frac{\pi}{2|B|}n,$$

where  $n$  is an integer.

◇

**Remark 2.2.22 Other Forms of Transformations.** For functions of the form

$$y = A \cdot \tan(Bx - E) + D, \quad y = A \cdot \cot(Bx - E) + D,$$

$$y = A \cdot \csc(Bx - E) + D, \quad \text{and} \quad y = A \cdot \sec(Bx - E) + D,$$

the transformations are the same as above, except for the phase shift and vertical asymptotes where you replace  $C$  with  $\frac{E}{B}$ . If  $\frac{E}{B} < 0$  the phase shift is to the right, and if  $\frac{E}{B} > 0$  it is to the left.

- For tangent and cotangent, vertical asymptotes occur at

$$x = \frac{E}{B} + \frac{\pi}{|B|}n,$$

where  $n$  is an integer.

- For cosecant and secant, vertical asymptotes occur at

$$x = \frac{E}{B} + \frac{\pi}{2|B|}n,$$




where  $n$  is an integer.

**Example 2.2.23 Vertical Stretch/Compression and Reflection about the  $x$ -axis.** Graph each function

1.  $y = \tan(x)$
2.  $y = 2 \tan(x)$
3.  $y = \frac{1}{2} \tan(x)$
4.  $y = -\tan(x)$

**Solution.**



**Figure 2.2.24** The transformations of the tangent function graph, starting with a baseline of  $y = \tan(x)$  along with graphs with a vertical stretch, a vertical compression, and a reflection about the  $x$ -axis.

□

**Example 2.2.25 Horizontal Stretch/Compression and Reflection about the  $y$ -axis.** Identify the period and graph one period for each of the following functions:

1.  $y = \cot(x)$
2.  $y = \cot(2x)$
3.  $y = \cot\left(\frac{1}{2}x\right)$

4.  $y = \cot(-x)$

**Solution.**

1. The period for  $y = \cot(x)$  is  $\pi$

2. The period for  $y = \cot(2x)$  is


$$\text{period} = \frac{\pi}{|2|} = \frac{\pi}{2}$$

3. The period for  $y = \cot\left(\frac{1}{2}x\right)$  is

$$\text{period} = \frac{\pi}{\left|\frac{1}{2}\right|} = 2\pi$$

4. The period for  $y = \cot(-x)$  is

$$\text{period} = \frac{\pi}{|-1|} = \pi$$




**Figure 2.2.26** The transformations of the cotangent function graph, starting with a baseline of  $y = \cot(x)$  along with graphs with a horizontal stretch, a horizontal compression, and a reflection about the  $y$ -axis.

□

**Example 2.2.27 Phase Shifts and Vertical Shifts.** Graph the function  $y = \csc\left(x - \frac{\pi}{2}\right) + 2$ .


**Solution.**




**Figure 2.2.28** The graphs of  $y = \csc(x - \frac{\pi}{2}) + 2$  and  $y = \sin(x - \frac{\pi}{2}) + 2$  start with the plots of  $y = \csc(x)$  and  $y = \sin(x)$  and add a phase shift of  $\frac{\pi}{2}$  to the right and a vertical shift up by 2.

□

Explore the effects of various transformations using the interactive features in [Figure 2.2.29](#) and [Figure 2.2.30](#).



**Figure 2.2.29** Manipulate the graphs of tangent and cotangent by adjusting the sliders for  $A$ ,  $B$ ,  $C$ , and  $D$ . Observe the effects on period, phase and vertical shifts, as well as reflections about the  $x$ - and  $y$ -axes. Additionally, you can toggle between the tangent and cotangent graphs by selecting the corresponding function.




**Figure 2.2.30** Manipulate the graphs of cosecant and secant by adjusting the sliders for  $A$ ,  $B$ ,  $C$ , and  $D$ . Observe the effects on period, phase and vertical shifts, as well as reflections about the  $x$ - and  $y$ -axes. Additionally, you can toggle between the cosecant and secant graphs by selecting the corresponding function.

### 2.2.10 Exercises

**Exercise Group.** Graph the function.


1.  $y = \sec(x) - 4$

Answer.




2.  $y = \cot(x) - 1$

Answer.




3.  $y = \csc(x) + 3$

Answer.




4.  $y = \tan(x) + 2$

Answer.




5.  $y = \frac{1}{2} \tan(x)$

Answer.




6.  $y = -3 \csc(x)$

Answer.




7.  $y = 2 \cot(x)$

Answer.




8.  $y = 4 \sec(x)$

Answer.




9.  $y = -\frac{1}{2} \tan(2x) + 2$

Answer.




10.  $y = 2 \csc(3x) + 3$

Answer.




11.  $y = 3 \cot(\frac{1}{2}x) - 1$

Answer.



12.  $y = -3 \sec(2x) - 4$


Answer.



**Exercise Group.** Determine the period for each function, and sketch the graph.


13.  $y = \cot(\frac{1}{2}x)$

Answer. Period:  $2\pi$




14.  $y = \tan(\frac{\pi}{3}x)$

Answer. Period: 3




15.  $y = \sec(\frac{\pi}{4}x)$

Answer. Period: 8




16.  $y = \csc(\frac{\pi}{6}x)$

Answer. Period: 12




17.  $y = \tan(2x)$

Answer. Period:  $\frac{\pi}{2}$




18.  $y = \sec\left(\frac{\pi}{2}x\right)$

Answer. Period: 4




19.  $y = \csc\left(\frac{1}{4}x\right)$

Answer. Period:  $8\pi$








20.  $y = \cot\left(\frac{\pi}{6}x\right)$

Answer. Period: 6



**Exercise Group.** Match the given function to one of the graphs below.





21.  $y = \csc(2x) + 4$


Answer. C

23.  $y = 3 \cot(x - \frac{\pi}{2})$

Answer. E

25.  $y = -2 \csc(2x)$

Answer. F



22.  $y = 2 \tan(x - \frac{\pi}{4})$

Answer. A

24.  $y = -\frac{1}{3} \cot(x)$

Answer. B


26.  $y = \sec(x + \frac{\pi}{3}) - 2$

Answer. D

**Exercise Group.** Find the period, phase shift, and vertical shift of each function and sketch the graph.


27.  $y = 2 \tan(x) + 3$

Answer. Period:  $\pi$ ; Phase Shift: 0; Vertical Shift: 3




28.  $y = \frac{1}{2} \cot(\frac{1}{2}x) + 1$

Answer. Period:  $2\pi$ ; Phase Shift: 0; Vertical Shift: 1




29.  $y = -\csc(x + \frac{\pi}{4})$

Answer. Period:  $2\pi$ ; Phase Shift:  $-\frac{\pi}{4}$ ; Vertical Shift: 0




30.  $y = -\frac{1}{3} \sec(2x)$

Answer. Period:  $\pi$ ; Phase Shift: 0; Vertical Shift: 0




31.  $y = 2 \sec\left(\frac{\pi}{2}x\right) + 4$

**Answer.** Period: 4; Phase Shift: 0; Vertical Shift: 4




32.  $y = \cot\left(x - \frac{\pi}{3}\right) + 2$

**Answer.** Period:  $\pi$ ; Phase Shift:  $\frac{\pi}{3}$ ; Vertical Shift: 2




33.  $y = \tan\left(x - \frac{\pi}{6}\right) + 1$

**Answer.** Period:  $\pi$ ; Phase Shift:  $\frac{\pi}{6}$ ; Vertical Shift: 1




34.  $y = -\frac{1}{4} \sec(x)$

**Answer.** Period:  $2\pi$ ; Phase Shift: 0; Vertical Shift: 0




35.  $y = 3 \csc\left(x + \frac{\pi}{2}\right)$

**Answer.** Period:  $2\pi$ ; Phase Shift:  $-\frac{\pi}{2}$ ; Vertical Shift: 0



36.  $y = -\tan\left(2\left(x - \frac{\pi}{4}\right)\right) + 2$

**Answer.** Period:  $\frac{\pi}{2}$ ; Phase Shift:  $\frac{\pi}{4}$ ; Vertical Shift: 2



**Navigating with Shadows.** When navigating the open ocean, maintaining a straight course poses challenges due to limited visual markers. One technique involves the steersperson using the positions of shadows cast by objects on the canoe—such as crew members, railings, and sails—to keep them fixed on the deck, ensuring a straight trajectory. However, if the canoe veers off course, the changing position of the canoe relative to the sun leads to a shift in the shadows. Observing these shadow movements allows the steersperson to make course corrections. It's important to note that this method is effective only over a short duration, as the sun's continuous movement across the sky causes ongoing changes in shadow positions. To illustrate the limitations over extended periods, consider the example of the Samoan double-hulled voyaging va'a, Gaulofa,



with a 14-meter-high mast. The length of the shadow is modeled by

$$l(t) = 14 \left| \cot \left( \frac{\pi}{12} t \right) \right|,$$

where  $l$  is the shadow length in meters and  $t$  represents the hours since 6 am (assuming sunrise at 6 am and sunset at 6 pm). In each of the following questions, calculate the length of the shadow, rounded to the nearest tenth of a meter, for the given time.

37. 7:00am

**Answer.** 52.2 meters

39. 12:00pm

**Answer.** 0 meters

41. 4:00pm

**Answer.** 24.2 meters

38. 10:00am


**Answer.** 8.1 meters

40. 3:00pm


**Answer.** 14 meters

42. Graph the length of the shadow,  $l$ , throughout the day from 6:00am to 6:00pm ( $0 < t < 12$ )

**Answer.**



**Exercise Group.**




An observer on Rangiroa spots the Fa'afaite, a double-hulled voyaging canoe from Tahiti, sailing off the north coast of the atoll, maintaining a distance of three nautical miles from the shore and traveling east. Let  $\theta$  represent the angle formed between the line from the observer to the va'a and a line extending due north from the observer, measured in radians. The angle  $\theta$  is negative if the va'a is to the left of the observer and positive when to the right, as shown in the figure above. The distance (in nautical miles), denoted by  $d(\theta)$  from Fa'afaite to the observer is given by the function

$$d(\theta) = 3 \sec(\theta).$$

In each of the following questions, calculate the distance from the observer to Fa'afaite,  $d(\theta)$ , in nautical miles, for the given angle  $\theta$ . Round your answer to two decimal places.

43.  $\theta = -\frac{\pi}{3}$   
**Answer.** 6.00 NM
45.  $\theta = -\frac{\pi}{6}$   
**Answer.** 3.46 NM
47.  $\theta = \frac{\pi}{6}$   
**Answer.** 3.46 NM
49.  $\theta = \frac{\pi}{3}$   
**Answer.** 6.00 NM

44.  $\theta = -\frac{\pi}{4}$   
**Answer.** 4.24 NM
46.  $\theta = 0$   
**Answer.** 6.00 NM
48.  $\theta = \frac{\pi}{4}$   
**Answer.** 4.24 NM
50. Graph the function  $d(\theta)$  on the domain  $-\frac{\pi}{3} \leq \theta \leq \frac{\pi}{3}$ .  
**Answer.**



51. What happens to  $d$  as  $\theta$  approaches  $\frac{\pi}{2}$ ?  
**Answer.** As  $\theta$  approaches  $\frac{\pi}{2}$ , the function  $d(\theta) = 3 \sec(\theta)$  approaches positive infinity.

52. What is the closest distance Fa'afaite comes to shore? Where does this occur?  
**Answer.** 3 NM, when Fa'afaite is directly north of the observer ( $\theta = 0$ )

## 2.3 Sinusoidal Curve Fitting and Graphical Analysis

In this chapter, we have learned that sinusoidal patterns exist in various aspects throughout our world. One example is the moon, which undergoes phases oscillating from no illumination, to waxing (increasing illumination), reaching a fully lit moon, and then waning (decreasing illumination) until it completes its cycle with no illumination again. Indigenous cultures across the world have deeply connected with these lunar cycles, shaping cultural practices aligned with the moon's phases. For example, the Māori of New Zealand and the Hopi Tribe in northeastern Arizona, USA, both time activities like planting and harvesting specific plants for each moon cycle based on generations of lunar observations.

In [Section 2.1](#), we graphed sinusoidal functions and determined their values at any given time. The ability to formulate a sinusoidal equation modeling the moon's phases allows us to predict the moon's phase on any date. In this section, we will explore the process of developing sinusoidal functions based on provided information. This will enable us to model real-world phenomena using real data.

### 2.3.1 Finding Sinusoidal Equations from Characteristics

To begin finding sinusoidal equations of the form

$$y = A \cdot \sin(B(x - C)) + D \quad \text{and} \quad y = A \cdot \cos(B(x - C)) + D$$

we will refer to the characteristics of the sine and cosine functions described in ("sine-cosine-transformations") and given below to help us determine the parameter values for  $A$ ,  $B$ ,  $C$ , and  $D$ .

- Amplitude and Vertical Compression/Stretch:  $|A|$ 
  - $|A|$  is the value of the amplitude.
  - If  $|A| > 1$ , there is vertical stretching.
  - If  $0 < |A| < 1$ , there is vertical compression.
- Period and Horizontal Stretch/Compression:  $|B|$ 
  - The period is  $\frac{2\pi}{|B|}$ .
  - If  $|B| > 1$ , there is horizontal compression and the period is shortened.
  - If  $0 < |B| < 1$ , there is horizontal stretching and the period is lengthened.
- Phase Shift:  $C$ 
  - If  $C$  is positive, there is a shift to the right.
  - If  $C$  is negative, there is a shift to the left.
- Vertical Shift:  $D$ 
  - If  $D$  is positive, there is a shift upward.
  - If  $D$  is negative, there is a shift downward.

- Reflection about the  $x$ -axis:
  - If  $A$  is negative ( $A < 0$ ), there is a reflection about the  $x$ -axis.
- Reflection about the  $y$ -axis:
  - If  $B$  is negative ( $B < 0$ ), there is a reflection about the  $y$ -axis.

**Example 2.3.1 Finding an Equation of a Sine Function Using its Characteristics.** Write an equation of a sine function with an amplitude of 2, period of 8, phase shift of  $\frac{\pi}{3}$ , and vertical shift of 4.

**Solution.** To determine an equation of the form

$$f(x) = A \cdot \sin(B(x - C)) + D$$

we first examine the information that we are given: amplitude of 2, period of 8, phase shift of  $\frac{\pi}{3}$ , and vertical shift of 4.

Since  $|A|$  represents the value of the amplitude, we get

$$|A| = 2.$$

Thus we either have  $A = 2$  or  $A = -2$ . Since we are not given a reflection about the  $x$ -axis, we can conclude that  $A$  is not negative, thus

$$A = 2.$$

Next, since the period of a sine function is given by  $\frac{2\pi}{|B|}$ , we get

$$\frac{2\pi}{|B|} = 8.$$

Solving for  $|B|$ , we get

$$|B| = \frac{2\pi}{8} = \frac{\pi}{4}$$

Since there is no reflection about the  $y$ -axis, we have that  $B$  must be positive:

$$B = \frac{\pi}{4}.$$

Finally, since  $C$  and  $D$  represent the phase shift and vertical shift, respectively, we get

$$C = \frac{\pi}{3}, \quad D = 4.$$

Combining these, the sine function becomes:

$$f(x) = 2 \sin\left(\frac{\pi}{4}\left(x - \frac{\pi}{3}\right)\right) + 4.$$


□

### 2.3.2 Finding Sinusoidal Equations from Graphs

Sometimes we are not explicitly given the characteristics of the function, but are provided with the graph. Examining a graph can reveal its characteristics, allowing us to find the equation of a function.

In this next example, we'll explore how to find an equation of a cosine function based on the graph of a sine function.

**Example 2.3.2 Finding an Equation of a Cosine Function Using the Graph of a Sine Function.** Below is the graph of  $y = \sin(x)$ .



Find an equation of the form  $y = A \cos(B(x - C)) + D$  that fits the graph.

**Solution.** The characteristics of the cosine start its maximum when  $x = 0$ , decrease to its minimum at  $x = \pi$ , and then increase before completing one period at  $2\pi$ . This graph starts at 0 when  $x = 0$ , then increases to its maximum at  $x = \frac{\pi}{2}$ , and then follows the characteristics of the cosine graph. This represents a phase shift to the right. There no reflections about the  $x$ -axis or  $y$ -axis. We will find the equation of a function with characteristics of a phase shift to the right.

*Amplitude (A):* Since the amplitude of the cosine function will be the same as the amplitude of the sine function, which is 1, we have  $|A| = 1$ . Since there is no reflection about the  $x$ -axis, we choose to positive value to get  $A = 1$ .


*Vertical Shift (D):* The vertical shift of the cosine function will be the same as that of the sine function, which is 0. So, we have  $D = 0$ .

*Period (B):* The period of the cosine function will also be the same as the period of the sine function, which is  $2\pi$ . Since  $2\pi = \frac{2\pi}{|B|}$ , we have  $|B| = 1$ . Since there is no reflection about the  $y$ -axis, we get  $B = 1$ .

Therefore, we have

$$y = 1 \cos(1x) + 0 = \cos(x).$$

Overlapping the graph of  $y = \cos(x)$  onto the original graph will help us determine the phase shift.



*Phase Shift (C):* From the graph, the cosine function will have a phase shift of  $\frac{\pi}{2}$  radians to the right compared to the sine function. This is because the cosine function reaches its maximum value at  $x = 0$ , while the sine function reaches its maximum at  $x = \frac{\pi}{2}$ . So, we have  $C = \frac{\pi}{2}$ .

Therefore, the equation of the cosine function that fits the graph of  $y = \sin(x)$  is:

$$y = \cos\left(x - \frac{\pi}{2}\right)$$


□

**Remark 2.3.3** Since this graph is also the graph of  $y = \sin(x)$ , we get

$$\sin(x) = \cos\left(x - \frac{\pi}{2}\right).$$

This solution demonstrates that any sine function can also be written as a cosine function, with an appropriate phase shift. In this case, the phase shift of  $\frac{\pi}{2}$  radians to the right converts the sine function to its corresponding cosine function.

**Example 2.3.4 Finding an Equation of a Sine Function Using its Graphs.** Find an equation that represents the following graph of the form



(a)

$$f(x) = A \sin(B(x - C)) + D$$

**Solution.** The characteristics of the sine function start at the origin, increase to its maximum when  $x = \frac{\pi}{2}$ , decrease to its minimum at  $x = \frac{3\pi}{2}$ , and then increase again before completing one period at  $2\pi$ . In the given graph, the function starts at 8 when  $x = 0$ , then increases to its maximum value of 16 at  $x = 10$ , following the characteristics of the sine graph, before completing one period at  $x = 40$ . This indicates a vertical shift, a vertical stretch, and a horizontal stretching of the period. There are no reflections about the  $x$ -axis or  $y$ -axis. Therefore, we need to find the equation of a function with these characteristics.

*Amplitude ( $A$ ):* Recall from [Definition 2.1.24](#) that the amplitude is defined as half the difference between the maximum and minimum values of the function. Here, the maximum value is 16 and the minimum value is 0, so the amplitude is:

$$|A| = \frac{16 - 0}{2} = 8.$$

Since there are no reflections about the  $x$ -axis, we use the positive value to get  $A = 8$ .

*Vertical Shift ( $D$ ):* The midline, calculated as

$$y = \frac{\max + \min}{2} = \frac{16 + 0}{2} = 8,$$

represents the vertical shift of the graph. Therefore,

$$D = 8.$$

*Period ( $B$ ):* The period of a function is the length of one cycle. You can identify the period on the graph by measuring the horizontal distance

between corresponding points where the graph completes a cycle. In this case, we'll use two corresponding peaks at  $x = 10$  and  $x = 50$  to obtain:

$$\text{period} = 50 - 10 = 40.$$

Note that other points on the graph, such as the minimum values or where the graph crosses the midline, could also be used to determine the period.

Since [Definition 2.1.28](#) defines a period as  $\frac{2\pi}{|B|}$ , we have:

$$40 = \frac{2\pi}{|B|}$$

Thus,

$$|B| = \frac{2\pi}{40} = \frac{\pi}{20}.$$

Since there are no reflections about the  $y$ -axis, we get  $B = \frac{\pi}{20}$ .

*Phase Shift (C):* The phase shift of the graph refers to its horizontal translation. The characteristics of a sine function typically start at  $x = 0$  on the midline, increase to the maximum, decrease, pass the midline to the minimum, and then complete a cycle back at the midline. Since this graph follows these characteristics without any horizontal translation, there is no phase shift. Therefore,  $C = 0$ .

Thus, the graph can be described by the following sine function:

$$f(x) = 8 \sin\left(\frac{\pi}{20}x\right) + 8$$

(b)

$$g(x) = A \cos(B(x - C)) + D$$

**Solution.** Since the cosine function starts at the maximum value when  $x = 0$  before decreasing to its minimum and then increasing to complete one period, the graph provides several options for phase shifts. By selecting a peak and determining the direction and value of the phase shift needed for the cosine function to reach that peak, we can align it with the desired position. For example, if we choose the peak at  $x = -30$ , the cosine function will shift 30 units to the left. Similarly, for peaks at  $x = 10$  or  $x = 50$ , the cosine function would need to shift 10 or 50 units, respectively, to the right. Since the peak at  $x = 10$  aligns closely with the peak of the original cosine function when  $x = 0$ , we opt for this phase shift. Additionally, similar to the graph of the sine function, there is a vertical shift, a vertical stretch, and a horizontal stretch, with no reflections about the  $y$ -axis or the  $x$ -axis.

*Amplitude (A):* The amplitude of a cosine function is the same as that of the corresponding sine function. Thus,  $|A| = 8$ . Since there are no reflections about the  $x$ -axis, we use the positive value to get  $A = 8$ .

*Vertical Shift (D):* The vertical shift of a cosine function is the same as that of the corresponding sine function. Thus,  $D = 8$ .


*Period (B):* The period of a cosine function is also the same as that of the corresponding sine function. In this case,  $|B| = \frac{\pi}{20}$ . Since there are no reflections about the  $y$ -axis, we determine  $B = \frac{\pi}{20}$ .

Before we examine the phase shift, let's summarize what we found so far:

$$g(x) = 8 \cos\left(\frac{\pi}{20}(x)\right) + 8$$



and overlap this graph with the graph of the original.



*Phase Shift (C):* From the graph, we see that the starting point of the cosine function's cycle is at its maximum value, unlike the starting point of a sine function's cycle, which is at its midline. In this case, the graph has a horizontal shift to the right of 10 units. Thus,  $C = 10$ .

We can now describe the graph with the following cosine function:


$$g(x) = 8 \cos\left(\frac{\pi}{20}(x - 10)\right) + 8.$$

□

### 2.3.3 Finding Sinusoidal Equations from Data

**Example 2.3.5 Modeling the Daylight Hours in Munda.** In [Section 2.1](#) we learned that Earth's axis is tilted, and as the Earth orbits the sun, this axial tilt causes seasons, which are periodic. Another effect of the axial tilt and orbit is the amount of daylight each part of Earth experiences, which is also periodic.

The total sunlight duration in Munda, on the island of New Georgia in the Solomon Islands in 2025, is plotted in [Figure 2.3.6](#). On the 21st of June 2025 (the 172nd day of the year), which is the shortest day of the year in Munda, the total sunlight duration is 11 hours, 38 minutes, and 15 seconds (approximately 11.64 hours). Conversely, on the 22nd of December 2025 (the 356th day of the year), which is the longest day of the year, the total sunlight duration is 12 hours, 36 minutes, and 28 seconds (approximately 12.61 hours). Find a function of the form  $y = A \cos(B(x - C)) + D$  to model the total hours of daylight in Munda in 2025, assuming that one period represents one year or 365 days.



**Figure 2.3.6** Hours of daylight in Munda, Solomon Islands. Source: NOAA Solar Calculator.

**Solution.** *Amplitude (A):* From the definition of amplitude ([Definition 2.1.24](#)),

$$|A| = \frac{\max - \min}{2} = \frac{12.61 - 11.64}{2} = 0.485.$$

Assuming no reflections about the  $x$ -axis, we have:

$$A = 0.485.$$

*Period (B):* We assume a period of one year or 365 days. Additionally, we can assume no reflections about the  $y$ -axis, leading to a positive value for  $B$ . Thus we have  $\frac{2\pi}{B} = 365$  or

$$B = \frac{2\pi}{365} \approx 0.0172.$$

*Phase Shift (C):* The characteristic of a cosine function is that at  $x = 0$ , the function is at its maximum value. However, in this case, the maximum value occurs on Day 356. This represents a phase shift to the right by 356 days, thus:

$$C = 356.$$

*Vertical Shift (D):* The value of the midline represents the average duration of daylight, which is the vertical shift:

$$D = \frac{\max + \min}{2} = \frac{12.61 + 11.64}{2} = 12.125.$$

Therefore, the equation of our function is

$$y = 0.485 \cos(0.0172(x - 356)) + 12.125.$$

□


**Remark 2.3.7** It's important to note that the average duration of daylight is not exactly 12 hours but instead approximately 12.125 hours. This discrepancy arises due to atmospheric refractions, which cause the apparent sunrise and sunset to occur slightly before and after, respectively, the sun crosses the horizon—the actual sunrise and sunset times.

**Example 2.3.8 Modeling the Temperate in Christchurch.** Another effect of axial tilt besides daylight hours is temperature, which is also periodic. The average monthly temperature for Christchurch, New Zealand is given in [Table 2.3.9](#) (Source: National Institute of Water and Atmospheric Research (NIWA). Retrieved 18 March 2024). Find a sinusoidal function of the form  $y = A \cos(B(x - C)) + D$  to model the average monthly temperature of Christchurch.

**Table 2.3.9** The average monthly temperature for Christchurch, New Zealand.

Month, $x$	Temperature ( $^{\circ}\text{C}$ )
January, 1	17.5
February, 2	17.2
March, 3	15.5
April, 4	12.7
May, 5	9.8
June, 6	7.1
July, 7	6.6
August, 8	7.9
September, 9	10.3
October, 10	12.2
November, 11	14.1
December, 12	16.1

**Solution.** We begin by plotting the points in [Table 2.3.9](#).



Examining the plot of the data points, we see this looks like the graph of a cosine function with no reflections about the  $x$ -axis or  $y$ -axis.

*Amplitude (A):* The amplitude is

$$|A| = \frac{\max - \min}{2} = \frac{17.5 - 6.6}{2} = 5.45.$$

Since there are no reflections about the  $x$ -axis, we get  $A = 5.45$

*Period (B):* Since the temperatures repeat every 12 months, the period is 12 and so  $\frac{2\pi}{|B|} = 12$ . Since there are no reflections about the  $y$ -axis, we keep the positive value to obtain

$$B = \frac{2\pi}{12} = \frac{\pi}{6}.$$

*Vertical Shift (D):* The vertical shift is the value of the average data:


$$D = \frac{\max + \min}{2} = \frac{17.5 + 6.6}{2} = 12.05.$$

*Phase Shift (C):* The maximum temperature in the data occurs in January ( $x = 1$ ). However, since the graph of cosine reaches its maximum value at  $x = 0$ , we have a phase shift of 1 to the right to align the peak with the maximum temperature data point. Thus,  $C = 1$ .

Our function now becomes

$$y = 5.45 \cos\left(\frac{\pi}{6}(x - 1)\right) + 12.05.$$

Finally, we plot our function and the data together.



□

**Remark 2.3.10 Steps for Deriving Sinusoidal Models from Data.**

1. Graph the data points.

- Determine the characteristics of the data, including vertical and horizontal stretching, phase shifts, vertical shifts, and reflections about the  $x$ -axis or  $y$ -axis.
- Amplitude*: Calculate the amplitude using the formula

$$|A| = \frac{\max - \min}{2}.$$

Use the positive value if there is no reflection about the  $x$ -axis, and use the negative value if there is a reflection.

- Vertical Shift*: Determine the vertical shift which is the average value of the data, using the formula

$$D = \frac{\max + \min}{2}.$$

- Period*: Calculate the period from the data, then find

$$|B| = \frac{2\pi}{\text{period}}.$$

Use the positive value if there is no reflection about the  $y$ -axis, and use the negative value if there is a reflection.

- Phase Shift*: Determine the phase shift by comparing points from the data to those on the sine and cosine function, such as maximum and minimum values as well as those values at the midline. This information will give you the phase shift,  $C$ .

### 2.3.4 Finding Sinusoidal Equations with Technology

Some graphing utilities, such as the TI-83 calculator or [Desmos Graphing Calculator](#)<sup>1</sup>, have functions that allow you to find a sinusoidal best-fit function given data. While some devices can find the best-fit for both sine and cosine functions, others only calculate the best-fit line for sine. For example, the TI-83 uses the `SinReg` function to calculate the sine function.

**Example 2.3.11 Finding the Sinusoidal Equation using Desmos.** Utilize a graphing utility to determine the best-fit cosine function for the data provided in [Table 2.3.9](#).

**Solution.**

- Open a new table in the Desmos Graphing Calculator by either typing “table” in a blank expression line or clicking the Add Item menu in the upper left corner and selecting Table.
- Enter the values from [Table 2.3.9](#) into the table, where  $x_1$  represents the month and  $y_1$  represents the temperature.
- Use the Zoom Fit icon (a magnifying glass with a + symbol) at the bottom left corner of the table to automatically adjust the graph settings window to best display your data.
- In a blank expression line, type “ $y_1 \sim A \cos(B(x_1 - C)) + D$ ” to fit a cosine function to the data.

---

<sup>1</sup>[DesmosGraphingCalculator](#)

5. The parameters for the best-fit function will be returned:  $A = 5.2141$ ,  $B = 0.534145$ ,  $C = 1.19926$ ,  $D = 12.1509$ . Thus, the best-fit cosine function is:

$$y = 5.2141 \cos(0.534145(x - 1.19926)) + 12.1509.$$

Figure 2.3.12 displays an interactive Desmos Graphing Calculator with the completed table and the line of best fit plotted together.




Figure 2.3.12 Given data points in a table, Desmos can create a sinusoidal function to model the data. □

### 2.3.5 Exercises

**Exercise Group.** Write the equation of a sine function with the following characteristics:

- |   |   |
|---|---|
| 1. Amplitude: 2; Period: $\pi$ .<br><b>Answer.</b> $y = 2 \sin(2x)$   | 2. Amplitude: 3; Period: $\frac{\pi}{2}$ .<br><b>Answer.</b> $y = 3 \sin(4x)$   |
| 3. Amplitude: 1.5; Period: $\frac{\pi}{6}$ ; Reflection about the $x$ -axis.<br><b>Answer.</b> $y = -1.5 \sin(12x)$                                     | 4. Amplitude: 2; Period: $3\pi$ ; Reflection about the $x$ -axis.<br><b>Answer.</b> $y = -2 \sin\left(\frac{2}{3}x\right)$                                    |
| 5. Amplitude: 4; Period: $\frac{2\pi}{3}$ ; Vertical Shift 2.<br><b>Answer.</b> $y = 4 \sin(3x) + 2$  | 6. Amplitude: 3; Period: $\frac{5}{2}$ ; Vertical Shift: $-\frac{2}{3}$ .<br><b>Answer.</b> $y = 3 \sin\left(\frac{4\pi}{5}x\right) - \frac{2}{3}$            |
| 7. Amplitude = $\frac{1}{2}$ ; Period = $2\pi$ ; Phase Shift = $\frac{\pi}{4}$ .<br><b>Answer.</b> $y = \frac{1}{2} \sin\left(x - \frac{\pi}{4}\right)$ | 8. Amplitude: 1; Period: $\frac{5\pi}{3}$ ; Phase Shift: $\frac{\pi}{6}$ .<br><b>Answer.</b> $y = \sin\left(\frac{6}{5}\left(x - \frac{\pi}{6}\right)\right)$ |


9. Amplitude: 2, Period:  $\frac{3\pi}{2}$ ; Phase Shift =  $\frac{\pi}{3}$ ; Vertical Shift = -1.  
**Answer.**  $y = 2 \sin\left(\frac{4}{3}\left(x - \frac{\pi}{3}\right) - 1\right)$
10. Amplitude = 2, Period =  $20$ ; Phase Shift =  $-\frac{\pi}{4}$ ; Vertical Shift = 3.  
**Answer.**  $y = 2 \sin\left(\frac{\pi}{10}\left(x + \frac{\pi}{4}\right) + 3\right)$

**Exercise Group.** Write the equation of a cosine function with the following characteristics:

11. Amplitude = 1.8; Period =  $2\pi$ .  
**Answer.**  $y = 1.8 \cos(x)$
12. Amplitude = 2.5; Period =  $\frac{\pi}{4}$ .  
**Answer.**  $y = 2.5 \cos(8x)$
13. Amplitude = 2; Period =  $\frac{7\pi}{4}$ ; Reflection in the  $x$ -axis.  
**Answer.**  $y = -2 \cos\left(\frac{8}{7}x\right)$
14. Amplitude = 1.5; Period =  $\frac{4\pi}{5}$ ; Reflection in the  $x$ -axis.  
**Answer.**  $y = -1.5 \cos\left(\frac{5}{2}x\right)$
15. Amplitude = 4; Period =  $\frac{5\pi}{3}$ ; Vertical Shift = -3.  
**Answer.**  $y = 4 \cos\left(\frac{6}{5}x\right) - 3$
16. Amplitude = 3.5; Period =  $\frac{4\pi}{3}$ ; Vertical Shift = 2.  
**Answer.**  $y = 3.5 \cos\left(\frac{3}{2}x\right) + 2$
17. Amplitude = 3; Period =  $\frac{3\pi}{2}$ ; Phase Shift =  $\frac{\pi}{6}$ .  
**Answer.**  $y = 3 \cos\left(\frac{4}{3}\left(x - \frac{\pi}{6}\right)\right)$
18. Amplitude = 2.5; Period =  $\frac{\pi}{4}$ ; Phase Shift =  $\frac{\pi}{4}$ .  
**Answer.**  $y = 2.5 \cos\left(6\left(x - \frac{\pi}{4}\right)\right)$
19. Amplitude = 2; Period =  $3\pi$ ; Phase Shift =  $-\frac{\pi}{2}$ ; Vertical Shift = 2.  
**Answer.**  $y = 2 \cos\left(\frac{2}{3}\left(x + \frac{\pi}{2}\right) + 2\right)$
20. Amplitude = 1.2; Period =  $\frac{5\pi}{2}$ ; Phase Shift =  $-\frac{\pi}{3}$ ; Vertical Shift = 3.  
**Answer.**  $y = 1.2 \cos\left(\frac{4}{5}\left(x + \frac{\pi}{3}\right) + 3\right)$


**Exercise Group.** For each given graph, identify the amplitude, period, phase shift, and vertical shift. Write an equation that represents these characteristics of the form  $y = A \sin(B(x - C)) + D$ .

21.




**Answer.** Amplitude:  $A = 4$ ; Period: 8; Phase Shift:  $C = 0$ ; Vertical Shift:  $D = 0$ ;  
 $y = 4 \sin\left(\frac{\pi}{4}x\right)$

22.




**Answer.** Amplitude:  $A = 3$ ; Period:  $\pi$ ; Phase Shift:  $C = 0$ ; Vertical Shift:  $D = 1$ ;  
 $y = 3 \sin(2x) + 1$

23.



**Answer.** Amplitude:  
 $A = 1.5$ ; Period:  $\pi$ ; Phase  
 Shift:  $C = \frac{\pi}{2}$ ; Vertical Shift:  
 $D = -1$ ;  
 $y = 1.5 \sin\left(2\left(x - \frac{\pi}{2}\right)\right) - 1$


24.



**Answer.** Amplitude:  $A = 4$ ;  
 Period: 12; Phase Shift:  
 $C = 3$ ; Vertical Shift:  $D = 2$ ;  
 $y = 4 \sin\left(\frac{\pi}{6}(x - 3)\right) + 2$


**Exercise Group.** For each given graph, identify the amplitude, period, phase shift, and vertical shift. Write an equation that represents these characteristics of the form  $y = A \cos(B(x - C)) + D$ .

25.




**Answer.** Amplitude:  $A = 2$ ;  
 Period:  $\frac{2\pi}{3}$ ; Phase Shift:  
 $C = 0$ ; Vertical Shift:  $D = 1$ ;  
 $y = 2 \cos(3x) + 1$

26.




**Answer.** Amplitude:  $A = 5$ ;  
 Period: 12; Phase Shift:  
 $C = 2$ ; Vertical Shift:  $D = 0$ ;  
 $y = 5 \cos\left(\frac{\pi}{6}(x - 2)\right)$

27.



**Answer.** Amplitude:  $A = 2$ ;  
 Period:  $4\pi$ ; Phase Shift:  
 $C = \pi$ ; Vertical Shift:  
 $D = -4$ ;  
 $y = 2 \cos\left(\frac{1}{2}(x - \pi)\right) - 4$

28.



**Answer.** Amplitude:  $A = 4$ ;  
 Period:  $8\pi$ ; Phase Shift:  
 $C = 2\pi$ ; Vertical Shift:  $D = 2$ ;  
 $y = 4 \cos\left(\frac{1}{4}(x - 2\pi)\right) + 2$


**Hours of daylight.** For each of the following questions, the number of daylight hours for a pair of islands in 2025 is given. These islands share the same latitude or are close to it, with one island located north of the equator and the other south of it. Find a sinusoidal function of the form  $y = A \cos(B(x - C)) + D$  to



model the daylight hours for each island. This data is sourced from the NOAA Solar Calculator.

29. Pohnpei, situated at  $6.9^\circ$  North latitude in the Federated States of Micronesia, experiences its longest day, lasting 12.52 hours, on 21 June 2025 (the 172nd day of the year), and its shortest day, lasting 11.72 hours, on 22 December 2025 (the 356th day of the year).

Nanumanga, located at  $6.3^\circ$  South latitude in Tuvalu, experiences its longest day, lasting 12.49 hours, on 22 December 2025 (the 356th day of the year), and its shortest day, lasting 11.76 hours, on 21 June 2025 (the 172nd day of the year).




**Answer.** Pohnpei:  $y = 0.4 \cos(0.0172(x - 172)) + 12.12$ ; Nanumanga:  $y = 0.365 \cos(0.0172(x - 356)) + 12.125$

30. Saipan, positioned at  $15.2^\circ$  North latitude in the Northern Mariana Islands, has its longest day, lasting 13.03 hours, on 21 June 2025 (the 172nd day of the year), and its shortest day, lasting 11.23 hours, on 22 December 2025 (the 356th day of the year).

Espiritu Santo, located at  $15.4^\circ$  South latitude in Vanuatu, experiences its longest day, lasting 13.04 hours, on 22 December 2025 (the 356th day of the year), and its shortest day, lasting 11.21 hours, on 21

June 2025 (the 172nd day of the year).




**Answer.** Saipan:  $y = 0.9 \cos(0.0172(x - 172)) + 12.13$ ; Espiritu Santo:  $y = 0.915 \cos(0.0172(x - 356)) + 12.125$

- 31.** Kaua'i, situated at  $22.1^\circ$  North latitude in Hawai'i, has its longest day, lasting 13.48 hours, on 20 June 2025 (the 171st day of the year), and its shortest day, lasting 10.78 hours, on 21 December 2025 (the 355th day of the year).

Mangai, located at  $21.9^\circ$  South latitude in the Cook Islands, experiences its longest day, lasting 13.47 hours, on 21 December 2025 (the 355th day of the year), and its shortest day, lasting 10.79 hours, on 20 June 2025 (the 171st day of the year).

It's noteworthy that Kaua'i and Mangai are situated east of the International Date Line, causing them to experience the winter and summer solstice one day earlier than islands located west of the Inter-

national Date Line.



**Answer.** Kaua'i:  $y = 1.35 \cos(0.0172(x - 171)) + 12.13$ ; Mangaia:  $y = 1.34 \cos(0.0172(x - 355)) + 12.13$

32. What patterns do you observe when comparing the graphs of daylight hours for pairs of islands across different latitudes? Additionally, how does the variation in daylight hours change as latitude moves from closer to the equator to further away?

**Answer.** Antipodal islands exhibit mirrored patterns in daylight hours, where one island experiences longer days while the other experiences shorter days. This is due to their opposite positions relative to the equator.

As latitude increases (moving away from the equator), the variation in daylight hours also increases. Islands closer to the equator experience less variation in daylight hours throughout the year, while islands further away from the equator experience more significant changes in daylight hours between seasons.

**Exercise Group.** The islands of O'ahu and Rarotonga are located at similar distances from the equator. However, they experience different climates due to their locations relative to the equator. O'ahu is situated at a latitude of 21.3 degrees north, and Rarotonga is located at a latitude of 21.2 degrees south. The table below gives the average monthly temperatures (in °C) for each island. Use the data in the table to answer the following questions. Source: <http://www.worldclimate.com>, retrieved on 18 March, 2024.

Month ( $x$ )	O'ahu (21.3° N)	Rarotonga (21.2° S)
January (1)	22.7	25.7
February (2)	22.7	26.1
March (3)	23.5	25.8
April (4)	24.3	24.9
May (5)	25.2	23.5
June (6)	26.3	22.3
July (7)	26.9	21.7
August (8)	27.4	21.6
September (9)	27.2	22.1
October (10)	26.4	22.9
November (11)	25.1	23.8
December (12)	23.3	24.8

- 33.** Determine a sinusoidal function of the form  $y = A \sin(B(x - C)) + D$  to represent the average monthly temperatures provided in the table for each island.

**Answer.** O'ahu:  $y = 2.35 \sin(0.5236(x - 5)) + 25.05$ ; Rarotonga:  $y = 2.25 \sin(0.5236(x - 11)) + 23.85$

- 34.** Utilize a graphing utility to identify the best-fit sinusoidal function of the form  $y = A \sin(B(x - C)) + D$  for each island.

**Answer.** O'ahu:  $y = 2.3727 \sin(0.502443(x - 4.72583)) + 25.005$ ; Rarotonga:  $y = 2.24202 \sin(0.521233(x + 1.16396)) + 23.7744$

- 35.** How do the temperature patterns of O'ahu, situated in the northern hemisphere, compare with those of Rarotonga, positioned in the southern hemisphere?

**Answer.** During O'ahu's winter, Rarotonga experiences its summer, and similarly, during O'ahu's summer, Rarotonga experiences its winter.

**Exercise Group.** Use the table below, which gives the average monthly temperatures (in Celsius) at various latitudes in the South Pacific, to answer the following questions. Express your answers in the form  $y = A \sin(B(x - C)) + D$ . Source: Data for Apia, Suva, Nuku'alofa, and Rapa Nui obtained from <http://www.worldclimate.com>, retrieved on 18 March 2024; Data for Whangārei and Dunedin obtained from [National Institute of Water and Atmospheric Research \(NIWA\)](http://www.niwa.co.nz)<sup>2</sup>, retrieved on 18 March 2024.

<sup>2</sup>NationalInstituteofWaterandAtmosphericResearch(NIWA)

Month	Apia 13.8°S	Suva 18.1°S	Nuku'alofa 21.1°S	Rapa Nui 27.1°S	Whangārei 35.7°S	Dunedin 45.9°S
Jan (1)	27.6	26.7	25.6	23.3	19.9	15.3
Feb (2)	27.6	26.9	26	23.7	20.2	15
Mar (3)	27.8	26.7	25.8	23.1	18.8	13.7
Apr (4)	27.8	26	24.9	21.9	16.6	11.7
May (5)	27.4	24.8	23.1	20.1	14.4	9.3
Jun (6)	27.1	24	22.4	18.9	12.4	7.3
Jul (7)	26.7	23.2	21.3	18	11.6	6.6
Aug (8)	26.5	23.2	21.2	17.9	11.9	7.7
Sep (9)	26.7	23.7	21.7	18.3	13.3	9.5
Oct (10)	27	24.4	22.4	19	14.6	10.9
Nov (11)	27.4	25.2	23.5	20.4	16.4	12.4
Dec (12)	27.4	26.1	24.7	21.8	18.5	13.9

**36.** Determine a sinusoidal function to represent the average monthly temperature in

(a) Apia, Sāmoa (13.8°S)

**Answer.**  $y = 0.65 \sin(0.5236(x - 11)) + 27.15$

(b) Suva, Fiji (8.1°S)

**Answer.**  $y = 1.85 \sin(0.5236(x - 11)) + 25.05$

(c) Nuku'alofa, Tonga (21.1°S)

**Answer.**  $y = 2.4 \sin(0.5236(x - 11)) + 23.6$

(d) Rapa Nui (27.1°S)

**Answer.**  $y = 2.9 \sin(0.5236(x - 11)) + 20.8$

(e) Whangārei, New Zealand (35.7°S)

**Answer.**  $y = 4.3 \sin(0.5236(x - 11)) + 15.9$

(f) Dunedin, New Zealand (45.9°S)

**Answer.**  $y = 4.35 \sin(0.5236(x - 11)) + 10.95$

**37.** Utilize a graphing utility to identify the best-fit sinusoidal function in

(a) Apia, Sāmoa (13.8°S)

**Answer.**  $y = 0.599155 \sin(0.590429(x - 0.197373)) + 27.2133$

(b) Suva, Fiji (8.1°S)

**Answer.**  $y = 1.86982 \sin(0.532431(x + 1.0423)) + 25.0509$

(c) Nuku'alofa, Tonga (21.1°S)

**Answer.**  $y = 2.41339 \sin(0.521333(x + 1.07987)) + 23.5577$

(d) Rapa Nui (27.1°S)

**Answer.**  $y = 2.93283 \sin(0.494705(x + 1.55737)) + 20.6621$

(e) Whangārei, New Zealand (35.7°S)

**Answer.**  $y = 4.27205 \sin(0.503071(x + 1.88988)) + 15.8716$

(f) Dunedin, New Zealand (45.9°S)

**Answer.**  $y = 4.07644 \sin(0.524652(x + 1.8183)) + 11.1006$

- 38.** Based on the sinusoidal functions that you found, what can you conclude about the relationship between temperature and latitude?

**Answer.** As latitude increases (moving away from the equator towards the poles), the sinusoidal function's amplitude increases, signifying greater temperature variability, while its vertical shift decreases, indicating lower average temperatures at higher latitudes.

## 2.4 Inverse Trigonometric Functions

Swells are a crucial navigational tool, providing a consistent means of maintaining a straight course over extended periods. Unlike stars, which may be obscured during the day or on cloudy nights, or winds that can change direction frequently, swells tend to remain relatively constant. Crew members can navigate a straight course by keeping the angle at which the swell passes the canoe constant.


Even in poor visibility, when the crew may not see the swells, they can feel them. Parts of the vaka (canoe) lift and lower as the swells pass beneath. Skilled navigators use these movements to maintain course, even without directly seeing the waves.

The side-to-side rocking motion of the canoe is known as **roll**. This motion occurs when a swell approaches from either the port (left) or starboard (right) side, causing the vaka to initially lift the corresponding hull (port or starboard), followed by the opposite hull, depending on the direction of the swell.

When a swell approaches from the bow (front) of the canoe, it lifts the front, causing the canoe to tilt backward before tilting forward. This motion is known as **pitch**. Conversely, if the swell comes from the stern (back), the canoe tilts forward and then backward.

When a swell approaches the canoe from an angle that is not perpendicular to any side, a combination of pitch and roll occurs. The specific motion experienced depends on the precise angle of the swell. For instance, if the swell comes from an angle and lifts the starboard bow (front right), it may subsequently lift the port bow, followed by the starboard stern, and finally the port stern. However, with a slightly different angle, the sequence may change as well. After lifting the starboard bow, it could lead to the starboard stern being lifted next, followed by the port bow, and finally the port stern. This twisting motion is referred to as a **corkscrew** effect due to its combination of motions.

The video in [Figure 2.4.1](#) demonstrates how the vaka Paikea moves as swells pass under the hulls. A change in vaka motion can indicate either a change in the canoe's direction or a shift in the direction of ocean swells. In such cases, the crew must assess the situation and, when conditions allow, utilize celestial markers, such as the rising and setting of stars, to determine the direction of the swells.




**Figure 2.4.1** As swells pass under the vaka Paikea, the canoe pitches, rolls, and corkscrews, depending on the angle of the swell. A navigator can use these movements to keep a straight course.

As the vaka Paikea sails north in the Cook Islands from Rarotonga to Aitutaki with a heading of  $0^\circ$ , the swells are approaching the vaka from the southwest and moving towards the northeast. Referring to [Figure 1.2.17](#), which

provides heading angles, we observe that the swells have a heading ranging between  $0^\circ$  (north) and  $90^\circ$  (east).


Additional observations of the crew reveal the swells hit Paikea in the following order: 1) port stern (back left); 2) starboard stern (back right); 3) port bow (front left); and 4) starboard bow (front right), as shown in [Figure 2.4.2](#). If Paikea is 14.8 m (50 ft) in length and 6.2 m (20 ft) in width, what is the possible range of headings from which the swells may be approaching?



**Figure 2.4.2** Swells moving towards the northeast pass under the vaka Paikea heading.

To determine the range of headings, we start by considering one boundary, which occurs when the swells are moving directly north with a heading of  $0^\circ$ . To identify the other boundary, we want to find the angle at which the swell intersects both the starboard stern and port bow simultaneously as it passes beneath the vaka. Essentially, we want to find the angle that diagonally traverses the canoe from one corner to another. To simplify this, we represent the vaka as a rectangle and create a triangle by connecting the corners. The angle, denoted as  $\theta$ , is formed between the side adjacent to the 14.8 m length of the canoe and the diagonal. This angle corresponds to the heading of the swells, as shown in [Figure 2.4.3](#).





**Figure 2.4.3** The deck of the vaka Paikea can be simplified to a rectangle, and its diagonal represents the threshold between different sequences of motions as swells pass. The angle  $\theta$  corresponds to the heading of the swells.

You may notice that for our given angle,  $\theta$ , we have the opposite side of length 6.2 m and the adjacent side of length 14.8 m, which are both related to the tangent function:

$$\tan \theta = \frac{14.8}{6.2}$$

Until now, the angles for the trigonometric problems we encountered were given. However, to solve for the angle  $\theta$ , we cannot simply divide by “tan” since it is part of the tangent function, which takes an angle as input and provides a ratio of sides as output. To find the angle  $\theta$ , we need to use the inverse function, which takes the ratio of sides as input and provides an angle as output. In this section, we will explore inverse trigonometric functions, including their properties and usage.

### 2.4.1 Inverse Trigonometric Functions


Recall from algebra that for a function  $f$  and its inverse function  $f^{-1}$  we have

1. The Domain of  $f^{-1} = \text{Range of } f$
2. The Range of  $f^{-1} = \text{Domain of } f$
3. If  $f(a) = b$  then  $f^{-1}(b) = a$

In terms of trigonometric functions, for example, if  $f(x) = \sin x$  then  $f^{-1}(x) = \sin^{-1} x$ . Now consider  $\sin\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2}$ , then  $\frac{\pi}{4} = \sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$ .


**Remark 2.4.4 Be Careful.** Do not confuse the inverse trigonometric notations with an exponent, in other words,  $\sin^{-1} x \neq \frac{1}{\sin x}$ . To avoid this, we will use parentheses around the trigonometric function to denote the power of negative one:  $(\sin x)^{-1}$ .

Also recall from algebra that for a function,  $f$ , to have an inverse,  $f^{-1}$ , it must be one-to-one, meaning no horizontal line intersects the graph more than once. Since this is not true for trigonometric functions, they do not have inverses. We need functions to be one-to-one because both  $\sin \frac{\pi}{4} = \frac{\sqrt{2}}{2}$  and  $\sin \frac{3\pi}{4} = \frac{\sqrt{2}}{2}$  but if we took the inverse of sine, we wouldn't know to use  $\sin^{-1} \left( \frac{\sqrt{2}}{2} \right) = \frac{\pi}{4}$  or  $\sin^{-1} \left( \frac{\sqrt{2}}{2} \right) = \frac{3\pi}{4}$ . To avoid this confusion and to ensure the function is one-to-one, we can put restrictions on the domains of each trigonometric function so they attain all the values in the range only once, making the function one-to-one, and thus have an inverse (see Figure 2.4.5).




**Figure 2.4.5** The graph of  $f(x) = \sin x$  does not pass the horizontal line test and is not one-to-one. If we restrict the graph to  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$  so that each value in the range  $[-1, 1]$  is attained only once, then the function is one-to-one and has an inverse.


We will restrict the domain of  $y = \sin x$  to the interval  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ , the domain of  $y = \cos x$  to  $[0, \pi]$ , and the domain of  $y = \tan x$  to the interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$ . Notice that the domain for each trigonometric function includes one quadrant where the function is positive and one quadrant where it is negative. The domains and the corresponding graphs for sine, cosine, and tangent are shown in Figure 2.4.6, Figure 2.4.7, and Figure 2.4.8, respectively.



**Figure 2.4.6** The domain of  $y = \sin x$  (left) and its graph on the restricted domain (right).



**Figure 2.4.7** The domain of  $y = \cos x$  (left) and its graph on the restricted domain (right).



**Figure 2.4.8** The domain of  $y = \tan x$  (left) and its graph on the restricted domain (right).

With these new restrictions on the domains, we now have trigonometric functions that are one-to-one and so we can define their inverse functions:

**Definition 2.4.9 Inverse Sine.** The **inverse sine function** is symbolized by

$$y = \sin^{-1} x \quad \text{and means} \quad x = \sin y.$$

The inverse sine function is also called the **arcsine function**, and is denoted by  $\arcsin x$ .  $\diamond$

**Definition 2.4.10 Inverse Cosine.** The **inverse cosine function** is symbolized by

$$y = \cos^{-1} x \quad \text{and means} \quad x = \cos y.$$

The inverse cosine function is also called the **arccosine function**, and is denoted by  $\arccos x$ .  $\diamond$

**Definition 2.4.11 Inverse Tangent.** The *inverse tangent function* is symbolized by

$$y = \tan^{-1} x \quad \text{and means} \quad x = \tan y.$$

The inverse tangent function is also called the **arctangent function**, and is denoted by  $\arctan x$ .  $\diamond$

**Definition 2.4.12 Inverse Cosecant, Secant, and Cotangent.** Inverse cosecant, inverse secant, and inverse cotangent functions are not as common as the other trigonometric functions and we will just summarize them.

$$\begin{array}{lll} y = \csc^{-1} x & \text{means} & x = \csc y \\ y = \sec^{-1} x & \text{means} & x = \sec y \\ y = \cot^{-1} x & \text{means} & x = \cot y \end{array}$$

◇

**Definition 2.4.13 Domain and Range for Inverse Trigonometric Functions.** The domain and range for each function is

Function	Domain	Range
$\sin^{-1}$	$[-1, 1]$	$[-\frac{\pi}{2}, \frac{\pi}{2}]$
$\cos^{-1}$	$[-1, 1]$	$[0, \pi]$
$\tan^{-1}$	$[-\infty, \infty]$	$(-\frac{\pi}{2}, \frac{\pi}{2})$
$\csc^{-1}$	$(-\infty, -1] \cup [1, \infty)$	$[-\frac{\pi}{2}, \frac{\pi}{2}], y \neq 0$
$\sec^{-1}$	$(-\infty, -1] \cup [1, \infty)$	$[0, \pi], y \neq \frac{\pi}{2}$
$\cot^{-1}$	$(-\infty, \infty)$	$(0, \pi)$

◇

### 2.4.2 Finding the Exact Value of an Inverse Trigonometric Function

**Example 2.4.14 Evaluating Inverse Trigonometric Functions.** Find the exact value

(a)  $\cos^{-1}(\frac{1}{2})$

**Solution.** Let  $\theta = \cos^{-1}(\frac{1}{2})$ . Then evaluating the problem is the same as determining the angle,  $\theta$ , for which

$$\cos \theta = \frac{1}{2}.$$

Although there are infinite values of  $\theta$  that satisfy the equation, such as  $\theta = \frac{\pi}{3}$  and  $\theta = \frac{5\pi}{3}$ , there is only one value that lies in the interval  $[0, \pi]$ . Thus,  $\cos^{-1}(\frac{1}{2}) = \frac{\pi}{3}$ .

(b)  $\tan^{-1} \sqrt{3}$

**Solution.** Let  $\theta = \tan^{-1} \sqrt{3}$ . Then we must find  $\theta$  that satisfies  $\tan \theta = \sqrt{3}$  as well as satisfies the range of  $\tan^{-1}$ . Because  $\tan \frac{\pi}{3} = \sqrt{3}$  and  $-\frac{\pi}{2} < \frac{\pi}{3} < \frac{\pi}{2}$ , we conclude that  $\tan^{-1} \sqrt{3} = \frac{\pi}{3}$ .

(c)  $\sin^{-1}(-\frac{\sqrt{2}}{2})$

**Solution.** The angle,  $\theta$ , in the interval  $[-\frac{\pi}{2}, \frac{\pi}{2}]$  that satisfies  $\sin \theta = -\frac{\sqrt{2}}{2}$  is  $\theta = \sin^{-1}(-\frac{\sqrt{2}}{2}) = -\frac{\pi}{4}$ .

□

### 2.4.3 Approximations of Inverse Trigonometric Functions

To evaluate inverse trigonometric functions that do not have special angles, we will need to use a calculator.

**Remark 2.4.15 Using a Calculator.** Using a calculator for inverse trigonometric functions: Most calculators will have a special key for the inverse sine, cosine, and tangent functions. Depending on your calculator, you may see the following keys for inverse the inverse trigonometric functions

Function	Calculator Key
inverse sine	$\boxed{\text{SIN}^{-1}}$ , $\boxed{\text{INV}}\boxed{\text{SIN}}$ , $\boxed{\text{ARCSIN}}$ , or $\boxed{\text{ASIN}}$
inverse cosine	$\boxed{\text{COS}^{-1}}$ , $\boxed{\text{INV}}\boxed{\text{COS}}$ , $\boxed{\text{ARCCOS}}$ , or $\boxed{\text{ACOS}}$
inverse tangent	$\boxed{\text{TAN}^{-1}}$ , $\boxed{\text{INV}}\boxed{\text{TAN}}$ , $\boxed{\text{ARCTAN}}$ , or $\boxed{\text{ATAN}}$

Often, the inverse trigonometry key can be found by first pressing  $\boxed{2\text{nd}}$  or  $\boxed{\text{SHIFT}}$ , followed by the trigonometry function key. For example, to get the  $\boxed{\text{SIN}^{-1}}$  key, press  $\boxed{2\text{nd}}\boxed{\text{SIN}}$  or  $\boxed{\text{SHIFT}}\boxed{\text{SIN}}$ .

Many calculators do not have specific keys for the inverse cosecant, secant, and cotangent functions. Instead, you can use the Reciprocal Identities (Definition 1.4.2) to get

Function	Calculator Key
inverse cosecant	$\boxed{1} / \boxed{\text{SIN}^{-1}}$
inverse secant	$\boxed{1} / \boxed{\text{COS}^{-1}}$
inverse cotangent	$\boxed{1} / \boxed{\text{TAN}^{-1}}$

**Example 2.4.16 Finding an Approximate Value of Inverse Trigonometric Functions.** Use a calculator to approximate the value of each expression in radians, rounded to two decimals.

(a)  $\cos^{-1}(-0.39)$

**Solution.** First verify the mode of the calculator is in radians. Then, press the following keys  $\boxed{\text{COS}^{-1}}\boxed{(}\boxed{-}\boxed{0.39}\boxed{)}\boxed{\text{ENTER}}$  to get

$$\cos^{-1}(-0.39) \approx 1.9714279195$$

If your calculator does not have the  $\boxed{\text{COS}^{-1}}$  key, then use the appropriate key(s) for inverse cosine, such as  $\boxed{\text{INV}}\boxed{\text{COS}}$ ,  $\boxed{\text{ARCCOS}}$ , or  $\boxed{\text{ACOS}}$ .

On some calculators,  $\boxed{\text{COS}^{-1}}$  is pressed first, then  $\boxed{(}\boxed{-}\boxed{0.39}\boxed{)}\boxed{\text{ENTER}}$ ; while other calculators the sequence is reversed with  $\boxed{(}\boxed{-}\boxed{0.39}\boxed{)}\boxed{\text{ENTER}}$  pressed first, then  $\boxed{\text{COS}^{-1}}$ . Verify with your calculator's manual.

(b)  $\tan^{-1} 12$

**Solution.** First verify the mode of the calculator is in radians. Then, using the appropriate key for inverse tangent, press the following keys  $\boxed{\text{TAN}^{-1}}\boxed{(}\boxed{12}\boxed{)}\boxed{\text{ENTER}}$  to get

$$\tan^{-1} 12 \approx 1.48765509491$$

(c)  $\sin^{-1} 0.8$

**Solution.** First verify the mode of the calculator is in radians. Then, using the appropriate key for inverse sine, press the following keys  $\boxed{\text{SIN}^{-1}}\boxed{(}\boxed{0.8}\boxed{)}\boxed{\text{ENTER}}$  to get

$$\sin^{-1} 0.8 \approx 0.927295218002$$

□

**Example 2.4.17 Swells Approaching Paikea.** At the start of this section, we discussed the swells passing through Paikea in the following order: 1) port stern (back left); 2) starboard stern (back right); 3) port bow (front left); and 4) starboard bow (front right). Given that Paikea is 14.8 m (50 ft) in length and 6.2 m (20 ft) in width, what is the possible range of headings from which the swells may be approaching?

**Solution.** If the swells are moving directly north, the heading will be  $0^\circ$ .

To find the other boundary, we want to determine the angle at which the swell simultaneously intersects both the starboard stern and port bow as it passes beneath the vaka. This angle, denoted as  $\theta$ , is formed between the side adjacent to the 14.8 m length of the canoe and the diagonal. Using the tangent function we have:

$$\tan \theta = \frac{14.8}{6.2}$$

Solving for  $\theta$  using the inverse tangent function we get:


$$\tan^{-1} \left( \frac{14.8}{6.2} \right) \approx 67.27^\circ$$

Therefore, the possible range of headings from which the swells may be approaching is between  $0^\circ$  (directly north) and approximately  $67.27^\circ$ .  $\square$

**Example 2.4.18 A gift for Mau.** In 1999, after Mau Pialug, also known as Papa Mau, sailed from Hawai'i to his home in Satawal, Micronesia aboard the Makali'i, the crew of Nā Kālai Wa'a expressed their gratitude to the man who shared his knowledge of navigation by constructing a sister canoe to Makali'i. This vessel, named the Alingano Maisu, was a 56-foot long double-hulled voyaging canoe. In 2007, accompanied by the Hōkūle'a, the Alingano Maisu embarked on its inaugural journey to Satawal, continuing Papa Mau's legacy of navigation in his home islands.

During this journey, the canoes sailed directly towards Johnston Atoll, using it as a sighting point without making a stop, before proceeding to their first destination in Majuro, Marshall Islands. The island of Majuro is situated 1,108 nautical miles west and 575 nautical miles south of Johnston Atoll.

What house do you need to sail in and what distance will you need to sail? If the wa'a travels at 5 knots, how many days will it take to reach the destination? Note that 1 knot = 1 nautical mile/hour.



We will use the tangent function because we are given the side opposite to  $\theta$  (575 NM) and the side adjacent to  $\theta$  (1,100 NM) and express it as

$$\tan \theta = \frac{575}{1,100}$$

**Solution.** Since we were able to write  $\tan \theta = \frac{575}{1,108}$ , we can use a calculator or other technology to evaluate the inverse tangent to find our angle:

$$\theta = \tan^{-1} \left( \frac{575}{1,108} \right) \approx 27.4^\circ.$$

Because this angle is in Quadrant III, we find its value on the Unit Circle by adding  $180^\circ + 27.4^\circ = 207.4^\circ$ . Next we refer to the Star Compass with angles (Figure 1.2.4) to conclude we will need to sail towards the House ‘Aina Kona.

To determine the distance,  $d$ , we use the Pythagorean Theorem:


$$d = \sqrt{575^2 + 1,108^2} \approx 1,248\text{NM}.$$

Finally, we note that since  $(\text{speed}) = (\text{distance})/(\text{duration})$ , we can rearrange the terms to get  $(\text{duration}) = (\text{distance})/(\text{speed})$ . If we travel at 5 knots (5 NM/hr), we can calculate the duration as

$$\begin{aligned} \text{duration} &= \frac{1,248.3 \text{ NM}}{5 \text{ kt}} && \frac{\text{distance}}{\text{speed}} \\ &= \frac{1,248.3 \text{ NM}}{5 \text{ NM/hr}} && 1 \text{ kt} = 1 \text{ NM/hr} \\ &= 249.7 \text{ hours} && \frac{\text{NM}}{\text{NM/hr}} = \text{NM} \cdot \frac{\text{hr}}{\text{NM}} = \text{hr} \\ &= 249.7 \text{ hours} \cdot \frac{\text{day}}{24 \text{ hours}} \\ &\approx 10.4 \text{ days.} \end{aligned}$$

□

**Example 2.4.19 Finding Land.** As Hōkūle‘a is sailing towards Rapa Nui, the navigator uses a process called **dead reckoning** to determine their position based on the latitude, measured by the stars, and other factors such as the estimated distance traveled, speed, and direction. Once the navigator has determined they are in the vicinity of land, her attention is now focused on looking for signs of land. One method navigators will use is to look for land-based seabirds such as the manu-o-kū (fairy tern) and the noio (noddy tern), which go out to sea in the morning to fish and return to land at night. However, Rapa Nui’s seabird population has been reduced so she will look for other signs such as drifting land vegetation; clouds that form over islands; the loom of the island when white sand and still lagoons reflect the sun or moon upwards; and distinctive patterns of swells bending (refracting) around and/or reflecting off islands. Land will be spotted when the navigator first sees Maunga Terevaka, the tallest point in Rapa Nui, which stands at 1,665 ft. Figure 2.4.20 depicts the relation between the canoe and island (left) as well as what is seen from the deck of the canoe (right).




**Figure 2.4.20** Watch as the canoe approaches an island from two different views. On the left, see a side view as the island gradually comes into sight over the horizon. On the right, experience the perspective from the canoe’s deck, observing the island appearing to rise from the water. This dual perspective provides a unique glimpse into the legend of Maui, the demigod who pulled the islands from the ocean with his fish-hook.

1. On the deck of Hōkūle‘a, a navigator stands at 9 ft above sea level. If she looks out to the sea, how far is she from horizon? Assume the radius of the earth at Rapa Nui is 20,911,171 feet.
2. How far is Hōkūle‘a from Maunga Terevaka when it first becomes visible over the horizon to someone standing 9ft above sea level?


**Solution.**

1. We begin by assuming that the earth is a sphere with radius  $R$ . Standing on Hōkūle‘a, the line from the navigator’s eye to the horizon is tangent to the circle of radius  $R$ .



Here  $h = 9$  ft, represents the height of the navigator’s eye above sea level,  $R = 20,911,171$  ft is the radius of the earth at Rapa Nui, and  $s_1$  is the arc length or distance along the surface of the earth from the navigator to the horizon. We have written the distances in feet since the height of the navigator is in feet. Recall from [Theorem 1.2.25](#) the formula for finding the arc length is  $s_1 = 2\pi R \cdot \left(\frac{\theta_1}{360}\right)$ . Since we know  $R$ , we only need to find  $\theta_1$ . Notice this forms a right triangle.





Since we know the adjacent side and hypotenuse of this triangle, we can use cosine:

$$\cos \theta_1 = \frac{R}{R+h} = \frac{20,911,171}{20,911,171+9} = \frac{20,911,171}{20,911,180}.$$

To solve for  $\theta_1$ , we use the inverse cosine

$$\theta_1 = \cos^{-1} \left( \frac{20,911,171}{20,911,180} \right) \approx 0.053^\circ.$$

Now we are ready to calculate the arc length,  $s_1$ :


$$s_1 = 2\pi \cdot \frac{0.053^\circ}{360^\circ} \cdot 20,911,171 \text{ feet} \approx 19,343 \text{ feet}.$$

Converting to miles we get


$$s_1 \approx 19,401 \text{ feet} \cdot \frac{1 \text{ mile}}{5,280 \text{ feet}} \approx 3.7 \text{ miles}.$$

So the horizon is 3.7 miles from the navigator.

- Next, to determine how far the navigator is from Maunga Terevaka when it emerges over the horizon, we need to align the top of the mountain with line from the navigator's eye to the horizon.



Here,  $H = 1,665$  ft, is the height of Maunga Terevaka. To find the distance from the top of the mountain to the horizon, we will need to determine  $s_2$ . We begin by redrawing the triangle.



The angle is then given by

$$\begin{aligned}
 \theta_1 &= \cos^{-1} \left( \frac{R}{R+H} \right) \\
 &= \cos^{-1} \left( \frac{20,911,171}{20,911,173 + 1,665} \right) \\
 &= \cos^{-1} \left( \frac{20,911,171}{20,912,836} \right) \\
 &\approx 0.723^\circ
 \end{aligned}$$

We conclude that the distance from Maunga Terevaka to the horizon is

$$s_2 = 2\pi \cdot \frac{0.723^\circ}{360^\circ} \cdot (20,911,171 \text{ feet}) \cdot \frac{1 \text{ mile}}{5,280 \text{ feet}} \approx 50.0 \text{ miles}$$

Therefore, the total distance between the navigator and Maunga Terevaka is  $s_1 + s_2 = 3.7 + 50.0 = 53.7$  miles. Please note that this is the distance when the island may first be seen, however, weather conditions may reduce visibility.



□


#### 2.4.4 Graphs of Inverse Trigonometric Functions

Recall from algebra that

1. The point  $(a, b)$  is on the graph of  $f$  if and only if the point  $(b, a)$  is on the graph of  $f^{-1}$ .
2. The graphs of  $f^{-1}$  and  $f$  are reflections of each other about the line  $y = x$

The graph of each inverse trigonometric function can be obtained by reflecting the graph of the original function about the line  $y = x$ .





### 2.4.5 Composition of Inverse Trigonometric Functions

Recall Algebra that if  $f$  is a one-to-one function with inverse  $f^{-1}$ , then

1.  $f(f^{-1}(y)) = y$  for every  $y$  in the domain of  $f^{-1}$
2.  $f^{-1}(f(x)) = x$  for every  $x$  in the domain of  $f$

In terms of trigonometric functions,  $f(f^{-1}(y)) = y$  will work for all  $y$  in the domain, however, we need to be careful when evaluating  $f^{-1}(f(x)) = x$  because the domain of  $f^{-1}$  is restricted.

**Definition 2.4.21** Properties of Composition of Trigonometric Functions.

$$\begin{array}{lll}
 \sin(\sin^{-1} x) = x & \text{when} & -1 \leq x \leq 1 \\
 \cos(\cos^{-1} x) = x & \text{when} & -1 \leq x \leq 1 \\
 \tan(\tan^{-1} x) = x & \text{when} & -\infty \leq x \leq \infty \\
 \sin^{-1}(\sin x) = x & \text{only when} & -\frac{\pi}{2} \leq x \leq \frac{\pi}{2} \\
 \cos^{-1}(\cos x) = x & \text{only when} & 0 \leq x \leq \pi \\
 \tan^{-1}(\tan x) = x & \text{only when} & -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}
 \end{array}$$

◇

**Example 2.4.22** Composition of trigonometric function and the inverse of the same trigonometric function. Find the exact value of each expression

(a)  $\cos^{-1}\left(\cos \frac{\pi}{8}\right)$

**Solution.** Since  $\frac{\pi}{8}$  is in the interval  $[0, \pi]$ , then from the properties of compositions of inverse functions, we get

$$\cos^{-1}\left(\cos \frac{\pi}{8}\right) = \frac{\pi}{8}$$

(b)  $\tan(\tan^{-1} 7)$

**Solution.** Since 7 is in the interval  $(-\infty, \infty)$ ,

$$\tan(\tan^{-1} 7) = 7$$


(c)  $\sin^{-1}(\sin(-\frac{\pi}{7}))$

**Solution.** Since  $-\frac{\pi}{7}$  is in the interval  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ ,

$$\sin^{-1}(\sin(-\frac{\pi}{7})) = -\frac{\pi}{7}$$

(d)  $\sin^{-1}(\sin \frac{4\pi}{5})$

**Solution.** Note that  $\frac{5\pi}{7}$  is not in the interval  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ . In order to evaluate the expression, we first need to find an angle,  $\theta$ , such that  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$  and  $\sin \frac{5\pi}{7} = \sin \theta$ . Since  $\frac{\pi}{2} < \frac{5\pi}{7} < \pi$  means  $\frac{5\pi}{7}$  is in Quadrant II. Recall from Section 1.3 that our reference angle is  $\theta = \pi - \frac{5\pi}{7} = \frac{2\pi}{7}$ .



Since  $\frac{2\pi}{7}$  is in the interval  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ ,

$$\sin^{-1}(\sin \frac{5\pi}{7}) = \sin^{-1}(\sin \frac{2\pi}{7}) = \frac{2\pi}{7}$$

(e)  $\cos(\cos^{-1}(-0.283))$

**Solution.** Since  $-0.283$  is in the interval  $[-1, 1]$ ,

$$\cos(\cos^{-1}(-0.283)) = -0.283$$

□

Now we will look at what happens when we need to evaluate the composition of a trigonometric function and the inverse of a different trigonometric function.


**Example 2.4.23 Composition of trigonometric function and the inverse of a different trigonometric function.** Find the exact value of

(a)  $\cos(\tan^{-1}(\frac{4}{3}))$

**Solution.** Let  $\theta$  be an angle in the range of  $\tan^{-1}$ , that is, let  $\theta$  be in the interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$  such that  $\theta = \tan^{-1}(\frac{4}{3})$ . This equation is equivalent to  $\tan \theta = \frac{4}{3}$ . Since  $\tan \theta > 0$ , we know that  $\theta$  must be in Quadrant I ( $0 < \theta < \frac{\pi}{2}$ ). Let  $(x, y)$  be a point on the terminal side of  $\theta$ , then by the trigonometric ratios,

$$\tan \theta = \frac{\text{opposite}}{\text{adjacent}} = \frac{y}{x}$$

Since we have  $\tan \theta = \frac{4}{3}$ , we get  $x = 3$  and  $y = 4$ , as shown in the figure below.



Evaluating  $\cos(\tan^{-1}(\frac{4}{3}))$  is equivalent to evaluating

$$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}} = \frac{3}{r},$$

where  $r = \sqrt{x^2 + y^2} = \sqrt{3^2 + 4^2} = \sqrt{9 + 16} = \sqrt{25} = 5$ . So,


$$\cos\left(\tan^{-1}\frac{4}{3}\right) = \cos \theta = \frac{3}{5}$$

(b)  $\sin(\cos^{-1}(-\frac{3}{8}))$

**Solution.** Let  $\theta$  be an angle in the range of  $\cos^{-1}$ , that is, let  $\theta$  be in the interval  $[0, \pi]$  such that  $\theta = \cos^{-1}(-\frac{3}{8})$ . This equation is equivalent to  $\cos \theta = -\frac{3}{8}$ . Since  $\cos \theta < 0$ , we know that  $\theta$  must be in Quadrant II ( $\frac{\pi}{2} \leq \theta \leq \pi$ ). Let  $(x, y)$  be a point on the terminal side of  $\theta$ , then by the trigonometric ratios,

$$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}} = \frac{x}{r}$$

Since we have  $\cos \theta = -\frac{3}{8}$ , we get  $\frac{x}{r} = -\frac{3}{8}$  which gives us either  $x = -3$  and  $r = 8$  or  $x = 3$  and  $r = -8$ . Since  $\theta$  is in Quadrant II, we know that  $x$  is negative, thus  $x = -3$  and  $r = 8$ , as shown in the figure below.



Evaluating  $\sin(\cos^{-1}(-\frac{3}{8}))$  is equivalent to evaluating

$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}} = \frac{y}{8},$$

where

$$r^2 = x^2 + y^2$$

$$8^2 = (-3)^2 + y^2$$

$$64 = 9 + y^2$$



$$y^2 = 55$$

Thus  $y = \sqrt{55}$ . So,

$$\sin\left(\cos^{-1}\frac{-3}{8}\right) = \sin\theta = \frac{\sqrt{55}}{8}$$

Note this is positive since sine is positive in Quadrant II.

□

### 2.4.6 Exercises

**Exercise Group.** What are the domain and range of

- |  |   |
|--|---|
| 1. $y = \sin x$<br><b>Answer.</b> D: $-\infty < x < \infty$ ;<br>R: $-1 \leq y \leq 1$ | 2. $y = \sin^{-1} x$<br><b>Answer.</b> D: $-1 \leq y \leq 1$ ; R:<br>$-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ |
| 3. $y = \cos x$<br><b>Answer.</b> D: $-\infty < x < \infty$ ;<br>R: $-1 \leq y \leq 1$ | 4. $y = \cos^{-1} x$<br><b>Answer.</b> D: $-1 \leq y \leq 1$ ; R:<br>$-0 \leq y \leq \pi$                       |

**Exercise Group.** Determine the exact value of each expression, expressed in radians.


- |  |  |   |
|--|--|---|
| 5. $\tan^{-1}(0)$<br><b>Answer.</b> 0  | 6. $\tan^{-1}(-\sqrt{3})$<br><b>Answer.</b> $-\frac{\pi}{3}$                     | 7. $\cos^{-1}\left(-\frac{\sqrt{2}}{2}\right)$<br><b>Answer.</b> $\frac{3\pi}{4}$ |
| 8. $\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)$<br><b>Answer.</b> $\frac{\pi}{4}$    | 9. $\tan^{-1}(-1)$<br><b>Answer.</b> $-\frac{\pi}{4}$                            | 10. $\sin^{-1}\left(\frac{\sqrt{3}}{2}\right)$<br><b>Answer.</b> $\frac{\pi}{3}$  |
| 11. $\cos^{-1}(-1)$<br><b>Answer.</b> $\pi$  | 12. $\cos^{-1}\left(\frac{\sqrt{2}}{2}\right)$<br><b>Answer.</b> $\frac{\pi}{4}$ | 13. $\sin^{-1}(1)$<br><b>Answer.</b> $\frac{\pi}{2}$                              |
| 14. $\sin^{-1}\left(-\frac{\sqrt{3}}{2}\right)$<br><b>Answer.</b> $-\frac{\pi}{3}$ | 15. $\cos^{-1}\left(-\frac{1}{2}\right)$<br><b>Answer.</b> $\frac{2\pi}{3}$      | 16. $\tan^{-1}\left(\frac{\sqrt{3}}{3}\right)$<br><b>Answer.</b> $\frac{\pi}{6}$  |

**Exercise Group.** Use a calculator to approximate each expression. Provide your answer in radians, rounding to two decimal places.

- |   |  |  |
|---|--|--|
| 17. $\cos^{-1}\left(\frac{\sqrt{5}}{3}\right)$<br><b>Answer.</b> 0.73 | 18. $\sin^{-1}(0.63)$<br><b>Answer.</b> 0.68                   | 19. $\tan^{-1}(\sqrt{7})$<br><b>Answer.</b> 1.21 |
| 20. $\sin^{-1}\left(\frac{\sqrt{2}}{9}\right)$<br><b>Answer.</b> 0.16 | 21. $\tan^{-1}(0.7)$<br><b>Answer.</b> 0.61                    | 22. $\cos^{-1}(-0.9)$<br><b>Answer.</b> 2.69     |
| 23. $\sin^{-1}(-0.6)$<br><b>Answer.</b> -0.64                         | 24. $\cos^{-1}\left(\frac{1}{4}\right)$<br><b>Answer.</b> 1.32 | 25. $\cos^{-1}(-0.4)$<br><b>Answer.</b> 1.98     |
| 26. $\tan^{-1}(-45.6)$<br><b>Answer.</b> -1.55                        | 27. $\sin^{-1}(-0.8)$<br><b>Answer.</b> -0.93                  | 28. $\tan^{-1}(78.9)$<br><b>Answer.</b> 1.56     |


**Exercise Group.** In [Exercise Group 1.4.7.1–8](#), we determined the values of the six trigonometric functions for each triangle. Now, use a calculator to find the value of  $\theta$  in degrees, rounding to two decimal places.

29.




Answer.  $53.13^\circ$

30.




Answer.  $35.54^\circ$

31.




Answer.  $56.31^\circ$

32.




Answer. 22.62

33.




Answer.  $63.43^\circ$

34.




Answer.  $45^\circ$

35.


 Answer.  $50.77^\circ$ 

36.


 Answer.  $33.69^\circ$ 

**Exercise Group.** Use properties of composition of trigonometric functions to find the exact value of each expression. Write “not defined” if no value exists.

37.  $\sin^{-1}(\sin \frac{\pi}{7})$

 Answer.  $\frac{\pi}{7}$ 

38.  $\cos^{-1}(\cos(-\frac{\pi}{7}))$

 Answer.  $\frac{\pi}{7}$ 

39.  $\sin(\sin^{-1}(-0.8))$

 Answer.  $-0.8$ 

40.  $\tan^{-1}(\tan(-\frac{2\pi}{9}))$

 Answer.  $-\frac{2\pi}{9}$ 

41.  $\cos(\cos^{-1} 1.4)$

Answer. Not defined

42.  $\sin(\sin^{-1} 0.74)$

 Answer.  $0.74$ 

43.  $\tan(\tan^{-1} 24)$

 Answer.  $24$ 

44.  $\cos^{-1}(\cos(-\frac{\pi}{5}))$

 Answer.  $\frac{\pi}{5}$ 

45.  $\tan(\tan^{-1}(-4.3))$

 Answer.  $-4.3$ 

46.  $\tan^{-1}(\tan \frac{\pi}{8})$

 Answer.  $\frac{\pi}{8}$ 

47.  $\cos(\cos^{-1} 0.39)$

 Answer.  $0.39$ 

48.  $\sin^{-1}(\sin \frac{4\pi}{5})$

 Answer.  $\frac{\pi}{5}$ 

**Exercise Group.** Find the exact value of each composite function. Write “not defined” if no value exists. You may find [Table 1.5.18](#) useful.

49.  $\sin(\cos^{-1} \frac{1}{2})$

 Answer.  $\frac{\sqrt{3}}{2}$ 

50.  $\cot(\sin^{-1}(-\frac{\sqrt{3}}{2}))$

 Answer.  $-\frac{\sqrt{3}}{3}$ 

51.  $\csc(\tan^{-1} 1)$

 Answer.  $\sqrt{2}$ 

52.  $\sec(\sin^{-1}(-\frac{1}{2}))$

 Answer.  $\frac{2\sqrt{3}}{3}$ 

53.  $\cos(\tan^{-1}(-\sqrt{3}))$

 Answer.  $\frac{1}{2}$ 

54.  $\tan(\cos^{-1} 0)$

Answer. Not defined

**Exercise Group.** Find the exact value of each composite function. Write “not defined” if no value exists.

55.  $\sin(\tan^{-1} \frac{3}{4})$

 Answer.  $\frac{3}{5}$ 

56.  $\csc(\cos^{-1} \frac{\sqrt{3}}{7})$

 Answer.  $\frac{7\sqrt{46}}{46}$ 

57.  $\sec(\tan^{-1}(-\sqrt{5}))$

 Answer.  $\sqrt{6}$ 

58.  $\cos(\sin^{-1} \frac{\sqrt{7}}{3})$

 Answer.  $\frac{\sqrt{2}}{3}$ 

59.  $\tan(\sin^{-1}(-\frac{1}{3}))$

 Answer.  $-\frac{\sqrt{8}}{8}$ 

60.  $\cot(\cos^{-1}(-\frac{2}{5}))$

 Answer.  $-\frac{2\sqrt{21}}{21}$

61.  $\tan(\csc^{-1}(\frac{9}{8}))$

Answer.  $\frac{8\sqrt{17}}{17}$

62.  $\sec(\sin^{-1}(-\frac{2}{5}))$

Answer.  $\frac{3\sqrt{2}}{4}$

**Exercise Group.** Utilizing the fact that  $x = \frac{x}{1}$  and sketching a right triangle, find the exact value of the expression in terms of  $x$ .

63.  $\cos(\tan^{-1}x)$

Answer.  $\frac{\sqrt{x^2+1}}{x^2+1}$

64.  $\sin(\cos^{-1}x)$

Answer.  $\sqrt{1-x^2}$

65.  $\sec(\sin^{-1}x)$

Answer.  $\frac{\sqrt{1-x^2}}{1-x^2}$


66.  $\sin(\sec^{-1}x)$

Answer.  $\frac{\sqrt{x^2-1}}{x}$

**Exercise Group.** Graph the function.


67.  $y = 4 \cos^{-1}x$

Answer.




68.  $y = \tan^{-1}x + \frac{\pi}{2}$

Answer.




69.  $y = \sin^{-1}(\frac{1}{2}x)$

Answer.



70.  $y = -\sin^{-1}(x-1)$

Answer.



71. **Sunrise on Mauna Kea.** The summit of Mauna Kea is the highest point in Hawai'i and sits 13,803 ft above the sea level. If you stand at the summit, you will be able to see the sun rise before someone standing at the sea level just north or south of Mauna Kea (at the same latitude). In fact, you will see the sunrise at the same time as someone at sea level sailing in a wa'a to the east of Mauna Kea. Assume the radius of the earth is 20,917,655 feet.

- (a) How far is the horizon when someone 5 ft tall stands at sea level to watch the sun rise? Round your answer to the nearest tenth of a mile.

Answer. 2.7 miles

- (b) How far is the horizon when someone 5 ft tall stands on the summit to watch the sunrise? Round your answer to the nearest tenth of a mile.


**Answer.** 143.9 miles

- (c) How much faster would someone 5 ft tall standing on the top of the mountain see the sunrise than someone 5 ft tall standing at sea level? Express your answer in minutes, rounded to one decimal place.

**Hint.** The ratio of the distance to the horizon (your previous answers) to the circumference of the Earth ( $2\pi R$ ) is equal to the ratio of the time it takes to see the sunrise to the 24 hours (1,440 minutes) in Earth's rotation. Compute the time it takes to see the sunrise from the top of the mountain and the shore, then calculate the difference.

**Answer.** 8.2 minutes

- 72. Maui Capturing the Sun.** According to mo'olelo, or legend, the sun traveled very fast across the sky, leaving people with days so short there was not enough time to carry on with their daily lives. Determined to slow the sun, the demigod Maui climbed to the summit of Haleakalā, which stands at 10,023 feet, to snare the sun. Assume the radius of the Earth at Haleakalā is the same as Mauna Kea at 20,917,655 feet.



- (a) How far is the horizon when Maui stands at the summit? Assume Maui's eyes are 5 ft from the ground. Round your answer to the nearest tenth of a mile.

**Answer.** 122.6 miles

- (b) How much faster would Maui see the sun emerge over the horizon than someone 5 ft tall and standing at the seashore? Express your answer in minutes, rounded to one decimal place.

**Answer.** 6.9 minutes

# Chapter 3

## Analytic Trigonometry

### 3.1 Trigonometric Identities

In this chapter, we explore trigonometric identities and formulas, essential tools that enable us to algebraically manipulate and solve complex trigonometric equations. These identities and formulas enable us to analyze expressions in various forms, often simplifying complex expressions into ones that are easily solvable and interpretable. By doing so, we increase our ability to accurately model the world around us.

#### 3.1.1 Fundamental Trigonometric Identities

An **identity** in mathematics is an equation that remains true for every valid values of its variables. We begin by reviewing some of the basic trigonometric identities from [Chapter 1](#), collectively known as the fundamental trigonometric identities.

**Definition 3.1.1** The fundamental trigonometric identities are

1. **Reciprocal Identities** ([Definition 1.4.2](#))

$$\begin{array}{lll} \sin \theta = \frac{1}{\csc \theta} & \cos \theta = \frac{1}{\sec \theta} & \tan \theta = \frac{1}{\cot \theta} \\ \csc \theta = \frac{1}{\sin \theta} & \sec \theta = \frac{1}{\cos \theta} & \cot \theta = \frac{1}{\tan \theta} \end{array}$$

2. **Quotient Identities** ([Definition 1.4.3](#))

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta}$$

3. **Pythagorean Identities** ([Definition 1.5.20](#))

$$\begin{aligned} \sin^2 \theta + \cos^2 \theta &= 1 \\ 1 + \tan^2 \theta &= \sec^2 \theta \\ 1 + \cot^2 \theta &= \csc^2 \theta \end{aligned}$$

4. **Odd-Even Identities** ([Definition 1.5.22](#))

The cosine and secant functions are **even**

$$\cos(-\theta) = \cos \theta \qquad \sec(-\theta) = \sec \theta$$

The sine, cosecant, tangent, and cotangent functions are **odd**

$$\begin{aligned}\sin(-\theta) &= -\sin \theta & \csc(-\theta) &= -\csc(\theta) \\ \tan(-\theta) &= -\tan \theta & \cot(-\theta) &= -\cot(\theta)\end{aligned}$$

### 5. Cofunction Identities (Definition 1.4.7)

$$\begin{aligned}\sin \theta &= \cos\left(\frac{\pi}{2} - \theta\right), & \cos \theta &= \sin\left(\frac{\pi}{2} - \theta\right) \\ \tan \theta &= \cot\left(\frac{\pi}{2} - \theta\right), & \cot \theta &= \tan\left(\frac{\pi}{2} - \theta\right) \\ \sec \theta &= \csc\left(\frac{\pi}{2} - \theta\right), & \csc \theta &= \sec\left(\frac{\pi}{2} - \theta\right)\end{aligned}$$

◇

## 3.1.2 Simplifying Trigonometric Expressions

We use a combination of trigonometric identities, formulas, and techniques from algebra to manipulate and simplify trigonometric expressions.

**Example 3.1.2** Simplify

$$\tan^2(x) \cdot \csc^2(x)$$

**Solution.** We can simplify this expression by writing each function in terms of sine and cosine functions

$$\tan^2(x) \cdot \csc^2(x) = \frac{\sin^2(x)}{\cos^2(x)} \cdot \frac{1}{\sin^2(x)} = \frac{1}{\cos^2(x)} = \sec^2(x)$$

□

**Example 3.1.3** Simplify

$$\sin^2(x)(\cot^2(x) - 1)$$

**Solution.** We can simplify this expression by first using the Pythagorean Identity and then using the Reciprocal Identity:

$$\sin^2(x)(\cot^2(x) - 1) = \sin^2(x)(-\csc^2(x)) = \sin^2(x)\left(-\frac{1}{\sin^2(x)}\right) = -1$$

□

## 3.1.3 Verifying Trigonometric Identities

To verify trigonometric identities, we begin with an expression on one side of the equation and manipulate it using trigonometric identities and algebraic techniques until it matches the expression on the other side.

**Remark 3.1.4 Steps for Verifying Trigonometric Identities.** To verify that an equation is an identity:

1. Pick an expression on one side of the equation. Often it is the more complicated expression.
2. Transform the expression using techniques such as trigonometric identities, rewriting in terms of sine and cosine functions, factoring, common denominator, or multiplying numerator and denominator by the same

term.

3. Continue manipulating until the transformed expression matches the other side of the equation.
4. If you have trouble making one side resemble the other, try manipulating both sides separately, and make them match to reach the same result.

Note: Unlike solving equations where we perform the same operation on both sides of the equal sign, when verifying trigonometric identities, we work with only one side and manipulate it to resemble the other side.

**Example 3.1.5 Verify the Identity by Rewriting in Terms of Sine and Cosine.** Verify the identity

$$\frac{\sin(x)}{\tan(x)} = \cos(x)$$

**Solution.** We use the Quotient Identity to rewrite  $\tan(x)$  in terms of  $\sin(x)$  and  $\cos(x)$ :

$$\frac{\sin(x)}{\tan(x)} = \frac{\sin(x)}{\frac{\sin(x)}{\cos(x)}} = \sin(x) \cdot \frac{\cos(x)}{\sin(x)} = \cos(x).$$

□

**Example 3.1.6 Verify the Identity by Factoring.** Verify the identity

$$\cos^4(x) + \sin^2(x) \cos^2(x) = \cos^2(x).$$

**Solution.** First notice that both terms in  $\cos^4(x) + \sin^2(x) \cos^2(x)$  contain  $\cos^2(x)$ . Then

$$\begin{aligned} \cos^4(x) + \sin^2(x) \cos^2(x) &= \cos^2(x) \cdot \cos^2(x) + \sin^2(x) \cos^2(x) \\ &= \cos^2(x) \cdot (\cos^2(x) + \sin^2(x)) \\ &= \cos^2(x) \cdot 1 \\ &= \cos^2(x) \end{aligned}$$

□

**Example 3.1.7 Verify the Identity by Odd-Even Properties.** Verify the identity

$$\frac{\cos(x) - \sin(x)}{\cos(-x) + \sin(-x)} = 1$$

**Solution.** By the Odd-Even Properties, we have  $\sin(-x) = -\sin(x)$  and  $\cos(-x) = \cos(x)$ . Thus,

$$\frac{\cos(x) - \sin(x)}{\cos(-x) + \sin(-x)} = \frac{\cos(x) - \sin(x)}{\cos(x) - \sin(x)} = 1$$

□

**Example 3.1.8 Verify the Identity by Multiplying the Numerator and Denominator by the Same Term.** Verify the identity

$$\frac{\sin(x)}{\sin(x) + \cos(x)} = \frac{1}{1 + \cot(x)}$$



**Solution.** Multiplying both the numerator and denominator by  $\frac{1}{\sin(x)}$ , we get

$$\begin{aligned} \frac{\sin(x)}{\sin(x) + \cos(x)} \cdot \frac{\frac{1}{\sin(x)}}{\frac{1}{\sin(x)}} &= \frac{\sin(x) \cdot \frac{1}{\sin(x)}}{\sin(x) \cdot \frac{1}{\sin(x)} + \cos(x) \cdot \frac{1}{\sin(x)}} \\ &= \frac{1}{1 + \frac{\cos(x)}{\sin(x)}} \\ &= \frac{1}{1 + \cot(x)} \end{aligned}$$

□

**Example 3.1.9 Verify the Identity by Manipulating Both Sides Separately.** Verify the identity

$$\frac{1 - \cos x}{1 + \cos x} = (\csc x - \cot x)^2.$$

**Solution.** We begin by simplifying the right-hand side of the equation

$$\begin{aligned} (\csc x - \cot x)^2 &= \csc^2 x - 2 \csc x \cot x + \cot^2 x \\ &= \csc^2 x + \cot^2 x - 2 \csc x \cot x \\ &= \csc^2 x + \cot^2 x - 2 \frac{1}{\sin x} \frac{\cos x}{\sin x} \\ &= \csc^2 x + \cot^2 x - 2 \frac{\cos x}{\sin^2 x}. \end{aligned}$$

Next, we will manipulate the left-hand side of the equation to get it to look like  $\csc^2 x + \cot^2 x - 2 \frac{\cos x}{\sin^2 x}$ .

$$\begin{aligned} \frac{1 - \cos x}{1 + \cos x} &= \frac{(1 - \cos x)(1 - \cos x)}{(1 + \cos x)(1 - \cos x)} \\ &= \frac{1 - 2 \cos x + \cos^2 x}{1 - \cos^2 x} \\ &= \frac{1 + \cos^2 x - 2 \cos x}{\sin^2 x} \\ &= \frac{1}{\sin^2 x} + \frac{\cos^2 x}{\sin^2 x} - 2 \frac{\cos x}{\sin^2 x} \\ &= \csc^2 x + \cot^2 x - 2 \frac{\cos x}{\sin^2 x}. \end{aligned}$$

Thus, since the left-hand side and the right-hand side of the equation can both be manipulated to  $\csc^2 x + \cot^2 x - 2 \frac{\cos x}{\sin^2 x}$ , we have established the identity. □

### 3.1.4 Exercises

**Exercise Group.** Verify the identity.


1.  $\cos \theta \sec \theta = 1$
2.  $\cos x \csc x = \cot x$
3.  $\frac{\cos \theta \sec \theta}{\tan \theta} = \cot \theta$
4.  $\frac{\cot t \tan t}{\csc t} = \sin t$
5.  $(1 + \tan \theta)(1 - \tan \theta) + \sec^2 \theta = 2$

6.  $1 - \sin^2(x) = \cos^2(x)$
7.  $1 - \sec^2(\theta) = -\tan^2(\theta)$
8.  $\tan(t) \cdot \cot(t) = 1$
9.  $(\sin \theta + \cos \theta)^2 = 1 + 2 \sin \theta \cos \theta$
10.  $(1 - \cot \theta)^2 = \csc^2 \theta - 2 \cot \theta$
11.  $\frac{\sin \theta}{\csc \theta} + \frac{\cos \theta}{\sec \theta} = 1$
12.  $\sin^2 t (\csc^2 t + \sec^2 t) = \sec^2 t$
13.  $\sin^2(x) - \sin^2(x) \cos^2(x) = \sin^4(x)$
14.  $\sin^2(-x) + \cos^2(-x) = 1$
15.  $\cos(-t) + \sin(-t) = \cos(t) - \sin(t)$
16.  $(\sin \theta + \cos \theta)^2 - 2 \sin \theta \cos \theta = 1$
17.  $\cot^2 x (\sec^2 x - 1) = 1$
18.  $(1 + \sin(t))(1 + \sin(-t)) = \cos^2 t$
19.  $\tan^4 \theta = \tan^2 \theta \sec^2 \theta - \tan^2 \theta$
20.  $\frac{1}{1 - \sin x} + \frac{1}{1 + \sin x} = 2 \sec^2 x$
21.  $\frac{1}{\csc t + 1} - \frac{1}{\csc t - 1} = -2 \tan^2 t$
22.  $\frac{1}{1 - \cos \theta} + \frac{1}{1 + \cos \theta} = 2 \csc^2 \theta$
23.  $\frac{1}{1 - \cos \theta} + \frac{1}{1 + \cos \theta} = 2 + 2 \cot^2 \theta$
24.  $\frac{1 - \cos^2 x}{\cos x} = \sin x \tan x$
25.  $1 - \frac{\cos^2 \theta}{1 + \sin \theta} = \sin \theta$
26.  $\sec^2 t + \csc^2 t = \csc^2 t \sec^2 t$
27.  $\frac{1 + \tan x}{1 - \tan x} = \frac{\cot x + 1}{\cot x - 1}$
28.  $\frac{\cos \theta}{1 - \sin \theta} = \frac{1 + \sin \theta}{\cos \theta}$
29.  $1 - \frac{\cos^2 t}{1 + \sin t} = \sin t$
30.  $\frac{1 + \cos \theta}{\cos \theta} = \frac{\tan^2 \theta}{\sec \theta - 1}$

### 3.2 Addition and Subtraction Formulas

In [Section 1.4](#), we used right triangles to determine the deviation of a wa'a (canoe) from its course based on the angle of deviation. If a wa'a sails for 120 nautical miles (NM), we were able to calculate the deviation from its course using right triangles to get the equation:

$$\text{deviation} = (120 \text{ NM}) \cdot \sin(\theta).$$



Before setting sail, a voyager studies a table listing the deviation distances corresponding to different houses of deviation. It's crucial to understand that while adding angles may yield a third angle, adding their corresponding deviations will not accurately determine the total deviation. In other words:

$$\sin(\alpha) + \sin(\beta) \neq \sin(\alpha + \beta)$$


for some angles of deviation  $\alpha$  and  $\beta$ .

To illustrate this, let's calculate the deviation distances for 1, 2, and 3 houses respectively. Using the given formula, we have:

$$120 \sin(1 \text{ house}) = 120 \sin(11.25^\circ) \approx 23.4 \text{ NM}$$

$$120 \sin(2 \text{ houses}) = 120 \sin(22.5^\circ) \approx 45.9 \text{ NM}$$

$$120 \sin(3 \text{ houses}) = 120 \sin(33.75^\circ) \approx 66.7 \text{ NM}$$



However,

$$120 \sin(1 \text{ house}) + 120 \sin(2 \text{ houses}) \approx 23.4 + 45.9 \text{ NM} \approx 69.3 \text{ NM},$$

which differs from the actual deviation of 66.7 NM when deviating by 3 houses.

These calculations demonstrate that the deviation distances for multiple houses cannot be determined by simply adding individual deviations, highlighting the importance of understanding trigonometric principles for accurate navigation. In this section, we will explore formulas for the addition and subtraction of angles in trigonometric functions.

### 3.2.1 Addition and Subtraction Formulas for Cosine

First we will derive the addition and subtraction formulas for the cosine function.

#### Definition 3.2.1 Addition and Subtraction Formulas for Cosine.

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$


$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

◇

*Proof.* First we will prove the Subtraction Formula for Cosine

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta.$$

We begin by considering two points on the unit circle. Point  $P$  is at an angle of  $\beta$  in standard position with coordinates  $(\cos \beta, \sin \beta)$  and point  $Q$  is at an angle of  $\alpha$  in standard position with coordinates  $(\cos \alpha, \sin \alpha)$ .



We use the Distance Formula to calculate the distance between  $P$  and  $Q$  to get


$$\begin{aligned} d(P, Q) &= \sqrt{(\cos \alpha - \cos \beta)^2 + (\sin \alpha - \sin \beta)^2} \\ &= \sqrt{\cos^2 \alpha - 2 \cos \alpha \cos \beta + \cos^2 \beta + \sin^2 \alpha - 2 \sin \alpha \sin \beta + \sin^2 \beta} \\ &= \sqrt{(\cos^2 \alpha + \sin^2 \alpha) + (\cos^2 \beta + \sin^2 \beta) - 2 \cos \alpha \cos \beta - 2 \sin \alpha \sin \beta} \end{aligned}$$

Then by the Pythagorean Identity (Definition 1.5.20),  $\cos^2 \alpha + \sin^2 \alpha = 1$  and  $\cos^2 \beta + \sin^2 \beta = 1$ . Thus the distance becomes

$$d(P, Q) = \sqrt{1 + 1 - 2 \cos \alpha \cos \beta - 2 \sin \alpha \sin \beta}$$

$$= \sqrt{2 - 2 \cos \alpha \cos \beta - 2 \sin \alpha \sin \beta}$$

Next, consider two additional points on a second unit circle. Point  $A$  has coordinates at  $(1, 0)$  and Point  $B$  is at an angle of  $\alpha - \beta$  in standard position with coordinates  $(\cos(\alpha - \beta), \sin(\alpha - \beta))$ .



The distance between  $A$  and  $B$  is

$$\begin{aligned} d(A, B) &= \sqrt{(\cos(\alpha - \beta) - 1)^2 + (\sin(\alpha - \beta) - 0)^2} \\ &= \sqrt{\cos^2(\alpha - \beta) - 2 \cos(\alpha - \beta) + 1 + \sin^2(\alpha - \beta)} \\ &= \sqrt{(\cos^2(\alpha - \beta) + \sin^2(\alpha - \beta)) - 2 \cos(\alpha - \beta) + 1} \\ &= \sqrt{1 - 2 \cos(\alpha - \beta) + 1} \\ &= \sqrt{2 - 2 \cos(\alpha - \beta)} \end{aligned}$$

Note that since  $OP$ ,  $OQ$ ,  $OA$ , and  $OB$  are lines from the center to points on the unit circle, they are congruent and have length of 1. Also note that  $\angle POB = \angle AOB = \alpha - \beta$ . Then since two sides and the included angle of  $\triangle OPQ$  and  $\triangle OAB$  are equivalent, we can conclude by the Side-Angle-Side Theorem (SAS) in geometry that the two triangles are congruent. Thus, corresponding sides have the same lengths, giving us:  $d(PQ) = d(AB)$ . Substituting our results for  $d(P, Q)$  and  $d(A, B)$  we get

$$\begin{aligned} d(PQ) &= d(AB) \\ \sqrt{2 - 2 \cos \alpha \cos \beta - 2 \sin \alpha \sin \beta} &= \sqrt{2 - 2 \cos(\alpha - \beta)} \\ 2 - 2 \cos \alpha \cos \beta - 2 \sin \alpha \sin \beta &= 2 - 2 \cos(\alpha - \beta) \\ -2 \cos \alpha \cos \beta - 2 \sin \alpha \sin \beta &= -2 \cos(\alpha - \beta) \end{aligned}$$

Dividing both sides by -2 we arrive at the Subtraction Formula for Cosine

$$\cos \alpha \cos \beta + \sin \alpha \sin \beta = \cos(\alpha - \beta)$$

To prove the Addition Formula for Cosine, replace  $\beta$  with  $-\beta$  in the Subtraction Formula and use the Even and Odd Trigonometric Properties (Definition 1.5.22) where  $\sin(-\beta) = -\sin \beta$  and  $\cos(-\beta) = \cos \beta$  to get

$$\begin{aligned}\cos \alpha \cos(-\beta) + \sin \alpha \sin(-\beta) &= \cos(\alpha - (-\beta)) \\ \cos \alpha \cos(\beta) + \sin \alpha(-\sin \beta) &= \cos(\alpha + \beta) \\ \cos \alpha \cos \beta - \sin \alpha \sin \beta &= \cos(\alpha + \beta)\end{aligned}$$

■

**Example 3.2.2** Find the exact value of  $\cos 105^\circ$ .

**Solution.** First note that  $105^\circ = 60^\circ + 45^\circ$ . Now, using the Addition Formula for Cosine (Definition 3.2.1),

$$\begin{aligned}\cos 105^\circ &= \cos(60^\circ + 45^\circ) \\ &= \cos 60^\circ \cos 45^\circ - \sin 60^\circ \sin 45^\circ \\ &= \frac{1}{2} \cdot \frac{\sqrt{2}}{2} - \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} \\ &= \frac{1}{4} (\sqrt{2} - \sqrt{6})\end{aligned}$$

□

**Example 3.2.3** Find the exact value of  $\cos\left(\frac{\pi}{4} - \frac{\pi}{6}\right)$ .

**Solution.** Using the Subtraction Formula for Cosine we get

$$\begin{aligned}\cos\left(\frac{\pi}{4} - \frac{\pi}{6}\right) &= \cos \frac{\pi}{4} \cos \frac{\pi}{6} + \sin \frac{\pi}{4} \sin \frac{\pi}{6} \\ &= \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2} + \frac{\sqrt{2}}{2} \cdot \frac{1}{2} \\ &= \frac{1}{4} (\sqrt{6} + \sqrt{2})\end{aligned}$$

□

**Example 3.2.4** Find the exact value of the expression  $\cos 25^\circ \cos 35^\circ - \sin 25^\circ \sin 35^\circ$ .

**Solution.** Notice this expression is the Addition Formula for Cosine with  $\alpha = 25^\circ$  and  $\beta = 35^\circ$ . So

$$\cos 25^\circ \cos 35^\circ - \sin 25^\circ \sin 35^\circ = \cos(25^\circ + 35^\circ) = \cos 60^\circ = \frac{1}{2}$$

□

### 3.2.2 Addition and Subtraction Formulas for Sine

Next we will learn about the addition and subtraction formulas for the sine function.

**Definition 3.2.5** Addition and Subtraction Formulas for Sine.

$$\begin{aligned}\sin(\alpha + \beta) &= \sin \alpha \cos \beta + \cos \alpha \sin \beta \\ \sin(\alpha - \beta) &= \sin \alpha \cos \beta - \cos \alpha \sin \beta\end{aligned}$$

◇

We will prove the Addition Formula for Sine in [Example 3.2.10](#) and the Subtraction Formula can be established using the Even and Odd Properties ([Definition 1.5.22](#)).


**Example 3.2.6** Given  $\sin \alpha = -\frac{12}{13}$ , with  $\frac{3\pi}{2} < \alpha < 2\pi$  and  $\cos \beta = -\frac{3}{5}$ , with  $\frac{\pi}{2} < \beta < \pi$ , find the exact value of  $\sin(\alpha + \beta)$ .

**Solution.** The Addition Formula for Sine gives us

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

At this moment, we do not know the exact values of  $\cos \alpha$  and  $\sin \beta$  but we can compute them.

Given  $\sin \alpha = -\frac{12}{13}$  with  $\frac{3\pi}{2} < \alpha < 2\pi$  and  $\cos \beta = -\frac{3}{5}$  with  $\frac{\pi}{2} < \beta < \pi$ , we can draw the following triangles associated with  $\alpha$  and  $\beta$ , respectively:



Next, using the Pythagorean Theorem, we solve for the missing sides on the triangles

$$\begin{aligned} x^2 + (-12)^2 &= 13^2 \\ x + 144 &= 169 \\ x^2 &= 25 \\ x &= 5 \end{aligned}$$

Thus we get

$$\cos \alpha = \frac{5}{13}$$

$$(-3)^2 + y^2 = 5^2$$

$$9 + y^2 = 25$$

$$y^2 = 16$$

$$y = 4$$

Thus we get

$$\sin \beta = -\frac{4}{5}$$

We now have all the information needed to proceed.

$$\begin{aligned} \sin(\alpha + \beta) &= \sin \alpha \cos \beta + \cos \alpha \sin \beta \\ &= \left(-\frac{12}{13}\right) \left(-\frac{3}{5}\right) + \left(\frac{5}{13}\right) \left(\frac{4}{5}\right) \\ &= \frac{36}{65} + \frac{20}{65} \\ &= \frac{56}{65} \end{aligned}$$

Notice we did not have to know the values of  $\alpha$  or  $\beta$  to do this example.  $\square$

### 3.2.3 Addition and Subtraction Formulas for Tangent

Now we will learn about the addition and subtraction formulas for the tangent function.

#### Definition 3.2.7 Addition and Subtraction Formulas for Tangent.

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$\tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

$\diamond$

*Proof.* Recall that  $\tan \theta = \frac{\sin \theta}{\cos \theta}$  as long as  $\cos \theta \neq 0$ . Using this fact, and our new formulas for the sum of sine and cosine, we get

$$\begin{aligned} \tan(\alpha + \beta) &= \frac{\sin(\alpha + \beta)}{\cos(\alpha + \beta)} \\ &= \frac{\sin \alpha \cos \beta + \cos \alpha \sin \beta}{\cos \alpha \cos \beta - \sin \alpha \sin \beta} \\ &= \frac{\sin \alpha \cos \beta + \cos \alpha \sin \beta}{\frac{\cos \alpha \cos \beta}{\cos \alpha \cos \beta} - \frac{\sin \alpha \sin \beta}{\cos \alpha \cos \beta}} \\ &= \frac{\sin \alpha \cos \beta}{\cos \alpha \cos \beta} + \frac{\cos \alpha \sin \beta}{\cos \alpha \cos \beta} \\ &= \frac{\cancel{\cos \alpha} \cos \beta}{\cancel{\cos \alpha} \cos \beta} - \frac{\sin \alpha \sin \beta}{\cos \alpha \cos \beta} \end{aligned}$$



$$\begin{aligned} & \frac{\sin \alpha}{\cos \alpha} + \frac{\sin \beta}{\cos \beta} \\ &= \frac{\sin \alpha \cos \beta + \sin \beta \cos \alpha}{\cos \alpha \cos \beta} \\ &= \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} \end{aligned}$$

The subtraction formula can be established using the Even and Odd Properties (Definition 1.5.22). ■

**Example 3.2.8** Find the exact value of  $\tan\left(\frac{3\pi}{4} + \frac{\pi}{6}\right)$

**Solution.** Using the Addition Formula for Tangent (Definition 3.2.7), we get

$$\begin{aligned} \tan\left(\frac{3\pi}{4} + \frac{\pi}{6}\right) &= \frac{\tan \frac{3\pi}{4} + \tan \frac{\pi}{6}}{1 - \tan \frac{3\pi}{4} \tan \frac{\pi}{6}} \\ &= \frac{(-1) + \frac{\sqrt{3}}{3}}{1 - (-1) \cdot \frac{\sqrt{3}}{3}} \\ &= \frac{-1 + \frac{\sqrt{3}}{3}}{1 + \frac{\sqrt{3}}{3}} \\ &= \frac{\frac{-3 + \sqrt{3}}{3}}{\frac{3 + \sqrt{3}}{3}} \\ &= \frac{-3 + \sqrt{3}}{3 + \sqrt{3}} \end{aligned}$$

□

### 3.2.4 Cofunction Identities

Recall the Cofunction Identities (Definition 1.4.7):

$$\begin{aligned} \sin \theta &= \cos\left(\frac{\pi}{2} - \theta\right), & \cos \theta &= \sin\left(\frac{\pi}{2} - \theta\right) \\ \tan \theta &= \cot\left(\frac{\pi}{2} - \theta\right), & \cot \theta &= \tan\left(\frac{\pi}{2} - \theta\right) \\ \sec \theta &= \csc\left(\frac{\pi}{2} - \theta\right), & \csc \theta &= \sec\left(\frac{\pi}{2} - \theta\right) \end{aligned}$$

Armed with the knowledge of the subtraction formulas, we can prove the Cofunction Identities.

*Proof.* We will prove the Cofunction Identity for  $\sin \theta$  in Example 3.2.9. The proof for  $\cos \theta$  is given as Exercise 3.2.7.97. The Cofunction Identities for  $\tan \theta$  and  $\cot \theta$  can be found using the Quotient Identities (Definition 1.4.3) for  $\csc \theta$  and  $\sec \theta$  can be found using the Reciprocal Identities (Definition 1.4.2). ■


**Example 3.2.9** Use the Subtraction Formula for Sine to establish the identity  $\cos \theta = \sin\left(\frac{\pi}{2} - \theta\right)$ .

**Solution.** To establish an identity, we will start from one side of the equality and use properties to end up with the expression on the other side of the equality. So,

$$\sin\left(\frac{\pi}{2} - \theta\right) = \sin\left(\frac{\pi}{2}\right) \cos(\theta) - \cos\left(\frac{\pi}{2}\right) \sin(\theta)$$

$$\begin{aligned}
 &= 1 \cdot \cos(\theta) - 0 \cdot \sin(\theta) \\
 &= \cos(\theta)
 \end{aligned}$$

Visually, we have  $\cos \theta = \frac{x}{r} = \sin\left(\frac{\pi}{2} - \theta\right)$ :



□

**Example 3.2.10** Prove the Addition Formula for Sine

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

**Solution.** We begin by using the cofunction identity

$$\begin{aligned}
 \sin(\alpha + \beta) &= \cos\left(\frac{\pi}{2} - (\alpha + \beta)\right) \\
 &= \cos\left(\left(\frac{\pi}{2} - \alpha\right) - \beta\right)
 \end{aligned}$$

By the Subtraction Formula for Cosine:

$$\begin{aligned}
 &= \cos\left(\frac{\pi}{2} - \alpha\right) \cos \beta + \sin\left(\frac{\pi}{2} - \alpha\right) \sin \beta \\
 &= \sin \alpha \cos \beta + \cos \alpha \sin \beta
 \end{aligned}$$

where the last step we use the Cofunction Identity. □

### 3.2.5 Sums of Sines and Cosines

Sometimes we may come across functions of the form

$$a \sin x + b \cos x$$

It can often be useful to rewrite this expression as a single trigonometric function.

**Definition 3.2.11** For any real numbers  $a$  and  $b$ , let  $\theta$  be an angle in standard position where  $P(a, b)$  is a point on the terminal side of  $\theta$ . Then

$$a \sin x + b \cos x = \sqrt{a^2 + b^2} \sin(x + \theta).$$


◇

*Proof.* We begin by considering the triangle formed by the angle  $\theta$  and point  $P(a, b)$ , shown in Figure 3.2.12. By the Pythagorean Theorem, the hypotenuse of this triangle, with base  $a$  and height  $b$ , is  $\sqrt{a^2 + b^2}$ . According to Definition 1.4.1, we have:

$$\cos \theta = \frac{a}{\sqrt{a^2 + b^2}}, \quad \sin \theta = \frac{b}{\sqrt{a^2 + b^2}}$$

or equivalently:

$$a = \sqrt{a^2 + b^2} \cos \theta, \quad b = \sqrt{a^2 + b^2} \sin \theta$$



**Figure 3.2.12** A triangle is formed by angle  $\theta$  and point  $P(a, b)$ .

Therefore, using the addition formula for sine, we get

$$\begin{aligned} a \sin x + b \cos x &= \sqrt{a^2 + b^2} \cos \theta \sin x + \sqrt{a^2 + b^2} \sin \theta \cos x \\ &= \sqrt{a^2 + b^2} (\cos \theta \sin x + \sin \theta \cos x) \\ &= \sqrt{a^2 + b^2} \sin(x + \theta) \end{aligned}$$

■

**Example 3.2.13** Express

$$-\frac{\sqrt{3}}{2} \sin x + \frac{1}{2} \cos x$$

in terms of sine only.

**Solution.** To express the given expression in terms of sine only, we will use [Definition 3.2.11](#). Considering the point  $P(a, b) = \left(-\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ , which lies in Quadrant II, we determine the angle  $\theta$ . Using either [Table 1.5.18](#) or inverse trigonometric methods where

$$\tan \theta = \frac{b}{a},$$

we find

$$\theta = 150^\circ.$$

Therefore, by [Definition 3.2.11](#), we have:

$$\begin{aligned} -\frac{\sqrt{3}}{2} \sin x + \frac{1}{2} \cos x &= \sqrt{\left(-\frac{\sqrt{3}}{2}\right)^2 + \left(\frac{1}{2}\right)^2} \sin(x + 150^\circ) \\ &= \sqrt{\frac{3}{4} + \frac{1}{4}} \sin(x + 150^\circ) \\ &= \sin(x + 150^\circ) \end{aligned}$$

□

### 3.2.6 Summary

To review, the addition and subtraction formulas are

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$\tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

### 3.2.7 Exercises

**Exercise Group.** Use the Addition or Subtraction Formula to find the exact value of each expression.

1.  $\cos(15^\circ)$

**Answer.**  $\frac{\sqrt{6}+\sqrt{2}}{4}$

3.  $\sin(195^\circ)$

**Answer.**  $\frac{-\sqrt{6}+\sqrt{2}}{4}$

5.  $\tan(105^\circ)$

**Answer.**  $\frac{1+\sqrt{3}}{1-\sqrt{3}}$

7.  $\tan\left(\frac{5\pi}{12}\right)$

**Hint.**  $\frac{5\pi}{12} = \frac{\pi}{6} + \frac{\pi}{4}$

**Answer.**  $\frac{3+\sqrt{3}}{3-\sqrt{3}}$

9.  $\cos\left(\frac{\pi}{12}\right)$

**Hint.**  $\frac{\pi}{12} = \frac{\pi}{3} - \frac{\pi}{4}$

**Answer.**  $\frac{\sqrt{6}+\sqrt{2}}{4}$

11.  $\cos\left(\frac{17\pi}{12}\right)$

**Hint.**  $\frac{17\pi}{12} = \frac{\pi}{4} + \frac{5\pi}{3}$

**Answer.**  $\frac{\sqrt{2}-\sqrt{6}}{4}$

2.  $\sin(75^\circ)$

**Answer.**  $\frac{\sqrt{6}+\sqrt{2}}{4}$

4.  $\tan(165^\circ)$

**Answer.**  $\frac{1-\sqrt{3}}{1+\sqrt{3}}$

6.  $\cos(255^\circ)$

**Answer.**  $\frac{-\sqrt{6}+\sqrt{2}}{4}$

8.  $\sin\left(\frac{7\pi}{12}\right)$

**Hint.**  $\frac{7\pi}{12} = \frac{\pi}{3} + \frac{\pi}{4}$

**Answer.**  $\frac{\sqrt{6}+\sqrt{2}}{4}$

10.  $\sin\left(\frac{13\pi}{12}\right)$

**Hint.**  $\frac{13\pi}{12} = \frac{11\pi}{6} - \frac{3\pi}{4}$

**Answer.**  $\frac{\sqrt{2}-\sqrt{6}}{4}$

12.  $\tan\left(-\frac{\pi}{12}\right)$

**Hint.**  $-\frac{\pi}{12} = \frac{\pi}{4} - \frac{\pi}{3}$

**Answer.**  $\frac{1-\sqrt{3}}{1+\sqrt{3}}$

**Exercise Group.** Use the Addition or Subtraction Formula to find the exact value of each expression.

13.  $\sin(172^\circ) \cos(68^\circ) + \cos(172^\circ) \sin(68^\circ)$

**Answer.**  $-\frac{\sqrt{3}}{2}$

14.  $\sin(317^\circ) \cos(257^\circ) - \cos(317^\circ) \sin(257^\circ)$

**Answer.**  $\frac{\sqrt{3}}{2}$

15.  $\cos(337^\circ) \cos(22^\circ) + \sin(337^\circ) \sin(22^\circ)$

**Answer.**  $-\frac{1}{2}$

16.  $\cos(59^\circ) \cos(211^\circ) - \sin(59^\circ) \sin(211^\circ)$

**Answer.** 0

$$17. \frac{\tan(85^\circ) - \tan(25^\circ)}{1 + \tan(85^\circ)\tan(25^\circ)}$$

$$\text{Answer. } \sqrt{3}$$

$$18. \sin\left(\frac{5\pi}{16}\right)\cos\left(\frac{\pi}{16}\right) - \cos\left(\frac{5\pi}{16}\right)\sin\left(\frac{\pi}{16}\right)$$

$$\text{Answer. } \frac{\sqrt{2}}{2}$$



$$19. \cos\left(\frac{5\pi}{22}\right)\cos\left(\frac{3\pi}{11}\right) - \sin\left(\frac{5\pi}{22}\right)\sin\left(\frac{3\pi}{11}\right)$$

$$\text{Answer. } 0$$

$$20. \frac{\tan\left(\frac{\pi}{7}\right) + \tan\left(\frac{4\pi}{21}\right)}{1 - \tan\left(\frac{\pi}{7}\right)\tan\left(\frac{4\pi}{21}\right)}$$

$$\text{Answer. } \sqrt{3}$$

**Exercise Group.** Find the exact value of each expression given



$$21. \sin(\alpha)$$

$$\text{Answer. } \frac{3}{5}$$

$$23. \tan(\alpha)$$

$$\text{Answer. } \frac{3}{4}$$

$$25. \cos(\beta)$$

$$\text{Answer. } \frac{21}{29}$$

$$27. \sin(\alpha + \beta)$$

$$\text{Answer. } \frac{3}{5} \cdot \frac{21}{29} + \frac{4}{5} \cdot \frac{20}{29} = \frac{143}{145}$$

$$29. \tan(\alpha + \beta)$$

$$\text{Answer. } \frac{\frac{3}{4} + \frac{20}{21}}{1 - \frac{3}{4} \cdot \frac{20}{21}} = \frac{143}{24}$$

$$31. \cos(\alpha - \beta)$$

$$\text{Answer. } \frac{4}{5} \cdot \frac{21}{29} + \frac{3}{5} \cdot \frac{20}{29} = \frac{144}{145}$$

$$22. \cos(\alpha)$$

$$\text{Answer. } \frac{4}{5}$$

$$24. \sin(\beta)$$

$$\text{Answer. } \frac{20}{29}$$

$$26. \tan(\beta)$$

$$\text{Answer. } \frac{20}{21}$$

$$28. \cos(\alpha + \beta)$$

$$\text{Answer. } \frac{4}{5} \cdot \frac{21}{29} - \frac{3}{5} \cdot \frac{20}{29} = \frac{24}{145}$$


$$30. \sin(\alpha - \beta)$$

$$\text{Answer. } \frac{3}{5} \cdot \frac{21}{29} - \frac{4}{5} \cdot \frac{20}{29} = -\frac{17}{145}$$

$$32. \tan(\alpha - \beta)$$

$$\text{Answer. } \frac{\frac{3}{4} - \frac{20}{21}}{1 + \frac{3}{4} \cdot \frac{20}{21}} = -\frac{17}{24}$$

**Exercise Group.** Find the exact value of each expression given



33.  $\sin(\alpha)$

Answer.  $\frac{\sqrt{15}}{5}$

35.  $\tan(\alpha)$

Answer.  $\frac{\sqrt{6}}{2}$

37.  $\cos(\beta)$

Answer.  $\frac{2\sqrt{13}}{13}$

39.  $\sin(\alpha + \beta)$

Answer.  $\frac{\sqrt{15}}{5} \cdot \frac{2\sqrt{13}}{13} + \frac{\sqrt{10}}{5} \cdot \frac{3\sqrt{13}}{13} = \frac{2\sqrt{195} + 3\sqrt{130}}{65}$

41.  $\tan(\alpha + \beta)$

Answer.  $\frac{\frac{\sqrt{6}}{2} + \frac{3}{2}}{1 - \frac{\sqrt{6}}{2} \cdot \frac{3}{2}} = \frac{-30 - 13\sqrt{6}}{19}$

43.  $\cos(\alpha - \beta)$

Answer.  $\frac{\sqrt{10}}{5} \cdot \frac{2\sqrt{13}}{13} + \frac{\sqrt{15}}{5} \cdot \frac{3\sqrt{13}}{13} = \frac{2\sqrt{130} + 3\sqrt{195}}{65}$

34.  $\cos(\alpha)$

Answer.  $\frac{\sqrt{10}}{5}$

36.  $\sin(\beta)$

Answer.  $\frac{3\sqrt{13}}{13}$

38.  $\tan(\beta)$

Answer.  $\frac{3}{2}$

40.  $\cos(\alpha + \beta)$

Answer.


42.  $\sin(\alpha - \beta)$

Answer.  $\frac{\sqrt{10}}{5} \cdot \frac{2\sqrt{13}}{13} - \frac{\sqrt{15}}{5} \cdot \frac{3\sqrt{13}}{13} = \frac{2\sqrt{130} - 3\sqrt{195}}{65}$

44.  $\tan(\alpha - \beta)$

Answer.  $\frac{\frac{\sqrt{6}}{2} - \frac{3}{2}}{1 + \frac{\sqrt{6}}{2} \cdot \frac{3}{2}} = \frac{30 - 13\sqrt{6}}{19}$

**Exercise Group.** Find the exact value of each expression given



45.  $\sin(\alpha)$

**Answer.**  $\frac{3\sqrt{34}}{34}$

47.  $\tan(\alpha)$

**Answer.**  $\frac{3}{5}$

49.  $\cos(\beta)$

**Answer.**  $\frac{7}{8}$

51.  $\sin(\alpha + \beta)$

**Answer.**  $\frac{3\sqrt{34}}{34} \cdot \frac{7}{8} + \frac{5\sqrt{34}}{34} \cdot \frac{\sqrt{15}}{8} = \frac{21\sqrt{34} + 5\sqrt{510}}{272}$

53.  $\tan(\alpha + \beta)$

**Answer.**  $\frac{\frac{3}{5} + \frac{\sqrt{15}}{7}}{1 - \frac{3}{5} \cdot \frac{\sqrt{15}}{7}} = \frac{480 + 119\sqrt{15}}{545}$

55.  $\cos(\alpha - \beta)$

**Answer.**  $\frac{5\sqrt{34}}{34} \cdot \frac{7}{8} + \frac{3\sqrt{34}}{34} \cdot \frac{\sqrt{15}}{8} = \frac{35\sqrt{34} + 3\sqrt{510}}{272}$

46.  $\cos(\alpha)$

**Answer.**  $\frac{5\sqrt{34}}{34}$

48.  $\sin(\beta)$

**Answer.**  $\frac{\sqrt{15}}{8}$

50.  $\tan(\beta)$

**Answer.**  $\frac{\sqrt{15}}{7}$

52.  $\cos(\alpha + \beta)$

**Answer.**  $\frac{5\sqrt{34}}{34} \cdot \frac{7}{8} - \frac{3\sqrt{34}}{34} \cdot \frac{\sqrt{15}}{8} = \frac{35\sqrt{34} - 3\sqrt{510}}{272}$

54.  $\sin(\alpha - \beta)$

**Answer.**  $\frac{3\sqrt{34}}{34} \cdot \frac{7}{8} - \frac{5\sqrt{34}}{34} \cdot \frac{\sqrt{15}}{8} = \frac{21\sqrt{34} - 5\sqrt{510}}{272}$

56.  $\tan(\alpha - \beta)$

**Answer.**  $\frac{\frac{3}{5} - \frac{\sqrt{15}}{7}}{1 + \frac{3}{5} \cdot \frac{\sqrt{15}}{7}} = \frac{480 - 119\sqrt{15}}{545}$

**Exercise Group.** Find the exact of each expression given  $\sin \alpha = \frac{20}{29}$ ,  $0 < \alpha < \frac{\pi}{2}$  and  $\cos \beta = \frac{24}{25}$ ,  $0 < \beta < \frac{\pi}{2}$

57.  $\cos(\alpha)$

**Answer.**  $\frac{21}{29}$

59.  $\sin(\beta)$

**Answer.**  $\frac{7}{25}$

61.  $\sin(\alpha + \beta)$

**Answer.**  $\frac{627}{725}$

63.  $\cos(\alpha + \beta)$

**Answer.**  $\frac{364}{725}$

65.  $\tan(\alpha + \beta)$

**Answer.**  $\frac{627}{364}$

58.  $\tan(\alpha)$

**Answer.**  $\frac{20}{29}$

60.  $\tan(\beta)$

**Answer.**  $\frac{7}{24}$

62.  $\sin(\alpha - \beta)$

**Answer.**  $\frac{333}{725}$

64.  $\cos(\alpha - \beta)$

**Answer.**  $\frac{644}{725}$

66.  $\tan(\alpha - \beta)$

**Answer.**  $\frac{333}{644}$

**Exercise Group.** Find the exact of each expression given  $\tan \alpha = \frac{8}{15}$ ,  $\pi < \alpha < \frac{3\pi}{2}$  and  $\cos \beta = -\frac{3}{5}$ ,  $\frac{\pi}{2} < \beta < \pi$

67.  $\sin(\alpha)$

**Answer.**  $-\frac{8}{17}$

69.  $\sin(\beta)$

**Answer.**  $\frac{4}{5}$

71.  $\sin(\alpha + \beta)$

**Answer.**  $-\frac{36}{85}$

73.  $\cos(\alpha + \beta)$

**Answer.**  $\frac{77}{85}$

75.  $\tan(\alpha + \beta)$

**Answer.**  $-\frac{36}{77}$

68.  $\cos(\alpha)$

**Answer.**  $-\frac{15}{17}$

70.  $\tan(\beta)$

**Answer.**  $-\frac{4}{3}$

72.  $\sin(\alpha - \beta)$

**Answer.**  $\frac{84}{85}$

74.  $\cos(\alpha - \beta)$

**Answer.**  $\frac{13}{85}$

76.  $\tan(\alpha - \beta)$

**Answer.**  $\frac{84}{13}$

**Exercise Group.** Verify the identity.

$$77. \sin\left(\theta + \frac{\pi}{2}\right) = \cos \theta$$

$$79. \cos(\theta - \pi) = -\cos \theta$$

$$81. \tan\left(\frac{\pi}{4} - \theta\right) = \frac{1 - \tan \theta}{1 + \tan \theta}$$

$$83. \cos\left(\theta + \frac{\pi}{3}\right) = -\sin\left(x - \frac{\pi}{6}\right)$$

$$85. \frac{\sin(x+y)}{\cos x \cos y} = \tan x + \tan y$$

$$87. \sin(x+y) - \sin(x-y) = 2 \cos x \sin y$$

$$78. \sin(\theta - \pi) = -\sin \theta$$

$$80. \tan(\theta - \pi) = \tan \theta$$

$$82. \sin\left(\frac{\pi}{2} - \theta\right) = \sin\left(\frac{\pi}{2} + \theta\right)$$

$$84. \cos(x+y) \cos(x-y) = \cos^2 x - \sin^2 y$$

$$86. \cot(x-y) = \frac{\cot x \cot y + 1}{\cot y - \cot x}$$

$$88. \cos(x+y) + \cos(x-y) = 2 \cos x \cos y$$

**Exercise Group.** Write each expression in terms on sine only. Round your angles to one decimal.

$$89. -\frac{\sqrt{2}}{2} \sin x - \frac{\sqrt{2}}{2} \cos x$$

$$\text{Answer. } \sin(x + 225^\circ)$$

$$91. \frac{1}{2} \sin x + \frac{\sqrt{3}}{2} \cos x$$

$$\text{Answer. } \sin(x + 60^\circ)$$

$$93. 3 \sin x + 7 \cos x$$

$$\text{Answer. } \sqrt{58} \sin(x + 66.8^\circ)$$

$$95. 8 \sin x - 2 \cos x$$

$$\text{Answer. } \sqrt{68} \sin(x + 346.0^\circ)$$

$$90. -\frac{\sqrt{3}}{2} \sin x - \frac{1}{2} \cos x$$

$$\text{Answer. } \sin(x + 210^\circ)$$

$$92. -\frac{\sqrt{3}}{2} \sin x - \frac{1}{2} \cos x$$

$$\text{Answer. } \sin(x + 210^\circ)$$

$$94. -5 \sin x - 9 \cos x$$

$$\text{Answer. } \sqrt{106} \sin(x + 240.9^\circ)$$

$$96. -7 \sin x + 4 \cos x$$


$$\text{Answer. } \sqrt{65} \sin(x + 150.3^\circ)$$

97. Use the Subtraction Formula for Cosine to prove the Cofunction Identity for Sine:  $\sin \theta = \cos\left(\frac{\pi}{2} - \theta\right)$ .



### 3.3 Double-Angle and Half-Angle Formulas

Suppose we want to accurately position the Hawaiian Star Compass on the Unit Circle. In [Figure 1.1.4](#), the house for Manu is located at halfway between Hikina and 'Ākau, resulting in an angle of  $45^\circ$ . By applying right triangle trigonometry, we can determine the exact coordinates of Manu as  $(\cos(45^\circ), \sin(45^\circ)) = \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$ . However, as we move to the house of 'Āina, located halfway between Manu and Hikina, we encounter a problem. The angle for 'Āina is  $22.5^\circ$ , which is not explicitly listed in [Table 1.5.18](#). Therefore, we must resort to a calculator for numerical approximations.



In this section, we will learn about the double and half-angle formulas for trigonometry. These formulas allow us to determine exact trigonometric function values for angles that are double or half of known values. This will enable us to use our existing knowledge of trigonometric functions at  $45^\circ$  and apply the half-angle formulas to obtain exact values at  $22.5^\circ$ .

#### 3.3.1 Double-Angle Formulas

Recall the addition formula for sine:

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta \quad (3.3.1)$$

Consider the case when the two angles are equal. We will call this angle  $\theta$ , so let  $\alpha = \theta$  and  $\beta = \theta$ . Then [Eq \(3.3.1\)](#) becomes

$$\begin{aligned} \sin(\theta + \theta) &= \sin \theta \cos \theta + \cos \theta \sin \theta \\ \sin(2\theta) &= 2 \sin \theta \cos \theta \end{aligned}$$

Thus we obtain a formula for sine of twice the angle  $\theta$ .

**Definition 3.3.1 Double-Angle Formulas.**

$$\begin{aligned} \sin 2\theta &= 2 \sin \theta \cos \theta \\ \cos 2\theta &= \cos^2 \theta - \sin^2 \theta \\ \cos 2\theta &= 1 - 2 \sin^2 \theta \\ \cos 2\theta &= 2 \cos^2 \theta - 1 \\ \tan 2\theta &= \frac{2 \tan \theta}{1 - \tan^2 \theta} \end{aligned}$$

◇


The proofs for the Double-Angle Formulas for Cosine and Tangent are left as exercises ([Exercise 3.3.4.6-Exercise 3.3.4.9](#)).

**Remark 3.3.2** Notice that there are three variations of the double-angle formula for cosine. All three equations give the correct answer; however, one version may be more convenient depending on the given information. For example, if you are given the value of  $\sin \theta$ , it may be easier to select the version that solely involves  $\sin \theta$  and does not include  $\cos \theta$ .

**Example 3.3.3** Given  $\sin \theta = -\frac{5}{13}$  and  $\theta$  lies in Quadrant III, find the exact value of

(a)  $\sin(2\theta)$

**Solution.** By the double-angle formula, we have  $\sin(2\theta) = 2 \sin \theta \cos \theta$ . We are given the value of  $\sin \theta$ , but we do not have  $\cos \theta$ . To find  $\cos \theta$ , we will draw the triangle formed from  $\sin \theta = -\frac{5}{13}$  where  $\theta$  lies in Quadrant III.



Using the Pythagorean Theorem, we can solve the triangle:

$$\begin{aligned}x^2 + (-5)^2 &= 13^2 \\x^2 + 25 &= 169 \\x^2 &= 144 \\x &= 12\end{aligned}$$

Thus we have

$$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}} = -\frac{12}{13}$$

and

$$\tan \theta = \frac{\text{opposite}}{\text{adjacent}} = \frac{5}{12}.$$

With this new information, we can use the double-angle formula to find  $\sin(2\theta)$ :

$$\sin(2\theta) = 2 \sin \theta \cos \theta = 2 \left( -\frac{5}{13} \right) \left( -\frac{12}{13} \right) = \frac{120}{169}$$

(b)  $\cos(2\theta)$

**Solution.** To compute  $\cos 2\theta$ , notice there are three different formulas:  $\cos 2\theta = \cos^2 \theta - \sin^2 \theta$ ,  $\cos 2\theta = 1 - 2\sin^2 \theta$ , or  $\cos 2\theta = 2\cos^2 \theta - 1$ . Using any of the three equations will give us the correct answer. However, given that we know  $\sin \theta = -\frac{5}{13}$ , it may be easier to use  $\cos 2\theta = 1 - 2\sin^2 \theta$ , since the other two equations require us to know  $\cos \theta$ .

Without having to draw the triangle, we could get

$$\begin{aligned}\cos 2\theta &= 1 - 2\sin^2 \theta \\ &= 1 - 2\left(-\frac{5}{13}\right)^2 \\ &= 1 - 2\left(\frac{25}{169}\right) \\ &= \frac{169}{169} - \frac{50}{169} \\ &= \frac{119}{169}\end{aligned}$$

(c)  $\tan(2\theta)$

**Solution.** Using the double-angle formula for tangent, we get

$$\begin{aligned}\tan 2\theta &= \frac{2 \tan \theta}{1 - \tan^2 \theta} \\ &= \frac{2\left(\frac{5}{12}\right)}{1 - \left(\frac{5}{12}\right)^2} \\ &= \frac{\frac{10}{12}}{1 - \frac{25}{144}} \\ &= \frac{\frac{10}{12}}{\frac{119}{144}} \\ &= \frac{10}{12} \cdot \frac{144}{119} \\ &= \frac{120}{119}\end{aligned}$$

□

**Example 3.3.4** Write  $\cos(3\theta)$  in terms of  $\sin \theta$ .

**Solution.**

$$\begin{aligned}\sin(3\theta) &= \sin(2\theta + \theta) \\ &= \sin(2\theta) \cos \theta + \cos(2\theta) \sin \theta && \text{addition formula} \\ &= (2 \sin \theta \cos \theta) \cos \theta + (\cos^2 \theta - \sin^2 \theta) \sin \theta && \text{double-angle formula} \\ &= 2 \sin \theta \cos^2 \theta + \sin \theta \cos^2 \theta - \sin^3 \theta \\ &= 3 \sin \theta \cos^2 \theta - \sin^3 \theta \\ &= 3 \sin \theta(1 - \sin^2 \theta) - \sin^3 \theta && \text{Pythagorean Identity} \\ &= 3 \sin \theta - 3 \sin^3 \theta - \sin^3 \theta \\ &= 3 \sin \theta - 4 \sin^3 \theta\end{aligned}$$

□

### 3.3.2 Reducing Powers Formulas

You may notice that the double-angle formula for cosine expresses a trigonometric function in terms of the square of another trigonometric function. By rearranging the terms, we can derive formulas for reducing the powers of sine, cosine, and tangent expressions with even powers to terms involving only cosine. These formulas are particularly useful in calculus.

**Definition 3.3.5 Formulas for Reducing Powers.**

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

$$\tan^2 \theta = \frac{1 - \cos(2\theta)}{1 + \cos(2\theta)}$$

◇

*Proof.* To prove the first formula, solve for  $\sin^2 \theta$  in the double-angle formula:  $\cos 2\theta = 1 - 2\sin^2 \theta$ . The second formula is obtained similarly by solving for  $\cos^2 \theta$  in the formula:  $\cos 2\theta = 2\cos^2 \theta - 1$ . The first two formula can be used to obtain the third formula:

$$\tan^2 \theta = \frac{\sin^2 \theta}{\cos^2 \theta} = \frac{\frac{1 - \cos(2\theta)}{2}}{\frac{1 + \cos(2\theta)}{2}} = \frac{1 - \cos(2\theta)}{1 + \cos(2\theta)}$$

■

**Example 3.3.6** Write  $\sin^4 \theta$  as an expression that does not involve powers of sine or cosine greater than 1.

**Solution.** We will use the Reducing Powers Formula twice.

$$\begin{aligned} \sin^4 \theta &= (\sin^2 \theta)^2 \\ &= \left( \frac{1 - \cos(2\theta)}{2} \right)^2 && \text{reducing powers} \\ &= \frac{1}{4} (1 - 2\cos(2\theta) + \cos^2(2\theta)) \\ &= \frac{1}{4} \left( 1 - 2\cos(2\theta) + \frac{1 + \cos(4\theta)}{2} \right) && \text{reducing powers} \\ &= \frac{1}{4} \left( 1 - 2\cos(2\theta) + \frac{1}{2} + \frac{\cos(4\theta)}{2} \right) \\ &= \frac{1}{4} \left( \frac{3}{2} - \frac{4}{2}\cos(2\theta) + \frac{1}{2}\cos(4\theta) \right) \\ &= \frac{1}{8} (3 - 4\cos(2\theta) + \cos(4\theta)) \end{aligned}$$

□

### 3.3.3 Half-Angle Formulas

Another set of useful formulas are the half-angle formulas.

**Definition 3.3.7 Half-Angle Formulas.**

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}, \quad \cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}, \quad \tan \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}}$$

The choice of the + or - sign depends on the Quadrant in which  $\theta/2$  lies.  $\diamond$

*Proof.* We take the square root on both sides of the Formulas for Reducing Powers (Definition 3.3.5) and halve the angle ( $\theta$  becomes  $\frac{\theta}{2}$  and  $2\theta$  becomes  $\theta$ ) to arrive at our formulas.  $\blacksquare$

**Example 3.3.8 Locating ‘Āina.** We are now ready to revisit the problem posed at the start of this section when we were asked to determine the exact coordinates of the house ‘Āina on the Unit Circle.

**Solution.** We know the coordinates are at

$$(\cos 22.5^\circ, \sin 22.5^\circ).$$

To find the exact value of  $\cos 22.5^\circ$ , we will use the half-angle formula:

$$\begin{aligned} \cos 22.5^\circ &= \cos \left( \frac{45}{2} \right)^\circ \\ &= \sqrt{\frac{1 + \cos 45^\circ}{2}} \\ &= \sqrt{\frac{1 + \frac{\sqrt{2}}{2}}{2}} \\ &= \sqrt{\frac{\frac{2}{2} + \frac{\sqrt{2}}{2}}{2}} \\ &= \sqrt{\frac{2 + \sqrt{2}}{4}} \\ &= \frac{1}{2} \cdot \sqrt{2 + \sqrt{2}} \end{aligned}$$

Since the half-angle formula has  $\pm$ , we check the quadrant. In this case, our angle is  $22.5^\circ$ , which is in Quadrant I. Therefore, we choose the positive value.

Finding the exact value of  $\sin 22.5^\circ$  is left for [Exercise 3.3.4.1](#).  $\square$

**Example 3.3.9** Given  $\sin \theta = -\frac{5}{13}$  and  $\theta$  lies in Quadrant III, find the exact value of

(a)  $\sin \frac{\theta}{2}$

**Solution.** Notice the Half-Angle Formulas all require us to know  $\cos \theta$ . Since the given information describes the same triangle in [Example 3.3.3](#), we refer to that problem to get  $\cos \theta = -\frac{12}{13}$ .

Next, since  $\theta$  is in Quadrant III,  $180^\circ < \theta < 270^\circ$ , so dividing by 2 gives us  $\frac{180^\circ}{2} < \frac{\theta}{2} < \frac{270^\circ}{2}$  or  $90^\circ < \frac{\theta}{2} < 135^\circ$ . Therefore, we conclude that  $\frac{\theta}{2}$  lies in Quadrant II.

To calculate  $\sin \frac{\theta}{2}$ , we first note that because  $\frac{\theta}{2}$  lies in Quadrant II,  $\sin \frac{\theta}{2} > 0$  so we will choose the positive (+) sign in the Half-Angle Formula:

$$\sin \frac{\theta}{2} = \sqrt{\frac{1 - \cos \theta}{2}}$$

$$\begin{aligned}
&= \sqrt{\frac{1 - \left(-\frac{12}{13}\right)}{2}} \\
&= \sqrt{\frac{1 + \frac{12}{13}}{2}} \\
&= \sqrt{\frac{\frac{13}{13} + \frac{12}{13}}{2}} \\
&= \sqrt{\frac{\frac{25}{13}}{2}} \\
&= \sqrt{\frac{25}{26}}
\end{aligned}$$

(b)  $\cos \frac{\theta}{2}$

**Solution.** Since  $\frac{\theta}{2}$  is in Quadrant II, we know that  $\cos \frac{\theta}{2} < 0$  so we will choose the negative (-) sign in the Half-Angle Formula:

$$\begin{aligned}
\cos \frac{\theta}{2} &= -\sqrt{\frac{1 + \cos \theta}{2}} \\
&= -\sqrt{\frac{1 + \left(-\frac{12}{13}\right)}{2}} \\
&= -\sqrt{\frac{1 - \frac{12}{13}}{2}} \\
&= -\sqrt{\frac{\frac{13}{13} - \frac{12}{13}}{2}} \\
&= -\sqrt{\frac{\frac{1}{13}}{2}} \\
&= -\sqrt{\frac{1}{26}}
\end{aligned}$$

(c)  $\tan \frac{\theta}{2}$

**Solution.** Since  $\frac{\theta}{2}$  is in Quadrant II, we know that  $\tan \frac{\theta}{2} < 0$  so we will choose the negative (-) sign in the Half-Angle Formula:

$$\begin{aligned}
\tan \frac{\theta}{2} &= -\sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}} \\
&= -\sqrt{\frac{1 - \left(-\frac{12}{13}\right)}{1 + \left(-\frac{12}{13}\right)}} \\
&= -\sqrt{\frac{1 + \frac{12}{13}}{1 - \frac{12}{13}}} \\
&= -\sqrt{\frac{\frac{13}{13} + \frac{12}{13}}{\frac{13}{13} - \frac{12}{13}}} \\
&= -\sqrt{\frac{\frac{25}{13}}{\frac{1}{13}}}
\end{aligned}$$

$$\begin{aligned}
&= -\sqrt{\frac{25}{13} \cdot \frac{13}{1}} \\
&= -\sqrt{25} \\
&= -5
\end{aligned}$$

□

We can derive another formula for  $\tan \frac{\theta}{2}$  that does not involve  $\pm$ :

**Definition 3.3.10 Half-Angle Formulas for Tangent.**

$$\tan \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}} = \frac{1 - \cos \theta}{\sin \theta} = \frac{\sin \theta}{1 + \cos \theta}$$

◇

*Proof.* We begin by first multiplying both sides of the Sine formula for Reducing Powers by 2 and halving the angle:

$$1 - \cos \theta = 2 \sin^2 \left( \frac{\theta}{2} \right)$$

and applying the double-angle formula to:

$$\sin \theta = \sin 2 \cdot \frac{\theta}{2} = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}$$

Dividing the two preceding results

$$\frac{1 - \cos \theta}{\sin \theta} = \frac{2 \sin^2 \left( \frac{\theta}{2} \right)}{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}} = \frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}} = \tan \frac{\theta}{2}$$

Thus

$$\tan \frac{\theta}{2} = \frac{1 - \cos \theta}{\sin \theta}$$

Similarly, it can be shown that

$$\tan \frac{\theta}{2} = \frac{\sin \theta}{1 + \cos \theta}$$

■

**Example 3.3.11** Calculate  $\tan \frac{\theta}{2}$  from [Example 3.3.9](#) using the above formula.

**Solution.**


$$\tan \frac{\theta}{2} = \frac{1 - \cos \theta}{\sin \theta} = \frac{1 - \left( -\frac{12}{13} \right)}{-\frac{5}{13}} = \frac{\frac{25}{13}}{-\frac{5}{13}} = \frac{25}{13} \cdot \left( -\frac{13}{5} \right) = -5$$

□

Note: We obtained the same result for  $\tan \frac{\theta}{2}$  as we did in [Example 3.3.9](#). In this example, we did not have to determine if  $\tan \frac{\theta}{2}$  was positive or negative, however, we do need to know the values of both  $\sin \theta$  and  $\cos \theta$ .

### 3.3.4 Exercises

**Exercise Group.** The house 'Āina is located at  $22.5^\circ = \frac{45^\circ}{2}$  and the house Nā Leo is located at  $67.5^\circ = \frac{135^\circ}{2}$ . Use the half-angle formulas to evaluate the exact value of the given expression at each of these houses.



1.  $\sin(22.5^\circ)$   
**Answer.**  $\frac{1}{2} \cdot \sqrt{2 - \sqrt{2}}$
2.  $\tan(22.5^\circ)$   
**Answer.**  $\sqrt{3 - 2\sqrt{2}} = \sqrt{2} - 1$
3.  $\sin(67.5^\circ)$   
**Answer.**  $\frac{1}{2} \sqrt{2 + \sqrt{2}}$
4.  $\cos(67.5^\circ)$   
**Answer.**  $\frac{1}{2} \sqrt{2 - \sqrt{2}}$
5.  $\tan(67.5^\circ)$   
**Answer.**  $\sqrt{3 + 2\sqrt{2}} = \sqrt{2} + 1$
6. Use the Addition Formula,  $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$ , to prove the double angle formula for cosine:  

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta.$$
7. Use the Pythagorean Identity ( $\sin^2 \theta + \cos^2 \theta = 1$ ) and the result from [Exercise 3.3.4.6](#) to prove  

$$\cos 2\theta = 1 - 2\sin^2 \theta.$$
8. Use the Pythagorean Identity ( $\sin^2 \theta + \cos^2 \theta = 1$ ) and the result from [Exercise 3.3.4.6](#) to prove  


$$\cos 2\theta = 2\cos^2 \theta - 1.$$
9. Use the addition formula,  $\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$ , to prove the double



angle formula for tangent:

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}.$$

**Exercise Group.** Use the figure below to find the exact values for each of the following exercises.



10.  $\sin(\theta)$

**Answer.**  $\frac{5}{13}$

12.  $\tan(\theta)$

**Answer.**  $\frac{5}{12}$

14.  $\cos(2\theta)$

**Answer.**  $\frac{119}{169}$

16.  $\sin\left(\frac{\theta}{2}\right)$

**Answer.**  $\sqrt{\frac{1}{26}}$

18.  $\tan\left(\frac{\theta}{2}\right)$

**Answer.**  $\frac{1}{5}$

11.  $\cos(\theta)$

**Answer.**  $\frac{12}{13}$

13.  $\sin(2\theta)$

**Answer.**  $\frac{120}{169}$


15.  $\tan(2\theta)$

**Answer.**  $\frac{120}{119}$

17.  $\cos\left(\frac{\theta}{2}\right)$

**Answer.**  $\sqrt{\frac{25}{26}}$

**Exercise Group.** Use the figure below to find the exact values for each of the following exercises.



19.  $\sin(\theta)$

**Answer.**  $\frac{4}{5}$

21.  $\tan(\theta)$

**Answer.**  $\frac{4}{3}$

23.  $\cos(2\theta)$

**Answer.**  $-\frac{7}{25}$

25.  $\sin\left(\frac{\theta}{2}\right)$

**Answer.**  $\sqrt{\frac{1}{5}}$

20.  $\cos(\theta)$

**Answer.**  $\frac{3}{5}$

22.  $\sin(2\theta)$

**Answer.**  $\frac{24}{25}$

24.  $\tan(2\theta)$

**Answer.**  $-\frac{24}{7}$

26.  $\cos\left(\frac{\theta}{2}\right)$

**Answer.**  $\sqrt{\frac{4}{5}}$

27.  $\tan\left(\frac{\theta}{2}\right)$

Answer.  $\frac{1}{2}$

**Exercise Group.** Find the exact value of each expression given  $\cos\theta = -\frac{3}{5}$  and  $\theta$  is in Quadrant III.

28.  $\sin(\theta)$

Answer.  $-\frac{4}{5}$

30.  $\sin(2\theta)$

Answer.  $\frac{24}{25}$

32.  $\tan(2\theta)$

Answer.  $-\frac{24}{7}$

34.  $\cos\left(\frac{\theta}{2}\right)$

Answer.  $-\sqrt{\frac{1}{5}}$

29.  $\tan(\theta)$

Answer.  $\frac{4}{3}$

31.  $\cos(2\theta)$

Answer.  $-\frac{7}{25}$

33.  $\sin\left(\frac{\theta}{2}\right)$

Answer.  $\sqrt{\frac{4}{5}}$

35.  $\tan\left(\frac{\theta}{2}\right)$

Answer.  $-2$

**Exercise Group.** Find the exact value of each expression given  $\sin\theta = -\frac{8}{17}$  and  $270^\circ < \theta < 360^\circ$ .

36.  $\cos(\theta)$

Answer.  $\frac{15}{17}$

38.  $\sin(2\theta)$

Answer.  $-\frac{240}{289}$

40.  $\tan(2\theta)$

Answer.  $-\frac{240}{161}$

42.  $\cos\left(\frac{\theta}{2}\right)$

Answer.  $\sqrt{\frac{16}{17}}$

37.  $\tan(\theta)$

Answer.  $-\frac{8}{15}$

39.  $\cos(2\theta)$

Answer.  $\frac{161}{289}$

41.  $\sin\left(\frac{\theta}{2}\right)$

Answer.  $\sqrt{\frac{1}{17}}$

43.  $\tan\left(\frac{\theta}{2}\right)$

Answer.  $-\frac{1}{4}$

**Exercise Group.** Find the exact value of each expression given  $\cos\theta = \frac{2}{3}$  and  $\frac{3\pi}{2} < \theta < 2\pi$ .

44.  $\sin(\theta)$

Answer.  $-\frac{\sqrt{5}}{3}$

46.  $\sin(2\theta)$

Answer.  $-\frac{4\sqrt{5}}{9}$

48.  $\tan(2\theta)$

Answer.  $-4\sqrt{5}$

50.  $\cos\frac{\theta}{2}$

Answer.  $-\sqrt{\frac{5}{6}}$

45.  $\tan(\theta)$

Answer.  $\frac{\sqrt{5}}{2}$

47.  $\cos(2\theta)$

Answer.  $-\frac{1}{9}$

49.  $\sin\frac{\theta}{2}$

Answer.  $\sqrt{\frac{1}{6}}$

51.  $\tan\frac{\theta}{2}$

Answer.  $-\sqrt{\frac{1}{5}}$

**Exercise Group.** Use the Half-Angle Formula to find the exact value of each of the following

52.  $\tan 157.5^\circ$

Answer.  $1 - \sqrt{2}$

53.  $\sin 75^\circ$

Answer.  $\sqrt{\frac{2+\sqrt{3}}{4}}$

54.  $\tan 112.5^\circ$

Answer.  $-1 - \sqrt{2}$

56.  $\cos \frac{\pi}{8}$

Answer.  $\cos \frac{\pi}{8} = \sqrt{\frac{\sqrt{2}+2}{4}}$

58.  $\sin \frac{7\pi}{12}$

Answer.  $\sin \frac{7\pi}{12} = \frac{\sqrt{2}-1}{2}$

55.  $\cos 15^\circ$

Answer.  $\frac{\sqrt{2+\sqrt{3}}}{2}$

57.  $\tan \frac{11\pi}{12}$

Answer.  $\tan \frac{11\pi}{12} = -\sqrt{2} + 1$

59.  $\cos \frac{3\pi}{8}$

Answer.  $\cos \frac{3\pi}{8} = \sqrt{\frac{\sqrt{2}+2}{4}}$

**Exercise Group.** Write each of the following as expressions that do not involve powers of sine or cosine greater than 1.

60.  $\cos^4 \theta$

Answer.  $\frac{3}{8} + \frac{1}{2} \cos(2\theta) + \frac{1}{8} \cos(4\theta)$

62.  $\sin^2 \theta \cos^4 \theta$

Answer.  $\frac{1}{16}(1 + \cos(2\theta) - \cos(4\theta) - \cos(2\theta) \cos(4\theta))$

61.  $\sin^3(2\theta)$

Answer.  $\frac{1}{2} \sin(2\theta) \cdot (1 - \cos(4\theta))$

63.  $\sin^4 \theta \cos^2 \theta$

Answer.  $\frac{1}{16}(1 - \cos(2\theta) - \cos(4\theta) + \cos(2\theta) \cos(4\theta))$

64. Write  $\sin^2 \theta \cos^2 \theta$  expressions that does not involve powers of sine or cosine greater than 1.

(a) Using the Reducing Powers Formula (Definition 3.3.5)

Answer.  $\frac{1}{8}(1 - \cos(4\theta))$

(b) Using the Double Angle Formula (Definition 3.3.1) where  $\sin \theta \cos \theta = \frac{1}{2} \sin 2\theta$

Answer.  $\frac{1}{8}(1 - \cos(4\theta))$

65. Find the exact value of  $\sin^2(15^\circ)$

(a) Evaluate using the Reducing Powers Formula (Definition 3.3.5)


Answer.  $\frac{2-\sqrt{3}}{4}$

(b) Evaluate using the Half-Angle Formula (Definition 3.3.7)

Answer.  $\frac{2-\sqrt{3}}{4}$


**Exercise Group.** Use half angle to find the exact deviation for the indicated angle,  $\theta$ .

66. 2 Hours ( $\theta = 22.5^\circ$ )



Answer.  $60\sqrt{2 - \sqrt{2}}$

67. 6 Houses ( $\theta = 67.5^\circ$ )



Answer.  $60\sqrt{2 + \sqrt{2}}$

**Exercise Group.** Verify the identity

68.  $(\sin \theta + \cos \theta)^2 = 1 + \sin(2\theta)$

70.  $(\sin^2 \theta - 1)^2 = \cos(2\theta) + \sin^4 \theta$

72.  $\cos^4 \theta - \sin^4 \theta = \cos(2\theta)$

74.  $\cot(2\theta) = \frac{1 - \tan^2 \theta}{2 \tan \theta}$

76.  $\sec^2\left(\frac{\theta}{2}\right) = \frac{2}{1 + \cos \theta}$

69.  $\cos(2\theta) = \frac{\cot \theta - \tan \theta}{\cot \theta + \tan \theta}$

71.  $\cos^2(3\theta) - \sin^2(3\theta) = \cos(6\theta)$

73.  $\sin(6\theta) = 2 \sin(3\theta) \cos(3\theta)$

75.  $\csc^2\left(\frac{\theta}{2}\right) = \frac{2}{1 - \cos \theta}$

77.  $\frac{2 \tan \theta}{1 + \tan^2 \theta} = \sin(2\theta)$

### 3.4 Product-to-Sum and Sum-to-Product Formulas

In this section, we will learn how to convert sums of trigonometric functions to products of trigonometric functions, and vice versa. These techniques provide us with tools to simplify expressions and solve equations.

#### 3.4.1 Product to Sum Formulas

**Definition 3.4.1** Product to Sum Formulas.

$$\sin \alpha \cos \beta = \frac{1}{2}[\sin(\alpha + \beta) + \sin(\alpha - \beta)]$$

$$\cos \alpha \sin \beta = \frac{1}{2}[\sin(\alpha + \beta) - \sin(\alpha - \beta)]$$

$$\cos \alpha \cos \beta = \frac{1}{2}[\cos(\alpha + \beta) + \cos(\alpha - \beta)]$$

$$\sin \alpha \sin \beta = \frac{1}{2}[\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$

◇

*Proof.* We add the addition and subtraction formulas for cosine:

$$\begin{array}{r} \cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \\ + \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta \\ \hline \end{array}$$

$$\cos(\alpha + \beta) + \cos(\alpha - \beta) = 2 \cos \alpha \cos \beta$$

Dividing both sides by 2, we get

$$\frac{1}{2}[\cos(\alpha + \beta) + \cos(\alpha - \beta)] = \cos \alpha \cos \beta$$

Next, we add the addition and subtraction formulas for sine:

$$\begin{array}{r} \sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta \\ + \sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta \\ \hline \end{array}$$

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Dividing both sides by 2, we get

$$\frac{1}{2}[\sin(\alpha + \beta) + \sin(\alpha - \beta)] = \sin \alpha \cos \beta$$

Next, we subtract the addition and subtraction formulas for sine:

$$\begin{array}{r} \sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta \\ - \sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta \\ \hline \end{array}$$

$$\sin(\alpha + \beta) - \sin(\alpha - \beta) = 2 \sin \beta \cos \alpha$$

Dividing both sides by 2, we get

$$\frac{1}{2}[\sin(\alpha + \beta) - \sin(\alpha - \beta)] = \sin \beta \cos \alpha$$

Finally, we subtract the addition and subtraction formulas for cosine:

$$\begin{array}{r} \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta \\ - \cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \\ \hline \end{array}$$

$$\cos(\alpha - \beta) - \cos(\alpha + \beta) = 2 \sin \alpha \sin \beta$$

Dividing both sides by 2, we get

$$\frac{1}{2}[\cos(\alpha - \beta) - \cos(\alpha + \beta)] = \sin \alpha \sin \beta$$

**Example 3.4.2** Express the product of  $\cos(2x) \cos(5x)$  as a sum or difference of sine and cosine with no products. ■

**Solution.** Using the formula we get

$$\begin{aligned} \cos(2x) \cos(5x) &= \frac{1}{2}[\cos(2x + 5x) + \cos(2x - 5x)] \\ &= \frac{1}{2}[\cos(7x) + \cos(-3x)] \end{aligned}$$

This satisfies the requirement of expressing the product of  $\cos(2x) \cos(5x)$  as a sum or difference of sine and cosine with no products. However, we can simplify it further.

Since cosine is an even function,  $\cos(-3x) = \cos(3x)$ . Thus, we can simplify the expression to:

$$\cos(2x) \cos(5x) = \frac{1}{2}[\cos(7x) + \cos(3x)]$$

□

**Example 3.4.3** Express the product of  $\sin(6\theta) \cos(4\theta)$  as a sum or difference of sine and cosine with no products.

**Solution.** Using the formula we get

$$\begin{aligned} \sin(6\theta) \cos(4\theta) &= \frac{1}{2}[\sin(6\theta + 4\theta) + \sin(6\theta - 4\theta)] \\ &= \frac{1}{2}[\sin(10\theta) + \cos(2\theta)] \end{aligned}$$

□

**Remark 3.4.4 Negative Angles.** In the previous example, both forms  $\cos(2x) \cos(5x) = \frac{1}{2}[\cos(7x) + \cos(-3x)]$  and  $\cos(2x) \cos(5x) = \frac{1}{2}[\cos(7x) + \cos(3x)]$  are valid representations of the answer. However, it is more common to write the first form (where the angles are positive) because it simplifies the expression and aligns with standard conventions for representing trigonometric identities. Positive angles are often preferred for clarity and consistency in mathematical notation. Negative angles can be transformed into positive angles using the even-odd properties of trigonometric functions ([Definition 1.5.22](#)).

### 3.4.2 Sum to Product Formulas

**Definition 3.4.5 Sum-to-Product Formula.**

$$\begin{aligned}\sin \alpha + \sin \beta &= 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right) \\ \sin \alpha - \sin \beta &= 2 \cos \left( \frac{\alpha + \beta}{2} \right) \sin \left( \frac{\alpha - \beta}{2} \right) \\ \cos \alpha + \cos \beta &= 2 \cos \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right) \\ \cos \alpha - \cos \beta &= -2 \sin \left( \frac{\alpha + \beta}{2} \right) \sin \left( \frac{\alpha - \beta}{2} \right)\end{aligned}$$

◇

*Proof.* We first let  $\alpha = \frac{u+v}{2}$  and  $\beta = \frac{u-v}{2}$ . Then

$$\alpha + \beta = \frac{u+v}{2} + \frac{u-v}{2} = \frac{2u}{2} = u$$

and

$$\alpha - \beta = \frac{u+v}{2} - \frac{u-v}{2} = \frac{2v}{2} = v$$

Substituting these values for  $\alpha$ ,  $\beta$ ,  $\alpha + \beta$ , and  $\alpha - \beta$  into the Product-to-Sum Formulas, we get

$$\begin{aligned}\sin \left( \frac{u+v}{2} \right) \cos \left( \frac{u-v}{2} \right) &= \frac{1}{2} [\sin(u) + \sin(v)] \\ \cos \left( \frac{u+v}{2} \right) \sin \left( \frac{u-v}{2} \right) &= \frac{1}{2} [\sin(u) - \sin(v)] \\ \cos \left( \frac{u+v}{2} \right) \cos \left( \frac{u-v}{2} \right) &= \frac{1}{2} [\cos(u) + \cos(v)] \\ \sin \left( \frac{u+v}{2} \right) \sin \left( \frac{u-v}{2} \right) &= \frac{1}{2} [\cos(v) - \cos(u)]\end{aligned}$$

Multiplying both sides by 2 and substituting  $u$  with  $\alpha$  and  $v$  with  $\beta$ , we arrive at the Sum-to-Product Formula, where we negate the last equation. ■

**Example 3.4.6** Express the sum  $\sin(8x) + \sin(2x)$  as a product of sines or cosines.

**Solution.** Using the formula we get

$$\begin{aligned}\sin(8x) + \sin(2x) &= 2 \sin \left( \frac{8x+2x}{2} \right) \cos \left( \frac{8x-2x}{2} \right) \\ &= 2 \sin \left( \frac{10x}{2} \right) \cos \left( \frac{6x}{2} \right) \\ &= 2 \sin(5x) \cos(3x)\end{aligned}$$

□

**Example 3.4.7** Express the difference  $\cos(3t) - \cos(5t)$  as a product of sines or cosines.

**Solution.** Using the formula we get

$$\begin{aligned}\cos(3t) - \cos(5t) &= -2 \sin\left(\frac{3t+5t}{2}\right) \sin\left(\frac{3t-5t}{2}\right) \\ &= -2 \sin\left(\frac{8t}{2}\right) \sin\left(\frac{-2t}{2}\right) \\ &= -2 \sin(4t) \sin(-t) \\ &= 2 \sin(4t) \sin(t)\end{aligned}$$

□

### 3.4.3 Exercises

**Exercise Group.** Express each product as a sum or difference of sine and cosine.

- |   |   |
|---|---|
| 1. $\sin(3x) \cos(5x)$<br><b>Answer.</b> $\frac{1}{2}[\sin(8x) + \sin(-2x)] = \frac{1}{2}[\sin(8x) - \sin(2x)]$                               | 2. $\sin(7t) \cos(-2t)$<br><b>Answer.</b> $\frac{1}{2}[\sin(5t) + \sin(9t)]$  |
| 3. $\cos(-3t) \sin(7t)$<br><b>Answer.</b> $\frac{1}{2}[\sin(4t) - \sin(-10t)] = \frac{1}{2}[\sin(4t) + \sin(10t)]$                            | 4. $\cos(9\theta) \sin(6\theta)$<br><b>Answer.</b> $\frac{1}{2}[\cos(15\theta) - \cos(3\theta)]$  |
| 5. $\cos(-4x) \cos(6x)$<br><b>Answer.</b> $\frac{1}{2}[\cos(2x) + \cos(-10x)] = \frac{1}{2}[\cos(2x) + \cos(10x)]$                            | 6. $\cos(2\theta) \cos(4\theta)$<br><b>Answer.</b> $\frac{1}{2}[\cos(6\theta) + \cos(-2\theta)] = \frac{1}{2}[\cos(6\theta) + \cos(2\theta)]$ |
| 7. $\sin(-\theta) \sin(8\theta)$<br><b>Answer.</b> $\frac{1}{2}[\cos(-9\theta) - \cos(7\theta)] = \frac{1}{2}[\cos(9\theta) - \cos(7\theta)]$ | 8. $\sin(6t) \sin(-3t)$<br><b>Answer.</b> $\frac{1}{2}[\cos(9t) - \cos(3t)]$  |

**Exercise Group.** Express each sum or difference as a product.

- |  |  |
|--|--|
| 9. $\sin(-2\theta) + \sin(-9\theta)$<br><b>Answer.</b> $2 \sin\left(-\frac{11\theta}{2}\right) \cos\left(\frac{7\theta}{2}\right) = -2 \sin\left(\frac{11\theta}{2}\right) \cos\left(\frac{7\theta}{2}\right)$ | 10. $\sin(5x) + \sin(7x)$<br><b>Answer.</b> $2 \sin(6x) \cos(x) = 2 \sin(6x) \cos(x)$  |
| 11. $\sin(4\theta) - \sin(-7\theta)$<br><b>Answer.</b> $2 \cos\left(\frac{11\theta}{2}\right) \sin\left(-\frac{3\theta}{2}\right) = -2 \cos\left(\frac{11\theta}{2}\right) \sin\left(\frac{3\theta}{2}\right)$ | 12. $\sin(3x) - \sin(-4x)$<br><b>Answer.</b> $2 \cos\left(\frac{7x}{2}\right) \sin\left(-\frac{x}{2}\right) = -2 \cos\left(\frac{7x}{2}\right) \sin\left(\frac{x}{2}\right)$ |
| 13. $\cos(8\theta) + \cos(-5\theta)$<br><b>Answer.</b> $2 \cos\left(\frac{3\theta}{2}\right) \cos\left(\frac{13\theta}{2}\right)$  | 14. $\cos(9t) + \cos(2t)$<br><b>Answer.</b> $2 \cos\left(\frac{11t}{2}\right) \cos\left(\frac{7t}{2}\right)$   |
| 15. $\cos(6\theta) - \cos(8\theta)$<br><b>Answer.</b> $-2 \sin(7\theta) \sin(-\theta) = 2 \sin(7\theta) \sin(\theta)$  | 16. $\cos(6t) - \cos(-3t)$<br><b>Answer.</b> $-2 \sin\left(\frac{9t}{2}\right) \sin\left(\frac{3t}{2}\right)$  |

**Exercise Group.** Find the exact value of each expression.

- |  |  |
|--|--|
| 17. $\sin(195^\circ) \cos(105^\circ)$<br><b>Answer.</b> $\frac{1}{2} \left(-\frac{\sqrt{3}}{2} + 1\right)$ | 18. $\cos(225^\circ) \cos(195^\circ)$<br><b>Answer.</b> $\frac{\sqrt{3}}{4} - \frac{1}{4}$ |
|--|--|



19.  $\sin(195^\circ) - \sin(75^\circ)$

**Answer.**  $-\frac{\sqrt{6}}{2}$

21.  $\sin(285^\circ) + \sin(195^\circ)$

**Answer.**  $-\frac{\sqrt{6}}{2}$

20.  $\cos(165^\circ) - \cos(105^\circ)$

**Answer.**  $-\frac{\sqrt{2}}{2}$

22.  $\cos(255^\circ) + \cos(15^\circ)$

**Answer.**  $\frac{\sqrt{2}}{2}$

**Exercise Group.** Verify the identity

23.  $\sin \theta + \sin(3\theta) = 4 \sin \theta \cos^2 \theta$

24.  $\cos(3\theta) + \cos \theta = 2(\cos^3 \theta - \sin^2 \theta \cos \theta)$

25.  $6 \cos(5\theta) \sin(6\theta) = 3 \sin(11\theta) + 3 \sin(\theta)$

26.  $\frac{\sin \theta + \sin(3\theta)}{2 \sin(2\theta)} = \cos \theta$

27.  $\frac{\cos \theta + \cos(3\theta)}{2 \cos(2\theta)} = \cos \theta$

28.  $\frac{\cos \theta - \cos(3\theta)}{\sin \theta + \sin(3\theta)} = \tan \theta$

## 3.5 Basic Trigonometric Equations

**Trigonometric equations** are equations involving trigonometric functions such as sine, cosine, and tangent. These equations seek to find specific values, known as **solutions**, that satisfy the equation.

Trigonometric functions are inherently periodic, meaning they repeat their values at regular intervals. As a result, trigonometric equations may have multiple solutions due to this periodic nature. In fact, some equations may have infinitely many solutions.

To address all possible solutions, we use a technique known as a **general solution**. This method involves initially identifying solutions within a single period of the trigonometric function. We then extend these solutions by adding integer multiples of the period of the trigonometric function.

In this section, we will explore various techniques for effectively solving trigonometric equations, including methods for finding general solutions.

### 3.5.1 Solving Equations with a Single Trigonometric Function

**Example 3.5.1** Solve the equation

$$\sin \theta = \frac{1}{2}.$$

**Solution.** To solve the equation  $\sin \theta = \frac{1}{2}$ , our initial instinct might lead us to take the inverse sine of both sides:


$$\sin^{-1}(\sin \theta) = \sin^{-1}\left(\frac{1}{2}\right)$$

resulting in

$$\theta = \frac{\pi}{6}.$$


While this is a valid solution, it's important to recognize that there are additional solutions to consider.

Since the sine function is positive in both Quadrant I and Quadrant II, we can find another solution in Quadrant II by using the reference angle  $\frac{\pi}{6}$  and the methods described in [Subsection 1.5.2](#). Therefore, the equivalent angle in Quadrant II is  $\theta = \pi - \frac{\pi}{6} = \frac{5\pi}{6}$ .



However, these two angles,  $\theta = \frac{\pi}{6}$  or  $\theta = \frac{5\pi}{6}$ , are not the only solutions. Recall that the sine function has a period of  $2\pi$ , meaning that adding or

subtracting any integer multiples of  $2\pi$  to these angles will also give you solutions. For example,  $\theta = \frac{\pi}{6} + 4\pi$  and  $\theta = \frac{5\pi}{6} - 10\pi$  are both solutions.



Thus, the general solution to  $\sin \theta = \frac{1}{2}$  can be expressed as:

$$\theta = \frac{\pi}{6} + 2k\pi \quad \text{or} \quad \theta = \frac{5\pi}{6} + 2k\pi,$$

where  $k$  is any integer. □

**Example 3.5.2** Solve the equation  $2 \cos \theta + \sqrt{2} = 0$ , list six solutions.

**Solution.** First we will isolate  $\cos \theta$ .

$$\begin{aligned} 2 \cos \theta + \sqrt{2} &= 0 \\ 2 \cos \theta &= -\sqrt{2} \\ \cos \theta &= -\frac{\sqrt{2}}{2}. \end{aligned}$$

Thus we have the general solution  $\theta = \frac{3\pi}{4} + 2k\pi$  or  $\theta = \frac{5\pi}{4} + 2k\pi$  for any integer  $k$ .

To get specific solutions, we select specific values of  $k$ :

$$\begin{aligned} k = -1 : \theta &= \frac{3\pi}{4} + 2(-1)\pi = -\frac{5\pi}{4}, & \theta &= \frac{5\pi}{4} + 2(-1)\pi = -\frac{3\pi}{4} \\ k = 0 : \theta &= \frac{3\pi}{4} + 2(0)\pi = \frac{3\pi}{4}, & \theta &= \frac{5\pi}{4} + 2(0)\pi = \frac{5\pi}{4} \\ k = 1 : \theta &= \frac{3\pi}{4} + 2(1)\pi = \frac{11\pi}{4}, & \theta &= \frac{5\pi}{4} + 2(1)\pi = \frac{13\pi}{4} \end{aligned}$$

□

### 3.5.2 Solving Trigonometric Equations with Square Terms

**Example 3.5.3** Solve  $\cos^2 \theta = \frac{1}{2}$  where  $0 \leq \theta < 2\pi$  on the interval  $0 \leq \theta < 2\pi$ .

**Solution.** We will first solve for  $\cos \theta$ . We begin by taking the square root of both sides of the equation and simplify:

$$\begin{aligned}\sqrt{\cos^2 \theta} &= \pm \sqrt{\frac{1}{2}} \\ \cos \theta \pm \frac{1}{\sqrt{2}} &= \pm \frac{\sqrt{2}}{2}\end{aligned}$$

From [Table 1.5.18](#), when  $\cos \theta = \frac{\sqrt{2}}{2}$ , we have  $\theta = \frac{\pi}{4}$  or  $\theta = \frac{7\pi}{4}$ ; and when  $\cos \theta = -\frac{\sqrt{2}}{2}$ , we have  $\theta = \frac{3\pi}{4}$  or  $\theta = \frac{5\pi}{4}$ .

Thus, our solutions are:

$$\theta = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}.$$

□

### 3.5.3 Solving Trigonometric Equations by Factoring

**Example 3.5.4** Solve  $\cos^2 x - 4 \cos x + 3 = 0$ .

**Solution.** To solve this equation, let's make a substitution to simplify it. We'll let  $y = \cos x$ , so the equation becomes  $y^2 - 4y + 3 = 0$ .

Factoring the quadratic equation, we obtain  $(y - 3)(y - 1) = 0$ . Setting each factor equal to zero, we find two potential solutions:  $y = 3$  or  $y = 1$ . We are not done because we need to solve for  $x$  and not  $y$ .

Substituting back  $\cos x$  for  $y$ , we find that  $\cos x = 3$  is not a valid solution, as the range of cosine is limited to  $[-1, 1]$ . However,  $\cos x = 1$  yields a solution of  $x = 0$  for one period.

Therefore, the general solution to the equation is:

$$x = 0 + 2k\pi = 2k\pi,$$

where  $k$  is any integer.

□

**Example 3.5.5** Solve  $2 \cos \theta \sin \theta + \sqrt{3} \cos \theta = 0$ .

**Solution.** We begin by factoring  $\cos \theta$ :

$$\begin{aligned}2 \cos \theta \sin \theta + \sqrt{3} \cos \theta &= 0 \\ \cos \theta(2 \sin \theta + \sqrt{3}) &= 0\end{aligned}$$

Thus we get two equations:  $\cos \theta = 0$  and  $2 \sin \theta + \sqrt{3} = 0$ .

From  $\cos \theta = 0$  we get  $\theta = \frac{\pi}{2}$  or  $\theta = \frac{3\pi}{2}$ .

From the second equation, we isolate  $\sin \theta$ :

$$\begin{aligned}2 \sin \theta + \sqrt{3} &= 0 \\ 2 \sin \theta &= -\sqrt{3} \\ \sin \theta &= -\frac{\sqrt{3}}{2}\end{aligned}$$

Thus we get  $\theta = \frac{4\pi}{3}$  or  $\theta = \frac{5\pi}{3}$ .

We get the general solutions by adding integer multiples of  $2\pi$  to get

$$\theta = \frac{\pi}{2} + 2k\pi, \quad \theta = \frac{3\pi}{2} + 2k\pi, \quad \theta = \frac{4\pi}{3} + 2k\pi, \quad \theta = \frac{5\pi}{3} + 2k\pi$$

where  $k$  is any integer. □

### 3.5.4 Solving a Trigonometric Equation with a Calculator

**Example 3.5.6** Use a calculator to solve  $3 \tan \theta = 2$  on the interval  $0 \leq \theta < 2\pi$ . Express your answer in radians, rounded to two decimals.

**Solution.** We begin by isolating  $\tan \theta$  :

$$\tan \theta = \frac{2}{3}$$

Next, we take the inverse tangent and use a calculator to obtain

$$\theta = \tan^{-1}\left(\frac{2}{3}\right) \approx 0.588002603548$$

Rounding to two decimals, we get  $\theta = 0.59$  radians, which is in Quadrant I since  $0 < 0.59 < \frac{\pi}{2}$ . Another Quadrant where  $\tan \theta = \frac{2}{3}$  is in Quadrant III. Using the methods in [Subsection 1.5.2](#) we get the other angle:  $\theta = 0.59 + \pi$ . □

### 3.5.5 Exercises

**Exercise Group.** Solve each equation on the interval  $0 \leq \theta < 2\pi$

1.  $\sin \theta = \frac{\sqrt{3}}{2}$

**Answer.**  $\frac{\pi}{3}, \frac{2\pi}{3}$

3.  $\cot \theta = -\sqrt{3}$

**Answer.**  $\frac{5\pi}{6}, \frac{11\pi}{6}$

5.  $\sec \theta = -2$

**Answer.**  $\frac{2\pi}{3}, \frac{4\pi}{3}$

7.  $\csc \theta + \sqrt{2} = 0$

**Answer.**  $\frac{5\pi}{4}, \frac{7\pi}{4}$

9.  $2 \tan \theta + 2\sqrt{3} = 0$

**Answer.**  $\frac{2\pi}{3}, \frac{5\pi}{3}$

11.  $2 \sin \theta + 1 = 0$

**Answer.**  $\frac{7\pi}{6}, \frac{11\pi}{6}$

2.  $\tan \theta = -1$

**Answer.**  $\frac{3\pi}{4}, \frac{7\pi}{4}$

4.  $\csc \theta = 2$

**Answer.**  $\frac{\pi}{6}, \frac{5\pi}{6}$

6.  $\cos \theta = \frac{\sqrt{2}}{2}$

**Answer.**  $\frac{\pi}{4}, \frac{7\pi}{4}$

8.  $6 \cos \theta + 1 = -2$

**Answer.**  $\frac{2\pi}{3}, \frac{4\pi}{3}$

10.  $2 \cot \theta + 8 = 6$

**Answer.**  $\frac{3\pi}{4}, \frac{7\pi}{4}$

12.  $\sec \theta - \sqrt{2} = 0$

**Answer.**  $\frac{\pi}{4}, \frac{7\pi}{4}$

**Exercise Group.** Solve each equation, giving the general formula for each solution. List six specific solutions.

13.  $\sin \theta = -\frac{\sqrt{2}}{2}$

**Answer.**  $\frac{5\pi}{4} + 2k\pi, \frac{7\pi}{4} + 2k\pi; -\frac{3\pi}{4}, -\frac{\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}, \frac{13\pi}{4}, \frac{15\pi}{4}$

15.  $\tan \theta = -1$

**Answer.**  $\frac{\pi}{4} + k\pi; -\frac{3\pi}{4}, \frac{\pi}{4}, \frac{5\pi}{4}, \frac{9\pi}{4}, \frac{13\pi}{4}, \frac{17\pi}{4}$

14.  $\cos \theta = \frac{1}{2}$

**Answer.**  $\frac{\pi}{3} + 2k\pi, \frac{5\pi}{3} + 2k\pi; -\frac{5\pi}{3}, -\frac{\pi}{3}, \frac{\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}, \frac{11\pi}{3}$

16.  $\cot \theta = -\sqrt{3}$

**Answer.**  $\frac{5\pi}{6} + k\pi, \frac{11\pi}{6} + k\pi; -\frac{\pi}{6}, \frac{5\pi}{6}, \frac{11\pi}{6}, \frac{17\pi}{6}, \frac{23\pi}{6}, \frac{29\pi}{6}$

**Exercise Group.** Solve each equation on the interval  $0 \leq \theta < 2\pi$ . Express your answer in radians, rounded to two decimals.

17.  $\sin \theta = -0.82$

Answer. 4.10, 5.32

19.  $\tan \theta = 2.5$

Answer. 1.19, 4.33

18.  $\cos \theta = 0.3$

Answer. 1.27, 5.02

20.  $\cot \theta = -5$

Answer. 2.94, 6.09

**Exercise Group.** Solve each equation on the interval  $0 \leq \theta < 2\pi$ .

21.  $4 \cos^2 \theta - 3 = 0$

Answer.  $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ 

23.  $4 \tan^2 \theta - 1 = 0$

Answer.  $\frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$ 

25.  $\sec^2 \theta - 4 = 0$

Answer.  $\frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$ 

27.  $(2 \cos \theta - 1)(\csc \theta + 2) = 0$

Answer.  $\frac{\pi}{3}, \frac{7\pi}{6}, \frac{5\pi}{3}, \frac{11\pi}{6}$ 

29.  $4 \sin^2 \theta - 2 \sin \theta = 0$

Answer.  $0, \frac{\pi}{6}, \frac{5\pi}{6}, \pi$ 

31.  $\sin^3 \theta - \sin \theta = 0$

Solution.  $0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}$ 

33.  $2 \sin^2 \theta - 7 \sin \theta + 3 = 0$

Answer.  $\frac{\pi}{6}, \frac{5\pi}{6}$ 

35.  $\cos^2 \theta + 2 \cos \theta + 1 = 0$

Answer.  $\pi$ 

37.  $2 \sin^2 \theta - 3 \sin \theta - 2 = 0$

Answer.  $\frac{7\pi}{6}, \frac{11\pi}{6}$ 

22.  $3 \csc^2 \theta - 4 = 0$

Answer.  $\frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$ 

24.  $2 \cot^2 \theta - 6 = 0$

Answer.  $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ 

26.  $2 \cos^2 \theta - 1 = 0$

Answer.  $\frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$ 

28.  $(\tan \theta - \sqrt{3})(\cot \theta + 1) = 0$

Answer.  $\frac{\pi}{3}, \frac{3\pi}{4}, \frac{4\pi}{3}, \frac{7\pi}{4}$ 

30.  $2 \cos^2 \theta - \sqrt{3} \cos \theta = 0$

Answer.  $\frac{\pi}{6}, \frac{\pi}{2}, \frac{3\pi}{2}, \frac{11\pi}{6}$ 

32.  $2 \cos^2 \theta + \cos \theta - 1 = 0$

Answer.  $\frac{\pi}{3}, \pi, \frac{5\pi}{3}$ 

34.  $3 \tan^3 \theta - \tan \theta = 0$

Answer.  $0, \frac{\pi}{6}, \frac{5\pi}{6}, \pi, \frac{7\pi}{6}, \frac{11\pi}{6}$ 

36.  $\csc^5 \theta - 4 \csc \theta = 0$

Answer.  $\frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$ 

38.  $2 \cos^2 \theta - 3 \cos \theta + 1 = 0$

Answer.  $0, \frac{\pi}{3}, \frac{5\pi}{3}$

## 3.6 Trigonometric Equations - Advanced Techniques

In this section, we will solve trigonometric equations using various trigonometric identities, equations involving multiple angles, and graphical methods.

### 3.6.1 Solving a Trigonometric Equation by Using Fundamental Identities

**Example 3.6.1** Solve  $2 \sin^2 x + 3 \cos x - 3 = 0$ .

**Solution.** To solve the equation  $2 \sin^2 x + 3 \cos x - 3 = 0$ , we want to first write it in terms of only cosine or only sine. By the Pythagorean Identity, we substitute  $\sin^2 x = 1 - \cos^2 x$ , resulting in:

$$\begin{aligned} 2 \sin^2 x + 3 \cos x - 3 &= 0 \\ 2(1 - \cos^2 x) + 3 \cos x - 3 &= 0 \\ 2 - 2 \cos^2 x + 3 \cos x - 3 &= 0 \\ -2 \cos^2 x + 3 \cos x - 1 &= 0 \\ 2 \cos^2 x - 3 \cos x + 1 &= 0 \\ (2 \cos x - 1)(\cos x - 1) &= 0 \end{aligned}$$

Thus  $\cos x = \frac{1}{2}$  or  $\cos x = 1$ .

Solving for  $x$  in the first equation gives  $x = \frac{\pi}{3}$  or  $\frac{5\pi}{3}$ , while the second equation gives  $x = 0$  for one period. Finding all solutions, we arrive at  $x = \frac{\pi}{3} + 2k\pi$ ,  $\frac{5\pi}{3} + 2k\pi$ , or  $x = 0 + 2k\pi = 2k\pi$  for any integer  $k$ .  $\square$

**Remark 3.6.2** If you are having trouble factoring trigonometric functions, try substituting a simpler term and then factor. For example, consider the expression  $2 \cos^2 x - 3 \cos x + 1$  in the previous example. If factoring this expression directly is not clear, you can use the substitution  $A = \cos x$ . This transforms the expression into  $2A^2 - 3A + 1$ , which may be easier to factorize. Once factored, you can substitute  $\cos x$  back in for  $A$  to obtain the final solution.

**Example 3.6.3** Solve the equation  $\sin(2\theta) + \cos \theta = 0$  on the interval  $0 \leq \theta < 2\pi$ .

**Solution.** Notice the first term has  $2\theta$  so we will begin by using the double-angle formula:

$$\begin{aligned} \sin(2\theta) + \cos \theta &= 0 \\ 2 \sin \theta \cos \theta + \cos \theta &= 0 \end{aligned}$$

Then we factor our  $\cos \theta$  from all the terms:

$$\cos \theta(2 \sin \theta + 1) = 0.$$

Thus we get

$$\cos \theta = 0 \quad \text{or} \quad 2 \sin \theta + 1 = 0.$$

When  $\cos \theta = 0$  we get  $\theta = \frac{\pi}{2}, \frac{3\pi}{2}$ .

When  $2 \sin \theta + 1 = 0$ , it is equivalent to when  $\sin \theta = -\frac{1}{2}$ , thus we get  $\theta = \frac{7\pi}{6}, \frac{11\pi}{6}$ .

Combining these, our solutions are

$$\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{6}, \frac{11\pi}{6}.$$

□

### 3.6.2 Solving a Trigonometric Equation with Multiples of an Angle

**Example 3.6.4** Solve the equation  $2 \cos(2\theta) - \sqrt{3} = 0$  on the interval  $0 \leq \theta < 2\pi$ .

**Solution.** First we will isolate  $\cos(2\theta)$ . We begin by rearranging the equation:

$$\begin{aligned} 2 \cos(2\theta) - \sqrt{3} &= 0 \\ 2 \cos(2\theta) &= \sqrt{3} \\ \cos(2\theta) &= \frac{\sqrt{3}}{2} \end{aligned}$$

Since  $\cos(2\theta)$  equals  $\frac{\sqrt{3}}{2}$  for angles  $\frac{\pi}{6}$  and  $\frac{11\pi}{6}$ , the general solutions for  $2\theta$  are

$$2\theta = \frac{\pi}{6} + 2k\pi \quad \text{or} \quad 2\theta = \frac{11\pi}{6} + 2k\pi$$

for some integer  $k$ .

Note that we have only solved for  $2\theta$ . Dividing both sides by 2 to find  $\theta$ , we obtain:

$$\theta = \frac{\pi}{12} + k\pi \quad \text{or} \quad \theta = \frac{11\pi}{12} + k\pi.$$

Our restriction  $0 \leq \theta < 2\pi$  gives us the following solutions:

$$\frac{\pi}{12}, \frac{11\pi}{12}, \frac{13\pi}{12}, \frac{23\pi}{12}.$$

□

**Remark 3.6.5** In this example, our angle was a double-angle:  $2\theta$ . When dealing with multiple-angle trigonometric functions, such as  $\cos(2\theta)$ , it's essential to understand their graphical behavior. According to [Definition 2.1.29](#), the graph of  $\cos(2\theta)$  undergoes a horizontal compression by a factor of 2, and its period is now  $\frac{2\pi}{2} = \pi$ . Since we are asked to find solutions on the interval  $0 \leq \theta < 2\pi$ , we will need to consider two periods. Thus, we have four solutions (2 solutions for each period):

$$\frac{\pi}{12}, \frac{11\pi}{12}, \frac{13\pi}{12}, \frac{23\pi}{12}.$$

In general, if our trigonometric function has an angle  $k\theta$ , for some number  $k$ , we will need to consider the effect of this multiple angle on the period and number of solutions, ensuring that we adjust our solutions accordingly to cover all possible solutions within the given interval.

**Example 3.6.6** Solve the equation  $3 \tan \frac{\theta}{2} - \sqrt{3} = 0$  on the interval  $0 \leq \theta < 2\pi$ .

**Solution.** We begin by isolating  $\tan \frac{\theta}{2}$ :

$$\begin{aligned} 3 \tan \frac{\theta}{2} - \sqrt{3} &= 0 \\ 3 \tan \frac{\theta}{2} &= \sqrt{3} \\ \tan \frac{\theta}{2} &= \frac{\sqrt{3}}{3} \end{aligned}$$



Since  $\tan \frac{\theta}{2}$  equals  $\frac{\sqrt{3}}{3}$  for angles  $\frac{\pi}{6}$  and  $\frac{7\pi}{6}$ , the general solutions for  $\frac{\theta}{2}$  are

$$\frac{\theta}{2} = \frac{\pi}{6} + k\pi \quad \text{and} \quad \frac{\theta}{2} = \frac{7\pi}{6} + k\pi$$

for some integer  $k$ .

Solving for  $\theta$ , we multiply both sides by 2 to obtain:

$$\theta = \frac{\pi}{3} + 2k\pi \quad \text{and} \quad \theta = \frac{7\pi}{3} + 2k\pi.$$

Considering our restriction  $0 \leq \theta < 2\pi$ , we only have one solution:

$$\theta = \frac{\pi}{3}.$$

□

**Remark 3.6.7** In this example, our angle was  $\frac{\theta}{2}$ , which stretches the graph of the tangent function. When we have an angle of the form  $\frac{\theta}{2}$ , it effectively stretches the period of the tangent function by a factor of 2. Thus, for the given interval, we only have half a period to consider instead of the full period.

### 3.6.3 Solving a Trigonometric Equation with a Graphing Utility


Sometimes we will encounter equations where an exact solution is not possible. However, we may be able to get an approximation to the solution by graphing the equation.

**Example 3.6.8** Use a graphing utility to find the solutions to the equation  $\sin x + \cos x = \frac{1}{2}x$ . Express your answers in radians, rounded to two decimals.

**Solution.** To find the solution to  $\sin x + \cos x = \frac{1}{2}x$ , we graph the left-hand side and the right-hand side of the equation and identify their intersections. Let  $y_1$  represent the curve for the left-hand side and  $y_2$  represent the curve for the right-hand side:

$$y_1 = \sin x + \cos x, \quad y_2 = \frac{1}{2}x.$$

Use a graphing utility to plot  $y_1$  and  $y_2$ .



**Figure 3.6.9** Plotting  $y_1 = \sin x + \cos x$  and  $y_2 = \frac{1}{2}x$ , corresponding to the left-hand and right-hand sides of the equation, respectively.

Next, you may need to zoom in or out to better visualize the behavior of the curves. To find their intersection points, calculators often have a **TRACE** or **INTERSECT** button or command. In [Desmos Graphing Calculator](#)<sup>1</sup> you can click on either curve, and the points of intersection will be highlighted. Hovering your cursor over the intersection will display the coordinates of that point.

The equation  $\sin x + \cos x = \frac{1}{2}x$  has three solutions, which correspond to the points of intersection between the curves  $y_1 = \sin x + \cos x$  and  $y_2 = \frac{1}{2}x$ . The  $x$ -values of these intersections are:

$$x = -2.68, -1.24, 1.71.$$

□

### 3.6.4 Exercises

**Exercise Group.** Solve each equation on the interval  $0 \leq \theta < 2\pi$ .

- |  |  |
|--|--|
| 1. $\sin^2 \theta - \cos^2 \theta = 0$<br><b>Answer.</b> $\frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$ | 2. $\cos^2 \theta - \sin^2 \theta = 1 + \sin \theta$<br><b>Answer.</b> $0, \pi, \frac{7\pi}{6}, \frac{11\pi}{6}$ |
| 3. $3 \cot^2 \theta - 4 \csc \theta = 1$<br><b>Answer.</b> $\frac{\pi}{6}, \frac{5\pi}{6}$                               | 4. $\sin^2 \theta = 5 \cos \theta + 5$<br><b>Answer.</b> $\pi$   |
| 5. $\cos^2 \theta = 3 - 3 \sin \theta$<br><b>Answer.</b> $\frac{\pi}{2}$   | 6. $2 \sin^2 \theta = 3 \cos \theta + 3$<br><b>Answer.</b> $\frac{2\pi}{3}, \pi, \frac{4\pi}{3}$                 |

**Trigonometric Equations Involving Multiples of an Angle.** Solve the given trigonometric equation on the interval  $0 \leq \theta < 2\pi$ .

- |   |   |
|---|---|
| 7. $\cot 2\theta = -\sqrt{3}$<br><b>Answer.</b> $\frac{5\pi}{12}, \frac{11\pi}{12}, \frac{17\pi}{12}, \frac{23\pi}{12}$ | 8. $\sin 4\theta = \frac{\sqrt{3}}{2}$<br><b>Answer.</b> $\frac{\pi}{12}, \frac{\pi}{6}, \frac{7\pi}{12}, \frac{2\pi}{3}, \frac{13\pi}{12}, \frac{7\pi}{6}, \frac{19\pi}{12}, \frac{5\pi}{3}$ |
| 9. $\sqrt{2} \cos 2\theta = 1$<br><b>Answer.</b> $\frac{\pi}{8}, \frac{7\pi}{8}, \frac{9\pi}{8}, \frac{15\pi}{8}$       | 10. $\csc 3\theta = 2$<br><b>Answer.</b> $\frac{\pi}{18}, \frac{5\pi}{18}, \frac{13\pi}{18}, \frac{17\pi}{18}, \frac{25\pi}{18}, \frac{29\pi}{18}$  |
| 11. $\sec \frac{3\theta}{2} = -\sqrt{2}$<br><b>Answer.</b> $\frac{\pi}{2}, \frac{5\pi}{6}, \frac{11\pi}{6}$             | 12. $\cos \frac{\theta}{2} - 1 = 0$<br><b>Answer.</b> $0$   |

**Trigonometric Equations Involving Addition or Subtraction Formula.** Use the Addition and Subtraction Formulas to solve each equation on the interval  $0 \leq \theta < 2\pi$ .

13.  $\sin \theta \cos 2\theta + \cos \theta \sin 2\theta = \frac{\sqrt{3}}{2}$   
**Answer.**  $\frac{\pi}{9}, \frac{2\pi}{9}, \frac{7\pi}{9}, \frac{8\pi}{9}, \frac{13\pi}{9}, \frac{14\pi}{9}$
14.  $\sin 3\theta \cos 2\theta - \cos 3\theta \sin 2\theta = -\frac{1}{2}$   
**Answer.**  $\frac{7\pi}{6}, \frac{11\pi}{6}$
15.  $\cos 3\theta \cos \theta + \sin 3\theta \sin \theta = -\frac{\sqrt{2}}{2}$   
**Answer.**  $\frac{3\pi}{8}, \frac{5\pi}{8}, \frac{11\pi}{8}, \frac{13\pi}{8}$
16.  $\cos \theta \cos 3\theta - \sin \theta \sin 3\theta = 1$   
**Answer.**  $0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}$

<sup>1</sup>DesmosGraphingCalculator

**Trigonometric Equations Involving Double-Angle or Half-Angle Formula.** Use the Double-Angle and Half-Angle Formulas to solve each equation on the interval  $0 \leq \theta < 2\pi$ .

17.  $\sin 2\theta = \cos \theta$

**Answer.**  $\frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}, \frac{3\pi}{2}$

19.  $\cos 2\theta + \cos \theta - 2 = 0$

**Answer.**  $0$

21.  $\cos 2\theta + 2 = 2 \sin^2 \theta$

**Answer.**  $\frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$

18.  $\cos 2\theta = \cos \theta$

**Answer.**  $0, \frac{2\pi}{3}, \frac{4\pi}{3}$

20.  $\tan 2\theta = -2 \sin \theta$

**Answer.**  $0, \frac{\pi}{3}, \pi, \frac{5\pi}{3}$

22.  $\tan \frac{\theta}{2} = \sin \theta$

**Answer.**  $0, \frac{\pi}{2}, \frac{3\pi}{2}$

**Trigonometric Equations Involving Sum-to-Product Formula.** Use the Sum-to-Product Formulas to solve each equation on the interval  $0 \leq \theta < 2\pi$ .

23.  $\sin 3\theta + \sin \theta = 0$

**Answer.**  $0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}$

25.  $\cos 4\theta - \cos 2\theta = 0$

**Solution.**  $0, \frac{\pi}{3}, \frac{2\pi}{3}, \pi, \frac{4\pi}{3}, \frac{5\pi}{3}$

24.  $\sin 6\theta - \sin 2\theta = \cos 4\theta$

**Answer.**  $\frac{\pi}{12}, \frac{\pi}{8}, \frac{3\pi}{8}, \frac{5\pi}{12}, \frac{5\pi}{8}, \frac{7\pi}{8}, \frac{13\pi}{12}, \frac{9\pi}{8}, \frac{11\pi}{8}, \frac{17\pi}{12}, \frac{13\pi}{8}, \frac{15\pi}{8}$

26.  $\cos 3\theta + \cos \theta = 0$

**Solution.**  $\frac{\pi}{4}, \frac{\pi}{2}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{3\pi}{2}, \frac{7\pi}{4}$

**Exercise Group.** Use a graphing utility to solve each equation. Express your solutions in radians, rounded to two decimals.

27.  $\sin(2x) = 4 \cos x + x$

**Answer.**  $x = -1.34, 2.63, 3.90$

29.  $x^2 = \cos x$

**Answer.**  $x = -0.82, 0.82$

28.  $\sin x - x = \cos x$

**Answer.**  $x = -1.26$

30.  $x^3 + 2x^2 = \cos(2x)$

**Answer.**  $x = -2.11, -0.56, 0.48$

# Glossary (Math)

**amplitude.definition**

**period.definition**

# Glossary (Canoe and Wayfinding)

**heading**.definition

**latitude sailing**.definition

# Index

- addition and subtraction formulas,
  - 184
  - for cosine, 184
  - for sine, 186
  - for tangent, 188
- amplitude, 97
- angle, 12
- angular speed, 23
- arc length, 20
- arc minute, 13
- arc second, 13
- arccosine function, 159
- arcsine function, 159
- arctangent function, 159
- area
  - of a sector, 21
- axial tilt, 88
- azimuth, 17
  
- backwash, 62
  
- central angle, 16
- cofunction identities, 50
- cofunctions, 49, 50
- complementary angles, 49
- corkscrew, 155
- cosine
  - domain, 90
  - range, 90
- coterminal angles, 15
  
- dead reckoning, 163
- declination, 25
  - solar, 30, 88
- degree, 12
- double-angle formulas, 197
  
- equinox
  - fall, 88
  - spring, 88
  
- even and odd trigonometric properties, 81
  
- fundamental trigonometric identities, 178
  
- general solution, 214
  
- half-angle formulas
  - for cosine, 201
  - for sine, 201
  - for tangent, 201, 203
- heading, 16
- horizontal shift, 100
  
- initial side, 12
- inverse cosine, 159
- inverse sine, 159
- inverse tangent, 159
  
- Kūkuluokalani, 4
  
- latitude, 12, 28
- linear speed, 23
- longshore drift, 62
  
- Makali‘i, 87
- measure, 12
- midline, 97
- minute, 13
- moon phases, 30
  
- nautical mile, 29
- negative angle, 14
  
- odd-even identities, 81
  
- paafu, 1
- period, 98
  - wave, 110
- period functions, 78
- periodic properties, 78

- phase shift, 100
- pitch, 155
- positive angle, 14
- product-to-sum formulas, 209
- Pythagorean identities, 80
  
- quadrantal angle, 14
- quadrants, 14
- quotient identities, 48
  
- radian measure
  - of angles, 17
- ray, 12
- reciprocal identities, 47
- reducing powers formulas, 200
- reference angle, 67
- reference course, 47
- reflection
  - about  $x$ -axis, 95
  - about  $y$ -axis, 96
- roll, 155
  
- second, 13
- sine
  - domain, 90
  - range, 90
- sinus curves, 93
- sinusoidal graphs, 93
- solar declination, 30, 88
- solstice
  - summer, 89
  - winter, 88
- speed
  - angular, 23
  - linear, 23
- standard position, 14
- Star Compass
  - Cook Island, 9
  - Hawaiian, 4
  - Māori, 9
  - paafu, 1
  - Sāmoan, 9
- stick chart, 7
- sum-to-product formula, 211
- sums of sines and cosines, 190
- swash, 62
  
- terminal side, 12
- triangles
  - solving, 52
- trigonometric equations, 214
- trigonometric functions, 34
  - circle of radius  $r$ , 43
  - of angles, 35
  - of real numbers, 34
  - ratios, 47
  - special, 49
  - special values, 40
  
- unit circle, 33
  
- vertical asymptote, 115
- vertical compression, 97
- vertical shift, 94
- vertical stretch, 97
  
- wave height, 110
- wave period, 110